

## Research Article



# Growth performance and body composition of the African catfish (*Clarias gariepinus*) juveniles fed diets supplemented with *Afrostryrax lepidophyllus* fruit powder

Yemdjie Mane D.D.<sup>1\*</sup>; Nyadjeu P.<sup>1</sup>; Ndjuissi Tamko A.N.<sup>1</sup>; Agwah Ebile D.<sup>2</sup>; Kuate Simo F.<sup>1</sup>; Efole Ewouken T.<sup>1</sup>; Pouomogne V.<sup>1</sup>; Tomedi Eyang M.<sup>1</sup>

Received: April 2022

Accepted: August 2022

### Abstract

To minimize the dependence of pisciculture on imported feed and improve the production of fish through the valorization of local feeds, this study was conducted to evaluate the effects of dietary inclusion of *Afrostryrax lepidophyllus* fruit powder on the growth and chemical composition of *Clarias gariepinus* juveniles. The juveniles weighing  $13 \pm 3.1$  g were divided into four treatments in triplicates and fed with diets containing 0 (T<sub>0</sub> or control), 2 (T<sub>1</sub>), 4 (T<sub>2</sub>), and 6 (T<sub>3</sub>) g/kg *A. lepidophyllus* fruit powder for 63 days. The feeding rate was 5% of their body weight. The results showed that adding different levels of *A. lepidophyllus* fruit powder to the diets improved growth, feed utilization, and carcass quality attributes. The African catfish fed with T<sub>3</sub> diet recorded higher final weight ( $43.44 \pm 0.23$  g), weight gain (WG,  $31.43 \pm 0.41$  g), feed conversion ratio (FCR,  $0.91 \pm 0.05$ ), and protein efficiency ratio (PER,  $2.67 \pm 0.14$ ) than those recorded in the control group (40.20 g final weight,  $27.20 \pm 0.52$  g WG,  $1.1 \pm 0.05$  FCR, and  $2.24 \pm 0.10$  PER). The same trend was observed in the body macro-nutrient retention ( $32.33 \pm 0.58$  % crude protein,  $25.33 \pm 0.58$  % lipid, and  $23.33 \pm 0.58$  % ash) compared to the control group ( $20.67 \pm 0.58$  % crude protein,  $16.00 \pm 0.00$  % lipid, and  $17.33 \pm 0.58$  % ash). Finally, this study clearly showed that dietary inclusion of *A. lepidophyllus* fruit powder at 6 g/kg improves growth performance, feed nutrient utilization, and body composition in juvenile *C. gariepinus*.

**Keywords:** *Afrostryrax lepidophyllus*, Carcass quality, *Clarias gariepinus*, Zootechnical performance

1- Department of Aquaculture, Institute of Fisheries and Aquatic Sciences of Yabassi, University of Douala, P.O. Box: 7236-Douala, Cameroon.

2- Department of Animal Science, Laboratory of Animal Nutrition, University of Dschang, P.O. Box: 96-Dschang, Cameroon.

\*Corresponding author's Email: dyemdjie5@yahoo.fr

## Introduction

The amount of wild-caught fish has not increased since 1980; aquaculture has mainly contributed to the supply of fish for human consumption in the world (FAO, 2016). In Africa, fish production is hampered by the unavailability of aqua-feed which is mostly imported and very expensive (Atangana *et al.*, 2019). However, in Africa and particularly in Cameroon, there is a wide range of products and agricultural by-products rich in nutrients and the addition of some feed additives or dietary supplements can keep fish healthy and enhance growth rate (Sayed *et al.*, 2011; Adegbesan *et al.*, 2019). Research shows that herbs and spices (phytobiotics) are used as supplements or feed additives in animal diets (Muneendra *et al.*, 2014; Mohsen *et al.*, 2018).

Phytobiotics contain active secondary metabolites belonging to the classes of isoprene derivatives, flavonoids, and phenols compounds which have been suggested to act as antioxidants (Frankič *et al.*, 2009; Sumczynski *et al.*, 2015). Beneficial effects of phytobiotics in farm animals result from feed intake improvement and secretion of digestive enzymes, immune stimulation, anti-bacterial, anthelmintic, antiviral, anti-inflammatory, and antioxidant properties (Muneendra *et al.*, 2014). Previous studies revealed that ginger (*Zingiber officinale*), garlic (*Allium sativum*), and scent leaf (*Ocimum gratissimum*) contain active substances such as alkaloids, tannins, flavonoids, saponins, and phenols compounds (Muhammad *et al.*, 2009; Abdou *et al.*,

2010). Those compounds are known to improve animal performances through their anti-oxidative and anti-microbial actions and improve feed palatability (Muneendra *et al.*, 2014). An example of a plant possessing secondary metabolites is *A. lepidophyllus* fruit (Moukette *et al.*, 2015; Namkona *et al.*, 2017; Sokamte *et al.*, 2018).

*A. lepidophyllus* is a plant of the Huaceae family and is commonly found in Equatorial and Tropical Africa (Cronquist, 1981; Moukette *et al.*, 2015; Namkona *et al.*, 2017). This plant is used in Congo as an antiseptic and in traditional medicine in treating gastroenteric diseases (Bouquet, 1969). In Cameroon, the seeds of this plant are traditionally used as a spice. Moreover, pharmacological studies showed that the extracts of *A. lepidophyllus* seed possess very interesting properties. Ngono (1999) demonstrated the antifungal activity and identified Afrostyraxthioside A, Afrostyraxthioside B, and Afrostyraxthioside C. Agbor *et al.* (2005) and Boufack *et al.* (2021) showed that the seed extracts possess antioxidant properties. In addition, Toumnou *et al.* (2012) reported the insecticidal activity of the extract from the seeds of *A. lepidophyllus*. Qualitative analysis of *A. lepidophyllus* fruit showed the presence of total polyphenols, flavonoids, tannins, and anthocyanins (Namkona *et al.*, 2017). These compounds possess significant antimicrobial activity (Sokamte *et al.*, 2018) and antioxidant properties (Oben *et al.*, 2010; Fogang *et al.*, 2014; Sokamte *et al.*, 2018) with

many other attributes including digestive enzyme stimulation, lipid metabolism, and microbial load modulation (Muneendra *et al.*, 2014).

In the view of the chemical composition of *A. lepidophyllus* fruit, its ability to balance the gut microbial load and stimulate digestive enzymes (Frankič *et al.*, 2009; Oben *et al.*, 2010), this phyto-additive can positively affect the growth performance of *C. gariepinus*. This study aims to determine the effects of *A. lepidophyllus* fruit powder on growth performance, feed utilization, and body composition of the African catfish (*Clarias gariepinus*) juveniles.

## Materials and methods

### *Site of study*

This study was conducted from April to May 2021 in the research and applied farm in fishery and aquaculture of Kanhé-Moyo (FRAPAIK) in Baham, the western region of Cameroon. This farm is located at 5°19' North and 10°24' East and an altitude of 1728 m above sea level. The annual average temperature is 23°C. Rainfall ranges between 1500-2000 mm per annum, over 9 months of the rainy season (March to November).

### *Experimental design*

360 *C. gariepinus* juveniles were distributed following a completely randomized design to four treatments of 90 fish each, the average weight was 13±3.10g. Each experimental treatment had three (03) experimental units (hapas) of 30 juveniles each placed in a non-fertilized pond of 80 m<sup>2</sup>.

### *Origin of A. lepidophyllus*

*A. lepidophyllus* fruit were bought from a local market then ground, sieved and the powder was incorporated at different levels in locally produced feed.

### *Animal material*

Three hundred and sixty *C. gariepinus* (360) juveniles, were purchased from a renowned fish farm in Douala. These juveniles were acclimatized for two weeks in 03 hapas of 1 m<sup>3</sup> each, they were fed during the first three days with a commercial imported feed (Gouessant, France; 2 mm) with 46% crude protein, 10 % crude fat, 2.83 % crude fiber, 8.1 % crude ash, 1 % calcium, 1.4% phosphorus, and 0.4 sodium as declared by the company. From the fourth day, 25% of their ration was substituted with local feed without *A. lepidophyllus*. From the seventh to the tenth day respectively 50 and 75 % of the imported feed was substituted. On the thirteenth and fourteenth days, the animals were entirely fed with local feed (100%). At the end of this acclimatization period, ten *C. gariepinus* juveniles were randomly sampled for the determination of the carcass quality characteristics.

### *Housing and equipment*

After acclimatization, juveniles were randomly transferred to 12 experimental hapas of 0.64 m<sup>2</sup>. The different feed rations were distributed manually at a frequency of three times per day (6:00 a.m., 12:00, and 5:00 p.m.); at a rate of 5% of the fish biomass. In order to monitor growth and adjust the quantities of feed distributed, using a landing net,

control fishing was done after 21 days and during cool hours of the day (6:00 a.m.). The different individual weights and lengths were respectively measured using a 1g precision electronic scale balance and an ichthyometer. A 50cm long Secchi disc, JBL Test Kits (JBL GmbH & Co. KG–Dieselstraße 3 - 67141 Neuhofen – Germany).

Maximum-minimum thermometer and pH meter were respectively used daily to measure water transparency, dissolved oxygen, temperature, and pH. The values of the physico-chemical parameters of water recorded during the experiment are resumed in Table 1.

**Table 1: Water quality parameters (Mean  $\pm$  SD) during the experimental period (63 days).**

Parameters	Experimental period (days)			
	0-21	21-42	42-63	Optimum values
T(°C)	22.55 $\pm$ 1.41	20.63 $\pm$ 1.60	22.2 $\pm$ 1.66	20-30
pH	7.03 $\pm$ 0.08	6.81 $\pm$ 0.49	7.07 $\pm$ 0,73	6.5-8
D.O(ppm)	5.85 $\pm$ 0.02	7,89 $\pm$ 0,55	7.63 $\pm$ 0,59	7.5-9.1
NO <sub>3</sub> <sup>-</sup> (mg/L)	30 $\pm$ 0	23 $\pm$ 2.64	25 $\pm$ 3	<50
Transparency (cm)	45.23 $\pm$ 6.62	43.04 $\pm$ 7.20	42.17 $\pm$ 7.04	<60

Temperature (T°C); Hydrogen potential pH; dissolved oxygen (D.O); nitrate (NO<sub>3</sub><sup>-</sup>)

#### Experimental diet

In addition to the locally prepared diet (T<sub>0</sub> or control), three isoprotein, isolipidic and isoenergetic diets including T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub> were made by incorporating 2, 4, and 6g/kg *A. lepidophyllus* fruit powder, respectively. The basal diet (T<sub>0</sub>) ingredients and proximate composition are summarized in Table 2.

#### Preparation of diets

The raw ingredient was finely ground in a mill. *A. lepidophyllus* previously crushed was incorporated in the mixtures while respecting the different proportions specified in Table 2. In each treatment, cold water (20% per treatment) was mixed, then stirred delicately before extrusion in a DGP60-C brand extruding machine, with a capacity of 120-150kg/hour, at a 105°C temperature equipped with a die of

2.5mm. The extrudates were sun-dried for 20 hours then placed in plastic bags, labeled, and kept away from humidity until use.

**Table 2: Composition of the basal diet.**

Ingredients	% in dry matter (DM)
Fishmeal	27
Soybean meal	15
Peanut meal	20
Cotton meal	8
Wheat bran	8
Maize meal	16
*Premix 5%	5
Palm oil	1
Total	100
<b>Chemical composition</b>	
Protein (% DM)	41.136 $\pm$ 3.68
Energy (kcal/kg DM)	2759.24 $\pm$ 6.04
Lipid (% DM)	11.18 $\pm$ 0.5
Ash (% DM)	11.101 $\pm$ 0.95
Moisture (% DM)	5.5 $\pm$ 0.57
Fiber (% DM)	6.31 $\pm$ 0.63
Dry matter (%)	94.5 $\pm$ 0.57

\*Premix 5%: Crude protein =40%; Lysine =3.30; Methionine= 2.40; Calcium= 8; Phosphorus= 2.05; Metabolized Energy = 2078 kcal/kg.

The formulated diet samples (10g) was analyzed following the procedures of AOAC (1990). Moisture content was determined by drying the sample in an oven at 105°C overnight. Crude protein was analyzed in a KJELTEC SYSTEM 1002 Distilling Unit made in Belgium following the Kjeldahl method while lipid content was determined by extraction with hexane using the Soxhlet method. The ash content was determined by combustion in a muffle furnace at 500°C for 6 h.

#### *Growth, survival rate, and feed utilization parameters*

Growth performance, survival rate, feed utilization, and nutrient retention were assessed per treatment by determining the weight gain (WG), specific growth rate (SGR), survival rate (SR), condition factor (K), feed intake (FI), feed conversion ratio (FCR), feed efficiency

ratio (FER), protein efficiency ratio (PER) and nutrient retention (NR). The calculations were done using the following formulae:

$$WG (g) = W_f - W_i$$

$$SGR (\%/day) = \frac{\ln W_f - \ln W_i}{T} \times 100$$

Where,  $W_f$  = final weight;  $W_i$  = initial weight;  $T$  = experimental period (days);

$$SR (\%) = \frac{\text{final number of fish}}{\text{initial number of fish}} \times 100$$

$$K = \frac{\text{Weight}}{\text{Length}^3} \times 100$$

$$FI (g/fish)$$

$$= \frac{\text{total dry feed distributed}}{\text{number of fish}}$$

$$FCR = \frac{\text{feed intake}}{\text{Fish weight gained}}$$

$$FER = \frac{\text{Fish weight gained}}{\text{feed intake}}$$

$$PER = \frac{\text{Fish weight gained}}{\text{Protein consumed}}$$

$$\text{Protein consumed} = \frac{\text{Total feed consumed} \times \text{Crude protein in feed}}{100}$$

$$NR (\%) = \frac{\text{Final carcass composition} - \text{Initial carcass composition}}{\text{Amount of nutrient fed}} \times 100$$

#### *Statistical analysis*

Results on growth performance, survival rate, feed utilization, and nutrient retention obtained from each replicate were used for statistical analysis. Data collected were submitted to a one-way Analysis of Variance test by the General Linear Model's procedure of Statistical Package for Social Science (SPSS 20.0) software. Where there was a significant difference between treatments, their means were separated using Duncan's multiple range test and probability


values less than 0.05 were considered significant.

## **Results**

### *Growth performance*

Independent of the incorporation level of *A. lepidophyllus* fruit powder in different levels, Figure 1 shows an inflection in the body weight and weight gain curves on the 42<sup>nd</sup> day of the experiment. After this point, the curves have a linear and increasing evolution until the end of the experiment. Moreover, the evolution of

weight and weight gain of *C. gariepinus* juveniles is proportional to the increasing incorporation level of *A. lepidophyllus* powder in the diets. Thus, after the 21<sup>st</sup> day, the curve of *C. gariepinus* juveniles fed diet supplemented with 6 g/kg *A. lepidophyllus* (T<sub>3</sub>) remained above all the other curves. At the end of the trial, the body weight (44.43±0.23 g) was significantly higher ( $p<0.05$ ) by 9.52%,



**Figure 1:** Body weight and weight gain of *Clarias gariepinus* juvenile fed different levels of *Afrostryrax lepidophyllus*. T<sub>0</sub>= control ration, T<sub>1</sub>= T<sub>0</sub>+2g/kg *A. lepidophyllus*, T<sub>2</sub>= T<sub>0</sub>+4 g/kg *A. lepidophyllus*, T<sub>3</sub>= T<sub>0</sub>+ 6 g/kg *A. lepidophyllus*.

The relationship between fish growth and dietary inclusion level of *A. lepidophyllus* powder is illustrated by the linear regression curve (Figure 2) below. *C. gariepinus* juvenile growth rate increased with increasing inclusion level of *A. lepidophyllus* in the diet ( $R^2=0.9671$ ).

The specific growth rate of *C. gariepinus* juveniles fed with different experimental diets for 63 days is illustrated in Figure 3. It is noted that irrespective of the treatments, the growth was faster during the first week of feeding and gradually decreased with the

4.82%, and 2.99% compared to those fed respectively T<sub>0</sub> (40.20±0.30 g), T<sub>1</sub> (42.29±0.14 g) and T<sub>2</sub> (43.10±0.2 g). On the other hand, weight gain (31.43±0.41 g) was significantly higher ( $p<0.05$ ) by 13.46%, 6.78%, and 4.23% compared to those fed with T<sub>0</sub> (27.20±0.52 g), T<sub>1</sub> (29.30±0.24 g), and T<sub>2</sub> (30.10±0.34 g), respectively.

increasing experimental period. At the end of the experimental period, fish fed 6 g/kg *A. lepidophyllus* powder (T<sub>3</sub>) had a specific growth rate of 1.95±0.06% g/d, significantly higher ( $p<0.05$ ) compared to the other treatments T<sub>0</sub> (1.79±0.02%g/d), T<sub>1</sub> (1.87 ± 0.009% g/d), T<sub>2</sub> (1.90 ± 0.01% g/d).

#### *Survival Rate and Feed Nutrient Utilization*

Table 3 resumes the survival rate and feed nutrient utilization of *C. gariepinus* juveniles at the end of the feeding period. It is noticed that, though no

significant difference was observed amongst treatments. Mortalities occurred in two treatments T<sub>1</sub> and T<sub>2</sub> with a survival rate of  $93.33 \pm 0.00\%$ . Determination of the feed nutrients utilization parameters (feed conversion ratio, feed efficiency ratio, and protein efficiency ratio) indicated that the fish fed diets supplemented with *A.*

*lepidophyllus* had better feed utilization characteristics ( $p < 0.05$ ) compared to the control diet. On the other hand, we notice a significant increase in the value of condition factor K concerning the dietary inclusion level of *A. lepidophyllus*.




Figure 2: Regression curve between *Clarias gariepinus* growth and dietary inclusion levels of *Afrostyrax lepidophyllus* fruit powder.




Figure 3: Specific growth rate of *Clarias gariepinus* juveniles fed different levels of *Afrostyrax lepidophyllus* fruit powder. T<sub>0</sub>=control ration, T<sub>1</sub>=T<sub>0</sub>+2g/kg *A. lepidophyllus*, T<sub>2</sub>=T<sub>0</sub>+4 g/kg *A. lepidophyllus*, T<sub>3</sub>=T<sub>0</sub>+ 6 g/kg *A. lepidophyllus*. Means on the same rearing period carrying different superscripts were significantly different at  $p < 0.05$ .

### Chemical (nutrient) Composition and Nutrient Retention of *C. gariepinus*

Nutrient composition of *C. gariepinus* juveniles produced at the end of the experimental period (63 days) is recorded in Table 4. As compared to fish nutrient composition at the beginning of the study, it is noted that apart from the moisture content that dropped significantly ( $p<0.05$ ) in the juveniles

fed diets supplemented with *A. lepidophyllus* powder, other macro-elements like the ash, protein, and lipid levels were significantly ( $p<0.05$ ) enhanced by increasing the incorporation levels of *A. lepidophyllus* powder in the diets. With the exception of the energy level that was significantly ( $p<0.05$ ) higher in the fish fed T<sub>2</sub> diet.

**Table 3: Survival rate and feed utilization of *Clarias gariepinus* fed diets supplemented with *Afrostryax lepidophyllus* fruit powder at the end of the feeding period.**

Parameters	Treatments				p- value
	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	
Ni	90	90	90	90	-
Nf	90	88	88	90	-
SR	100.00 ± 0.00	93.333 ± 0.00	93.33 ± 0.00	100.00 ± 0.00	-
FI (g/fish)	39 ± 0.00	38.409 ± 0.00	38.116 ± 0.00	39 ± 0.00	-
Wi (g)	13.02 ± 0.01 <sup>a</sup>	13.01 ± 0.02 <sup>a</sup>	13.02 ± 0.01 <sup>a</sup>	13.01 ± 0.1 <sup>a</sup>	0.977
Wf (g)	40.20 ± 0.30 <sup>d</sup>	42.29 ± 0.14 <sup>c</sup>	43.10 ± 0.2 <sup>b</sup>	44.43 ± 0.23 <sup>a</sup>	0.000
WG (g)	27.20 ± 0.52 <sup>d</sup>	29.29 ± 0.24 <sup>c</sup>	30.10 ± 0.34 <sup>b</sup>	31.43 ± 0.41 <sup>a</sup>	0.000
Lf (cm)	17.38 ± 0.02 <sup>d</sup>	17.59 ± 0.05 <sup>c</sup>	17.94 ± 0.02 <sup>a</sup>	17.72 ± 0.02 <sup>b</sup>	0.000
K	0.76 ± 0.01 <sup>bc</sup>	0.78 ± 0.01 <sup>ab</sup>	0.75 ± 0.004 <sup>c</sup>	0.80 ± 0.002 <sup>a</sup>	0.006
FCR	1.1 ± 0.05 <sup>a</sup>	0.94 ± 0.02 <sup>b</sup>	0.90 ± 0.05 <sup>b</sup>	0.91 ± 0.05 <sup>b</sup>	0.004
FER	0.92±0.04 <sup>b</sup>	1.06±0.03 <sup>a</sup>	1.11±0.06 <sup>a</sup>	1.11±0.06 <sup>a</sup>	0.007
PER	2.24±0.10 <sup>b</sup>	2.59±0.07 <sup>a</sup>	2.69±0.15 <sup>a</sup>	2.67±0.14 <sup>a</sup>	0.007

T<sub>0</sub>= control ration, T<sub>1</sub>=T<sub>0</sub>+2g/kg *A. lepidophyllus*, T<sub>2</sub>=T<sub>0</sub>+4 g/kg *A. lepidophyllus*, T<sub>3</sub>=T<sub>0</sub>+ 6 g/kg *A. lepidophyllus*.

Values are mean±standard deviation of three replicates of 30 fish each. Mean within the row with different superscripts are significantly different from other at  $p<0.05$ . Ni, initial number of fish; Nf, final number of fish; FI, feed intake; Lf, final length of fish; K, condition factor; FCR, feed conversion ratio; FER, feed efficiency ratio; PER, protein efficiency ratio.

**Table 4: Proximate composition (% or kJ/g WW) of carcass and nutrient retention in *Clarias gariepinus* juveniles fed different inclusion levels of *Afrostryax lepidophyllus* fruit powder.**

Parameters	Initial	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	p- value
	Chemical composition (% or KJ/g WW)					
Ash	2.47±0.38 <sup>d</sup>	2.66±0.34 <sup>c</sup>	2.76±0.24 <sup>b</sup>	2.76±0.26 <sup>b</sup>	2.99±0.84 <sup>a</sup>	0.000
Crude protein	11.34±0.22 <sup>e</sup>	11.96±1.09 <sup>d</sup>	14.32±0.40 <sup>c</sup>	14.72±0.18 <sup>b</sup>	15.08±1.05 <sup>a</sup>	0.000
Lipid	2.36±0.28 <sup>e</sup>	2.50±0.32 <sup>d</sup>	2.55±0.45 <sup>c</sup>	3.10±0.26 <sup>b</sup>	3.17±0.34 <sup>a</sup>	0.000
Energy	7.31±0.58 <sup>e</sup>	7.43±0.40 <sup>d</sup>	8.84±0.35 <sup>c</sup>	9.22±0.46 <sup>a</sup>	9.16±0.22 <sup>b</sup>	0.000
Moisture	81±0.01 <sup>a</sup>	81±0.01 <sup>a</sup>	77±0.01 <sup>b</sup>	77±0.01 <sup>b</sup>	77±0.01 <sup>b</sup>	0.000
Nutrient retention (% dry feed)						
Ash		17.33±0.58 <sup>c</sup>	19.67±0.58 <sup>b</sup>	20.00±0.00 <sup>b</sup>	23.33±0.58 <sup>a</sup>	0.000
Crude protein		20.67±0.58 <sup>d</sup>	28.67±0.58 <sup>c</sup>	30.33±0.58 <sup>b</sup>	32.33±0.58 <sup>a</sup>	0.000
Lipid		16.00±0.00 <sup>d</sup>	18.00±0.00 <sup>c</sup>	23.67±0.58 <sup>b</sup>	25.33±0.58 <sup>a</sup>	0.000
Energy		19.00±0.00 <sup>d</sup>	26.00±0.00 <sup>c</sup>	28.00±0.00 <sup>b</sup>	29.00±0.00 <sup>a</sup>	0.000
Dry matter		14.00±0.00 <sup>c</sup>	20.00±0.00 <sup>b</sup>	20.33±0.58 <sup>b</sup>	21.00±0.00 <sup>a</sup>	0.000

T<sub>0</sub>= control ration, T<sub>1</sub>= T<sub>0</sub>+2g/kg *A. lepidophyllus*, T<sub>2</sub>= T<sub>0</sub>+4 g/kg *A. lepidophyllus*, T<sub>3</sub>= T<sub>0</sub>+ 6 g/kg *A. lepidophyllus*; WW, wet weight.

Mean within the same row with different superscripts are significantly different at  $p<0.05$ . p= probability



Results on nutrient retention of *C. gariiepinus* carcass (Table 4) after 63 days of feeding revealed that incorporating *A. lepidophyllus* powder in fish diet affects protein, lipid, ash, energy, and dry matter contents. The fish fed T<sub>3</sub> diet had protein (32.33±0.58%), lipid (25.33±0.58%), ash (23.33±0.58%), energy (29.00±0.00%), and dry matter (21.00±0.00%) contents retention significantly higher compared to those fed the other experimental diets. Figure 4 illustrates the linear regression between nutrient retention in the body of

fish and the dietary inclusion level of *A. lepidophyllus* powder. Regardless of the nutrient, there is a relationship between nutrient retention and dietary inclusion level of *A. lepidophyllus* powder. This is illustrated by the regression coefficient which indicates the relationship between lipid retention ( $R^2=0.95$ ) and the inclusion level of *A. lepidophyllus* powder followed by ash retention ( $R^2=0.91$ ) then by protein retention ( $R^2=0.86$ ) and finally energy retention ( $R^2=0.84$ ).




Figure 4: Regression curves between nutrient retention of *Clarias gariiepinus* juveniles and dietary inclusion levels of *Afrostryax lepidophyllus* fruit powder.

## Discussion

Water quality parameters had a great influence on fish survival rate as well as

on their growth. Moreover, the recorded physicochemical parameters such as temperature, dissolved oxygen, pH, and

nitrate ( $\text{NO}_3^-$ ) were within the recommended range for freshwater fish breeding. According to the above observations, the mortalities observed in the fish fed with 2 g/kg ( $T_1$ ) and 4 g/kg ( $T_2$ ) dietary inclusion of *A. lepidophyllus* fruit powder could be probably attributed to stress-induced during the fish rearing. The application of phytochemical agents (herbal components) in aquaculture has greatly increased as they have been reported to prevent diseases and reduce the use of hazardous antibiotics (Sakai, 1999). Results from this study showed positive effects of dietary inclusion *A. lepidophyllus* on growth performance and feed nutrients utilization. Incorporating *A. lepidophyllus* fruit powder in *C. gariepinus* feed markedly enhanced body weight and body weight gain compared to those fed the control diet. The highest response was observed in fish fed with the highest level of phyto-additive. These results agree with Soosean *et al.* (2010) and Abbasi *et al.* (2017), who showed that the highest final weight and weight gain was recorded when the African catfish was fed the highest level of *Garcinia mangostana* and *Zingiber officinale* powder in common carp, *Cyprinus carpio* diets. This result can be explained by the presence of anti-oxidative and antimicrobial substances such as flavonoids and phenolic compounds (Fogang *et al.*, 2014; Namkona *et al.*, 2017; Sokamte *et al.*, 2018) which are known to maintain gut equilibrium and improve growth performance of animal (Frankič *et al.*, 2009). The highest

improvement in body weight was recorded with *C. gariepinus* fed 6 g/kg *A. lepidophyllus* powder. This could be attributed to the antimicrobial properties of their specific active compounds and their impact on gut function. Increasing feed inclusion levels of *A. lepidophyllus* increased flavonoids and phenolic compounds, which are known to improve animal performances by changing the intestinal ecosystem of the animal through their antimicrobial action (Odoemelam *et al.*, 2013). These compounds act by forming complexes with many proteins, causing the destruction of the bacterial membranes (Frankič *et al.*, 2009). The changes in the intestinal ecosystem due to their antimicrobial action could lead to greater availability of some nutrients for the host and consequently improve body weight gain and feed utilization. This is in agreement with Frankič *et al.* (2009), who noticed the growth-promoting effect of most herbs and spices that act by killing parasites that hinder the digestibility and growth performance of animals. On the other hand, the improvement in body weight of *C. gariepinus* juveniles obtained with high a level of incorporation of *A. lepidophyllus* could also be attributed to the potent antioxidant properties of their major components as reported by Fogang *et al.* (2014) and Moukette *et al.* (2015).

Some phyto-additives have been reported to improve specific growth rate (Dada and Ikurowo, 2009; Nyadjeu *et al.*, 2021). In this study, the fish fed diet containing 6 g/kg *A. lepidophyllus*

powder had the specific growth rate value significantly higher than all the other treatments. This result contradicts those obtained using dietary *Zingiber officinale* extract in beluga juvenile (*Huso huso*) diet (Vahedi *et al.*, 2017). The conflicting results of the present study with those of Vahedi *et al.* (2017) could be attributed to the different species, feeding programs, and farming conditions. On the other hand, it was noted that regardless of the treatment, juvenile growth was faster during the first week of feeding and gradually decreased with the increase in the experimental period. This was comparable to that obtained by Gnikpo *et al.* (2014) who reported that the specific growth rate decreases while the muscle mass increases.

The improvement in feed nutrient utilization in *C. gariepinus* juveniles induced by *A. lepidophyllus* inclusion compared to those fed the control diet could have emanated from the presence of bioactive compounds contained in the phytobiotic. These results agree with the findings of Kana *et al.* (2017) where the incorporation of *A. lepidophyllus* fruits in broilers diet enhanced body weight and body weight gain and tended to reduce feed conversion ratio compared to birds fed the control diet. The improvement in growth and feed nutrient utilization in animals could be attributed to gallic acid which was the most abundant phenolic acid present in *A. lepidophyllus* fruit (Sokamte *et al.*, 2018). According to Kang Yang *et al.* (2020), gallic acid influences the gut microbiome and modulates the immune

response to maintain intestinal health. The improvement of intestinal health reduced the animal's exposure to microbial toxins and other undesired microbial metabolites such as ammonia and biogenic amines (Windisch *et al.*, 2008). Thereby, animals are relieved from immune defense stress during critical situations and there is increased availability of essential nutrients for absorption. This is in accordance with the results from this study where there was an increase in nutrient retention (protein, lipid, ash, energy, and dry matter) in the fish fed diets supplemented with *A. lepidophyllus* compared to those in the control diet. Moreover, the fish fed diet containing 6 g/kg *A. lepidophyllus* powder registered the highest nutrient retention of these macro-elements. This result indicates that increasing *A. lepidophyllus* level in *C. gariepinus* diet increases the presence of bioactive compounds present in the phyto-additive such as flavonoids and phenolics compounds (Sokamte *et al.*, 2018). These secondary metabolites are known to maintain gut equilibrium (Frankič *et al.*, 2009) of the microbiome that plays a crucial role in the maintenance of physiological homeostasis within the gastrointestinal tract (Kang Yang *et al.*, 2020). Hence, they help the animals to grow better within the framework of their genetic potential.

The results of the present study revealed that supplementing *C. gariepinus* diet with *A. lepidophyllus* improved growth performance, feed nutrient utilization, and nutrient

retention in the African catfish juveniles. The highest effect was obtained with the highest level of inclusion. These observed beneficial effects have been attributed to the bioactive contents of *A. lepidophyllus* than to their nutritional properties. The present findings provided a ray of hope for dietary supplements that are widely and intensively used in aquaculture.

### References

- Abbasi Ghadikolaei, H., Kamali, A., Soltani, M. and Sharifian, M., 2017.** Effects of *Zingiber officinale* powder on growth parameters, survival rate and biochemical composition of body in juvenile common carp (*Cyprinus carpio*). *Iranian Journal of Fisheries Sciences*, 16(1), 67-85. DOI:10.22092/IJFS.2018.114641
- Abdou Bouda, A., Njintang, Y.N., Scher, J. and Mbofung, C.M.F., 2010.** Phenolic compounds and radical scavenging potential of twenty Cameroonian spices. *Agriculture and Biology Journal of North America*, 1, 213-224.
- Adegbesan, S.I., Obasa, S.O., Akintokun, A.K. and Abdulraheem, I., 2019.** Effects of Dietary Supplementation of *Zingiber officinale* Root-Powder on Growth, Nutrient Utilization and Intestinal Microbes of African Mud Catfish (*Clarias gariepinus*) Fingerlings. *Journal of Aquaculture & Fisheries*, 3, 16. DOI:10.24966/AAF-5523/100016
- Agbor, G.A., Oben, J.E., Ngogang, J. Y., Xinxing, C. and Vinson, J.A., 2005.** Antioxidant Capacity of Some Herbs/Spices from Cameroon: A Comparative Study of Two Methods. *Journal of Agricultural Food Chemistry*, 53, 819-6824. DOI:10.1021/jf050445c
- AOAC (Association of Official Analytical Chemist), 1990.** Official methods of analysis association of official analytical chemists. 15<sup>th</sup> ed. Association of Analytical Chemists, Washington, DC.1298 P.
- Atangana, K.J.A., Ducarme, C. and Micha, J.C., 2019.** La pisciculture au Cameroun : bilan et perspectives. *International Journal of Biology Chemistry and Sciences*, 13(2), 1140-1161. DOI:10.4314/ijbcs.v13i2.44
- Boufack, J. L., Guiekep, N. A.J., Vemo, B. N., Fonou, T.L., Tsambou, M.M.A., Menkem, B., Ngaleu, D.C.C., Yidjieu, N.F. and Kenfack, A., 2021.** Effects of ethanol extract of *Afrostryrax lepidophyllus* seeds on reproductive parameters and oxidative stress markers in male guinea pigs (*Cavia porcellus*) exposed to paraquat. *International Journal of Veterinary Sciences and Animal Husbandry*, 6(4), 26-33. DOI:10.22271/veterinary.2021.v6.i4 a.363
- Bouquet, A., 1969.** Féticheurs et médecines traditionnelles du Congo (Brazzaville), O.R.S.T.O.M, Paris, pp. 234-235.
- Cronquist, A., 1981.** An Integrated System of Classification of Flowering Plants; Columbia University Press: New York, pp. 396-397.

- Dada, A.A. and Ikuerowo, M., 2009.** Effects of ethanolic extracts of *Garcinia kola* seeds on growth and haematology of catfish (*Clarias gariepinus*) broodstock. *African Journal of Agricultural Research*, 4(4), 344-347.
- FAO (Food and Agriculture Organization of the United Nations), 2016.** Food and Agricultural Organization of United Nations. Contributing to Food Security and Nutrition for All. The State of World Fisheries and Aquaculture. Rome. 200 P.
- Fogang, H.P.D., Maggi, F., Tapondjou, L.A., Womeni, H.M., Papa, F. and Quassinti, L., 2014.** In vitro Biological Activities of Seed Essential Oils from the Cameroonian Spices *Afrostryrax lepidophyllus* Mildbr. and *Scorodophloeus zenkeri* Harms Rich in Sulfur -Containing Compounds. *Chemistry & Biodiversity*, 11(1), 161-169.
- Frankič, T., Voljc, M., Salobir, J. and Rezar, V., 2009.** Use of herbs, spices and their extracts in animal nutrition. *Acta argiculturae Slovenica*, 94(2), 95-102.
- Gnikpo, A., Adandedjan, D., d'Almeida A., Houessou, A.A., Fagbemi, M., Soude, E., Osse, O. J., Houndonougbo, F., Kayode, P., Chrysostome, C., Codjia, J. C. and Chikou, A., 2014.** Effet du probiotique « kpete-kpete » sur la survie et la croissance des larves *clarias gariepinus* (poisson chat, clariidae) élevées en bacs hors sol. *Cahiers du CBRST* N° 5 juin 2014, Bibliothèque Nationale du Bénin, 03 BP : 1665, Cotonou, Rép du Bénin, pp. 27-43
- Kana, J.R., Mube, K.H., Ngouana, T.R., Yangoue, A., Komguep, R., Tsafong, F. and Tegua, A., 2017.** Growth Performance and Serum Biochemical Profile of Broiler Chickens Fed on Diets Supplemented with *Afrostryrax lepidophyllus* Fruit and Bark as Alternative to Antibiotic Growth Promoters. *Journal of Veterinary Medicine and Research*, 4(6), 1095.
- Kang, Y., Limeng, Z., Pinfeng, L., Zaili, X., Fan, Z., Daniel, S., Zhongquan, X., Chengquan, T., Jinping, D., Yulong, Y., and Baichuan, D., 2020.** Impact of Gallic Acid on Gut Health: Focus on the Gut Microbiome, Immune Response, and Mechanisms of Action. *Journal Frontiers in Immunology*, 11, 580208.  
DOI:10.3389/fimmu.2020.580208
- Mohsen Mohammadi Gheisar and In Ho Kim, 2018.** Phytochemicals in poultry and swine nutrition— a review. *Italian Journal of Animal Sciences*, 17(1), 92-99,  
DOI:10.1080/1828051X.2017.1350120
- Moukette, B.M., Constant, A.P., Prosper, C.N.B., Vicky, J.A.M., Eustace, B. and Jeanne, Y.N., 2015.** *Afrostryrax lepidophyllus* extracts exhibit in vitro free radical scavenging, antioxidant potential and protective properties against liver enzymes ion mediated oxidative damage. *BMC Research Notes*, 8,

- 344, DOI:10.1186/s13104-015-1304-8
- Muhammad, J., Fazil-Raziq, D., Abdul, H., Rifatullah, K. and Ijaz, A., 2009.** Effects of aqueous extract of plant mixture on carcass quality of broiler chicks. *Journal of Agricultural Biology Sciences*, 4, 37-40.
- Muneendra, K., Vinod, K., Debashis, R., Raju, K., and Shalini, V., 2014.** Application of Herbal Feed Additives in Animal Nutrition - A Review. *International Journal of Livestock Research*, 4, 1-8. DOI:10.5455/ijlr.20141205105218
- Namkona, A.F., Bolevane, O.S.F., Moustapha, F., Worowounga, X., Ngaissona, P. and Koane, J.N., 2017.** Biological activities and phytochemical analysis of extracts *Afrostryax lepidophyllus* Mildbr. seeds. *The Journal of Phytopharmacology*, 6(2), 102-106.
- Ngono, N.A., 1999.** Contribution à l'étude des propriétés antifongiques et analyse phytochimique de cinq plantes médicinales camerounaises. Ph.D. Thèse, Université de Reims Champagne-Ardenne, France. 198 P.
- Nyadjeu, P., Yemdjie, D.D.M., Ndjussi, N.A.T., Nguenang, G.N., Dedou, N.Y.C. and Tabi-Tomedi, M.E., 2021.** Effect of *Zingiber officinale* and *Allium sativum* Powders as Natural Feed Additives Promoting Growth, Feed Utilization and Whole-Body Composition in *Clarias gariepinus* Fry. *Food and Nutrition Sciences*, 12, 526-543. DOI:10.4236/fns.2021126040
- Oben, J., Etoundi, C. B., Kuate, D. and Ngondi, J. L., 2010.** Anti-amylase, anti-lipase and antioxidant effects of aqueous extracts of some Cameroonian spices. *Journal Natural Products*, 3, 165-171.
- Odoemelam, V.U., Nwaogu, K.O., Ukachukwu, S.N., Esonu, B.O., Okoli, I.C. and Etuk, E.B., 2013.** Carcass and Organoleptic assessment of Broiler fed *Ocimum gratissimum* Supplemented diets. Proceedings of the 38th Conferences of Nigeria Society of Animal Production. 17-20th March 2013, Rivers state University of Sciences and Technology, Port Harcourt, Nigeria.
- Sakai, M., 1999.** Current research status of fish immunostimulants. *Aquaculture*, 172, 63-92.
- Sayed, S.H., Zakaria, A., Mohamed, G.A. and Mohamed, K.K., 2011.** Use of Probiotics as growth promoter, anti-bacterial and their effects on the physiological parameters and immune response of *Oreochromis niloticus* Lin. fingerlings. *Journal of the Arabian Aquaculture Society*, 6, 201-222.
- Sokamte, T.A., Mbougoung, P.D., Ntsamo, B.T.M., Noumo, N.T. and Tatsadjieu, N.L., 2018.** Antioxidant and antimicrobial activities of two edibles spices from Cameroon and quantification of their major phenolic compounds. *International Food Research Journal*, 25(6), 2352-2361.
- Soosean, C., Marimuthu, K., Sudhakaran, S. and Xavier, R., 2010.** Effects of mangosteen

- (*Garcinia mangostana L.*) extracts as a feed additive on growth and haematological parameters of African catfish (*Clarias gariepinus*) fingerlings. *European Review of Medicine Pharmacy Sciences*, 14, 605-611.
- Sumczynski, D., Bubelova, Z., Sneyd, J., Erb-Weber, S. and Mlcek, J., 2015.** Total phenolics, flavonoids, antioxidant activity, crude fibre and digestibility in non-traditional wheat flakes and muesli. *Food Chemistry*, 174, 319–325. DOI : 10.1016/j.foodchem.2014.11.065
- Toumno, A.L., Seck, D., Namkossere, S., Cisse, N., Kandioura, N. and Sembene, M., 2012.** Utilisation des plantes indigènes à effet insecticide pour la protection des denrées stockées contre des insectes ravageurs à Boukoko (Centrafrique). *International Journal of Biology Chemistry and Sciences*, 6(3), 1040-1050. DOI:10.4314/ijbcs.v6i3.11
- Vahedi, A.H., Hasanpour, M., Akrami, R. and Chitsaz, H., 2017.** Effect of dietary supplementation with ginger (*Zingiber officinale*) extract on growth, biochemical and hemato-immunological parameters in juvenile beluga (*Huso huso*). *Iranian Journal of Aquatic Animal Health*, 3(1), 26-46. DOI:10.18869/acadpub.ijaah.3.1.26
- Windisch, W., Schedle, K., Plitzner, C. and Kroismayr, A., 2008.** Use of phytogenic products as feed additives for swine and poultry. *Journal of Animal Sciences*, 86, 140-148. DOI:10.2527/jas.2007-0459
- Zidni, I., Iskandar, Buwono, I.D. and Mahargyani, B.P., 2019.** Water Quality in the Cultivation of Cat fish (*Clarias gariepinus*) and Nile Tilapia (*Oreochromis niloticus*) in the Aquaponic Bio floc System. *Asian Journal of Fisheries and Aquatic Research*, 4, 1-6. DOI:10.9734/AJFAR/2019/v4i230048