

# Prevalence of Intestinal Protozoan Parasites among Children Attending the Hevi Pediatric Hospital in Duhok Province, Kurdistan Region, Iraq

Ismael, SS<sup>1\*</sup>, Abdullah, BH<sup>1</sup>, Sadiq, AJ<sup>1</sup>, Ajaj, JS<sup>1</sup>, Ali, NS<sup>1</sup>, Omer, DM<sup>1</sup>, Nori, NY<sup>1</sup>

1. Department of Medical Laboratory Sciences, College of Health Sciences, University of Duhok, Iraq

**How to cite this article:** Ismael SS, Abdullah BH, Sadiq AJ, Ajaj JS, Ali NS, Omer DM, Nori NY. Prevalence of Intestinal Protozoan Parasites among Children Attending the Hevi Pediatric Hospital in Duhok Province, Kurdistan Region, Iraq. *Archives of Razi Institute Journal*. 2024;79(3):507-512. DOI: 10.32592/ARI.2024.79.3.507



Copyright © 2023 by



Razi Vaccine & Serum Research Institute

## ABSTRACT

The present study aimed to determine the prevalence rate of intestinal protozoan infection among children in Duhok Province, North of Iraq, from October 2022 to May 2023. This cross-sectional study was conducted on 740 children attending Hevi Pediatric Hospital, Duhok Province, Iraq. Firstly, all collected stool samples were examined directly by preparing a wet stool smear for the detection of the vegetative stage (Trophozoite) of both *Entamoeba histolytica* and *Giardia lamblia*. Thereafter, all stool samples were preserved at -22°C until further use for identifying cysts of protozoa and oocysts of *Cryptosporidium parvum* by the Telman concentration stool method. The children were within the age range of 1-14 years. Out of 740 stool samples collected, 205 (27.7%) cases were positive for intestinal protozoan infections as follows: 160 (78.0%) children tested positive for *E. histolytica*, 35 (17.1%) were positive for *C. parvum*, and only 10 (4.9%) were positive for *G. lamblia*. Nonetheless, the results indicated a statistically significant difference between the prevalence of *E. histolytica*, *C. parvum*, and *G. lamblia* infections across different age groups ( $P < 0.05$ ). The highest infection rates of intestinal protozoa, namely *E. histolytica*, *C. parvum*, and *G. lamblia*, were reported in the age ranges of 5-9 years (53.75%, 25.0%, and 60.0%), 1-4 years (28.75%, 20.0%, and 30.0%), and 10-14 years (17.55%, 8.6%, and 10.0%), respectively. In this study, the prevalence of amoebiasis and cryptosporidiosis was reported to be higher than in other studies performed in Iraq, while the prevalence of giardiasis was reported at a lower rate than the rate reported previously in Iraq, especially in Duhok. These infections continue to pose a challenging public health issue, necessitating the implementation of stronger and more effective preventative measures.

### Article Info:

Received: 30 July 2023

Accepted: 30 September 2023

Published: 30 June 2024

### Corresponding Author's E-Mail:

shameeran.ismael@uod.ac

**Keywords:** Children, Duhok Province, Intestinal Protozoa, Prevalence

## 1. Introduction

Gastrointestinal infections and diarrhea are widespread in developing countries, including various tropical regions. Infections of the gastrointestinal tract can be caused by viruses, bacteria, protozoa, helminths, and occasionally fungi (1). The World Health Organization reported that diarrhea affects more individuals than any other disease, even in high-income countries (2). Poor quality drinking water is one of the factors that have been linked to an increased prevalence of diarrheal diseases (3). Intestinal protozoan infections are generally common across the globe. They represent a major cause of diarrheal diseases, especially among children in developing countries (1). *Entamoeba histolytica*, *G. lamblia*, and *C. parvum* are the most frequently identified protozoan parasites in stool samples. *Cryptosporidium parvum*, first described in humans in 1976, is now recognized as an important cause of diarrhea in young children and immunocompromised adults (4, 5). The transmission dynamics of enteric protozoa vary considerably among geographic areas (1). Understanding the significance of different sources of infection is needed to set priorities for controlling measures. The present study aimed to investigate protozoan infections in children with diarrhea and its relationship with protozoan contamination of drinking water and other possible risk factors in Duhok Province.

## 2. Materials and Methods

This cross-sectional study was conducted on 740 children referred to Hevi Pediatrics Hospital, Duhok Province, Iraq, from October 2022 to May 2023.

### 2.1. Sample Collection and Processing

About 4 g of fresh stool samples were collected from all children and put in a plastic cube. Each stool cup was labeled with the main information of the child, including name, age, and gender. All diarrheic samples were examined directly under the microscope for the detection of the trophozoite stage of *E. histolytica* and *G. lamblia*. Following that, all stool samples were preserved at  $-22^{\circ}\text{C}$  until used for the identification of cysts of *E. histolytica* and *G. lamblia*, and oocysts of *C. parvaum*, by direct wet stool smear and concentration stool methods.

#### 2.1.1. Direct stool smear

A direct stool smear examination was performed using the methods previously mentioned (6). A small amount of stool sample was mixed with a few drops of normal saline, put on a cover slide, and examined directly under the microscope first at low power and then moved to other powers. From each sample, at least three slides were prepared and examined.

#### 2.1.2. Telmman stool concentration technique

Firstly, all frozen stool samples were left at room temperature to be thawed; thereafter, about 1 g of stool

sample was mixed well with 5 ml of acetic acid and set for a few minutes. It was then filtered through a fine sieve, and the filtered fluid was pureed into a clean test tube. Following that, the same volume of ether was added to the test tube, covered with a cover, and centrifuged at 2000 rpm for 3 min. After centrifugation, the supernatant was discarded and a drop of sediment was put on the slide and mixed with a few drops of normal saline, covered, and examined under the microscope for the detection of cysts. For the detection of the oocyst of *C. parvum*, a thin stool smear was prepared from the sediment and dried at room temperature. Subsequently, the slide was put in Methyl alcohol for 4 min for fixation, and the smear was then stained with Modified Ziehl-Neelsen Stain.

### 2.3. Statistical analysis

GraphPad Prism software (version 8.0) was used to evaluate the data gathered for the current investigation. The Chi-square test was used to examine the relationship of infections with age and gender.

## 3. Results

The children were in the age range of 1-14 years. Out of the 740 stool samples collected, 205 (27.7%) were positive for intestinal protozoan infections as follows: 160 (78.0%) children tested positive for *E. histolytica*, 35 (17.1%) were positive for *C. parvum*, and only 10 (4.9%) were positive for *G. lamblia*, as presented in table 1 and displayed in figures 1, 2, and 3. Nonetheless, the results showed no statistically significant difference between the prevalence of amoebiasis, cryptosporidiosis, and giardiasis across different age groups ( $P < 0.05$ ). As tabulated in table 2, the rate of amoebiasis was higher in females than in males (60.6% vs 39.4%), respectively. However, the rate of cryptosporidiosis was higher in males than in females (65.7% vs 34.4%), while the rate of giardiasis was equal in both genders (5%). These results were significant at  $P < 0.05$ . Table 3 gives information about the infection rate of intestinal protozoa among children according to the age group. The highest infection rates of intestinal protozoa, namely *E. histolytica*, *C. parvum*, and *G. lamblia*, were reported in the age range of 5-9 years (53.75%, 25.0%, and 60.0%), 1-4 years (28.75%, 20.0%, and 30.0%), and 10-14 years (17.55, 8.6%, and 10.0%), respectively. These differences according to the age groups were statistically significant at  $P < 0.05$ .


## 4. Discussion

The results of the present study demonstrated a relatively high prevalence rate of intestinal protozoan parasites among children in Duhok Province, Iraq. The overall frequency rate of protozoal infections was estimated at 27.7%, consistently reported by several studies performed in other countries (7).


**Table 1:** Prevalence of *E. histolytica*, *C. parvum*, and *G. lamblia* in Duhok City, Iraq (n=740)

Intestinal Protozoa	Frequency	Percent (%)
<i>E. histolytica</i>	160	78.0
<i>C. parvum</i>	35	17.1
<i>G. lamblia</i>	10	4.9
Total	205	100


The chi-square statistic is 0.0023. The *p*-value is 0.998869.  
The result is *not* significant at a  $P < 0.05$ .



**Fig 1:** Cyst stage of *Entamoeba histolytica*



**Fig 2:** Cyst stage of *Giardia lamblia*



**Fig 3:** Oocyst stage of *Cryptosporidium parvum* with acid fast stain

**Table 2:** Frequency of *E. histolytica*, *C. parvum*, and *G. lamblia* protozoan regarded to the gender

Intestinal Protozoa	Total No. (%)	Gender No. (%)	
		Males	Females
<i>E. histolytica</i>	160 (78)	63(39.4)	97(60.6)
<i>C. parvum</i>	35(17.1)	23(65.7)	12(34.3)
<i>G. lamblia</i>	10 (4.9)	5(50)	5(50)
Total No. (%)	205 (100)	91(44.4)	114(55.6)

The chi-square statistic is 21.3248. The  $p$ -value is 0.000023. The result is significant at a  $P < 0.05$ .

**Table 3:** Prevalence of intestinal parasites among children according to the age groups

Intestinal Protozoa	Total No. (%)	Age group (year) No. (%)		
		1-4	5-9	10-14
<i>E. histolytica</i>	160(78.0)	46(28.75)	86(53.75)	28(17.5)
<i>C. parvum</i>	35(17.1)	7(20.0)	25(71.4)	3(8.6)
<i>G. lamblia</i>	10(5.88)	3 (30.0)	6(60.0)	1(10.0)
Total No. (%)	205 (100)	56(27.3)	117(57.1)	32(15.6)

The results are significant at a  $P < 0.05$

In the present study, the most prominent intestinal protozoa infections were *E. histolytica* (78.0%) and *C. parvum* (17.1%), while the lowest infection rate was caused by *G. lamblia* (4.9%). Earlier studies in Iraq have revealed that the most prevalent intestinal protozoan infections among patients who visited Azadi Teaching Hospital in Duhok City Kurdistan Region were *E. histolytica* and *G. lamblia*. The prevalence of intestinal protozoan infections and the reported species vary in different studies conducted in Iraq. Nevertheless, most of these studies have reported that *E. histolytica* caused the highest prevalence of infection. The highest prevalence rate of *E. histolytica* recorded in the present research was in agreement with the results of a study performed by Salman et al. (2019) (8). They reported that the highest rate of Amoebiasis was 27.0% among diarrheic children attending a hospital in Al-Karkh region, Baghdad Province. The second-highest prevalence rate of *C. parvum* in the current study was in line with the results of a study conducted by Kanabe and Darogha (9) in Erbil City, Iraq, where the prevalence of *C. parvum* was 18.94%. Finally, the lowest infection rate of *G. lamblia* was calculated at 3.0% in the present study, which was consistent with that reported by Bazzaz et al. (10). The findings of the current research was in agreement with a previous study conducted in Erbil City in term of reporting similar species of intestinal parasites; however, prevalence rates were slightly lower than those found in our study (11). This research reported that 61.24% and 26.60% of subjects were infected with *E. histolytica* and *G. lamblia*, respectively. A further study conducted in Mosul, Iraq, on 2,296 subjects found that the rates of *E. histolytica* and *G. lamblia* in children associated with diarrhea were 14.5% and 0.96%, respectively (12). Another study performed in Duhok, Northern Iraq, revealed that the frequency rates of *E. histolytica* and *G. lamblia* among patients attending Azadi Teaching Hospital were 28.3% and 1.1%, respectively (13). The results of a study conducted in Mazandaran Province, Iran, demonstrated that the prevalence of *G. lamblia* infection among the general population was 4.6% (14). This variation in the rate of parasitic infections can be attributed to differences in the number of samples, design of the study, laboratory tests, sanitary conditions, type of water consumed, economic standards, climate and environmental factors, hygiene levels, behavior style, and type of drinking water. The present study pointed out that the prevalence of *E. histolytica* was higher than that of *G. lamblia* infection across different age groups. In a similar study performed by Omar and Ismael (15) in Duhok, it was found that the infection rate of amoebiasis was higher than that of giardiasis in children. These results were inconsistent with those of the two studies carried out in Iran by Kia et al. (16)

and Zaki (11), who recorded that the infection rate of giardiasis was higher than amoebiasis in children. The prevalence rate of *E. histolytica* was higher among females than males in this study. In a similar vein, Haji and Bamarni (17) reported that the infection rate was higher among female children. Inconsistent with the results of the present research, a related study reported that the rate of *E. histolytic* (61.7%) was higher in males than in females (18). The differences observed between genders in terms of susceptibility to infections can be attributed to variations in endocrine immune interactions, specifically in the levels of sex steroid hormones, such as androgens and estrogens. These hormones have been shown to modulate different aspects of the host immune response by regulating the expression of Toll-like receptors and the production of humoral immunity (19). In contrast to the current research findings, a study has reported a higher prevalence of giardiasis among males compared to females. In disagreement with the results of the present research, one study reported that the prevalence rates of giardiasis were 4.7% and 4.5% in males and females, respectively (13). The higher rate of infection with *C. parvum* in males in this study was in agreement with a study by Ahmed et al. performed in Zakho District, Duhok, Iraq (20). The results of our study found that the infection rate of amoebiasis, cryptosporidiosis, and giardiasis was higher in the age range of 5-9 years. The results of the present study agreed with a study conducted in Zakho City on primary school children. The referred study found that the highest infection rate occurred in 9-year-old children (21, 22). The higher prevalence of infection among 5-9-year-old students could be attributed to less attention paid by their mothers to personal hygiene as children in this age group are more active and tend to play outdoors, exposing them to more dirt and potential sources of infection. On the other hand, older individuals tend to be more conscious of personal hygiene and take better care of themselves. It is suggested that timely measures be taken by carrying out long-term control strategies, enhancing personal cleanliness standards, and improving living environments and sanitary conditions.

#### Acknowledgment

The authors would like to express their deepest gratitude to the Hevi Private Hospital team in Duhok City for their help and support during this study.

#### Authors' Contribution

Study concept and design: Ismael, SS. Abdullah, BH. & Sadiq, AJ.

Shameeran Salman Ismael, Bland Husamuldeen Abdullah and Amal Jamil Sadiq

Acquisition of data: Ajaj, JS. Ali, NS. Omer, DM. Nori, NY.

Analysis and interpretation of data: Sadiq, AJ.  
 Drafting of the manuscript: Ismael, SS. & Abdullah, BH  
 Critical revision of the manuscript for important intellectual content: Ismael, SS. & Abdullah, BH  
 Statistical analysis: Sadiq, AJ.  
 Administrative, technical, and material support: Ismael, SS.

### Ethics

The study proposal were approved by the ethics committee of the College of Health Sciences, University of Duhok, Duhok, Iraq

### Conflict of Interest

The authors declare that they have no conflict of interest.

### Data Availability

The data that support the findings of this study are available on request from the corresponding author.

### References

- Fletcher SM, Stark D, Harkness J, Ellis J. Enteric protozoa in the developed world: a public health perspective. *Clinical microbiology reviews*. 2012; 25(3):420-449.
- World Health Organization .The global burden of disease: (2004) update. WHO press, Geneva, Switzerland 2008.
- Levy K. Does Poor Water Quality Cause Diarrheal Disease?. *The American journal of tropical medicine and hygiene*. 2015; 93(5):899-900.
- Ortega YR, Eberhard ML, Kris H. Protozoan diseases: cryptosporidiosis, giardiasis and other intestinal. *Parasitologists United Journal*. 2008; 88 protozoan diseases. In: Heggenhougen K (ed) *International encyclopedia of public health*. Academic Press, Oxford, United Kingdom, pp: 354-366.
- Farthing MJG, Cevalls AM, Kelly P. (2009). Intestinal protozoa. In: Gook GC, Zumla A (eds) *Manson's tropical disease*, 22nd. WB Saunders, London. 2009; pp: 1375-1406.
- Fotadar R, Stark D, Beebe N, Marriott D, Ellis J, Harkness J. Laboratory diagnostic techniques for Entamoeba species. *Clinical microbiology reviews*. 2007;20(3):511-32, table of contents
- Murad MA, Al-Saeed ATM, Mustafa AA. Prevalence of Intestinal Parasites Among Patients Attended Azadi Teaching Hospital in Duhok City-Kurdistan Region/Iraq. *Journal of Duhok University*. 2019; 21(2):68-74.
- Salman AO, Mhaisen FT, Abdul-Rahman A, Al-Tae A. On the occurrence of some intestinal parasites among Diarrheic children attending a hospital in Al-Karkh Side, Baghdad City, Iraq. *Biological and applied Environmental Research* 2019; 3: 93-102.
- Kanabe LO, Darogha SNR. Epidemiology of Cryptosporidiosis among diarrheic children of Raparin Pediatric hospital, Erbil province-Kurdistan Region, Iraq. *Cihan University-Erbil Scientific Journal*. 2017; 2: 538.
- Bazzaz AA, Ouhood M, Shakir OM, Alabbasy RH. Prevalence of two gastrointestinal parasites *Entamoeba histolytica* and *Giardia lamblia* within Samarra City, Iraq. *Advances in Bioscience and Biotechnology*. 2017;8:399-410
- Zaki DM. Prevalence of *Entamoeba histolytica* and *Giardia Lamblia* Associated with Diarrhea in Children referring to Ibn Al-Atheer Hospital in Mosul, Iraq. *Archives of Razi Institute*. 2022; 77(1):73-9
- Dhubyan Mohammed Zaki Z. Prevalence of *Entamoeba histolytica* and *Giardia Lamblia* Associated with Diarrhea in Children referring to Ibn Al-Atheer Hospital in Mosul, Iraq. *Archives of Razi Institute*. 2022; 77(1):73-79. Published 2022.
- Hooshyar H, Rostamkhani P, Arbabi M, Delavari M. *Giardia lamblia* infection: review of current diagnostic strategies. *Gastroenterology and hepatology from bench to bench*. 2019; ;12(1):3-12.
- Siyadatpanah A, Sharif M, Daryani A, et al. Spatial distribution of *Giardia lamblia* infection among general population in Mazandaran Province, north of Iran. *Journal of parasitic diseases*. 2018;42(2):171-176.
- Omar AJ, Ismael SS. Detection of Pathogenic Strains of *Entamoeba Histolytica* in Children Using ELISA Technique in Duhok. *Detection of Pathogenic Strains of Entamoeba histolytica in Children Using ELISA Technique in Duhok*. *Journal of US-China Medical Science* 2011; 8 (2): 109-119
- Kia EB, Hosseini M, Nilforoushan MR, Meamar AR, Rezaeian M. Study of Intestinal Protozoan Parasites in Rural Inhabitants of Mazandaran Province, Northern Iran. *Iranian Journal of Parasitology* 2008; 3(1):21-25
- Haji HM, Bamarni SSI. Prevalence of *Entamoeba histolytica* and *Giardia lamblia* in Children in Duhok Province, Kurdistan Region, Iraq. *Journal of Duhok University*. 2023; 26(1), 274-280.
- Hegazi MA, Patel TA, El-Deek BS. Prevalence and characters of Entamoeba histolytica infection in Saudi infants and children admitted with diarrhea at 2 main hospitals at South Jeddah: a re-emerging serious infection with unusual presentation. *Braz J Infect Dis*. 2013;17(1):32-40.
- Klein SL. Hormones and mating system affect sex and species differences in immune function among vertebrates. *Behavioural processes*. 2000; 51(1-3):149-66.
- Ahmed HI, Salih Mero WM, Basheer Mohameed A. Prevalence of Cryptosporidiosis and It's Associated Risk Factors Among Human Population in Zakho District, Duhok Province, Kurdistan Region, Iraq. *Science Journal of University of Zakho*. 2022; 10(4), 153–158.
- Jameel AY, Issa AR, Amidy KSK, Mero WMS, Sabri MA, Khdihir MA. Prevalence of Intestinal Parasites under Fingernails of Primary School Children in Zakho, Kurdistan Region. *Science Journal of University of Zakho*. 2017;5(1):25-7.
- Ismael, S. Prevalence of Trichomoniasis and Vulvovaginal Candidiasis Among Married Women in Duhok City, Kurdistan Region, Iraq. *Archives of Razi Institute*. 2023.