



Epidemiological and Clinical Characteristics of Patients with Diphtheria Attending the Infectious Disease Hospital in Delhi

Kamble, BD¹, Singh, SK², Chellaiyan, VG^{3*}, Jethani, S², Gunjan, M², Jenniferbritto, J⁴

1. Department of Community Medicine & Family Medicine, All India Institute of Medical Sciences, Bibinagar, Hyderabad Metropolitan Region, Telangana-508126. India
2. Department of Community Medicine. North DMC Medical College & Hindu Rao Hospital, Delhi. India.
3. Department of Community Medicine, Chettinad Hospital and Research Institute, Chettinad Academy of Research and Education, Kelambakkam, Chennai. India.
4. Department of Obstetrics & Gynecology, Chettinad Hospital and Research Institute, Chettinad Academy of Research and Education, Kelambakkam, Chennai. India

How to cite this article: Kamble BD, Singh SK, Chellaiyan VG, Jethani S, Gunjan M, Jenniferbritto J. Epidemiological and Clinical Characteristics of Patients with Diphtheria Attending the Infectious Disease Hospital in Delhi. *Archives of Razi Institute Journal*. 2024;79(2):355-360. DOI: 10.32592/ARI.2024.79.2.355



Copyright © 2023 by



Razi Vaccine & Serum Research Institute

ABSTRACT

Diphtheria is a re-emerging disease with a changing epidemiology. It is thus essential to recognize diphtheria's clinical profile, patterns of morbidity and mortality, and the level of immunization. This study aimed to analyze the clinical-epidemiological profile of patients with diphtheria at the Infectious Disease Hospital, New Delhi, India. The present study was a cross-sectional investigation conducted on 94 patients diagnosed with diphtheria. The data collection tool was a questionnaire that had questions eliciting details such as socio-demographic characteristics, clinical history, examination findings, and vaccination history. The case fatality rate was calculated. Means and proportions were also measured, and the significance level was set at $P < 0.05$. The mean \pm SD age of the participants was 9 ± 4.4 years. Laryngeal involvement was found in 10 (11%) participants. Complete diphtheria vaccination doses were taken by 6 (6.4%) participants. The case fatality rate was 13%, and complications such as neuropathy were found in 21 (22%) cases, cardiac problems in 12 (13%), and respiratory problems in 13 (14%). Longer duration of the illness, delayed presentation, and complications were associated with poor clinical outcomes ($P < 0.05$). The majority of diphtheria cases were 6-10 years old and had a low socio-economic status. Most of them had a partial vaccination against diphtheria. Around one-third of the patients had complications related to diphtheria, and the case fatality rate was also high. Key preventive measures for controlling the deadly diphtheria disease include specialized immunization programs for areas with poor immunization coverage, as well as the early detection and treatment of suspected cases.

Keywords: Clinical profile, Complications, Diphtheria, Mortality

Article Info:

Received: 22 August 2023

Accepted: 15 October 2023

Published: 30 April 2024

Corresponding Author's E-Mail:
drchellaiyan@gmail.com

1. Introduction

Diphtheria is caused by gram-positive *Bacillus* known as *Corynebacterium diphtheriae* (1). It spreads from person to person, often via the respiratory route and occasionally through direct contact with infected skin lesions. Although diphtheria is a vaccine-preventable disease, it can become severe in rare cases and lead to several complications or even death. Diphtheria was a major cause of childhood morbidity and mortality before the introduction of its vaccine (2). However, despite universal immunization programs, this disease continues to be reported in many regions of India. Many outbreaks of diphtheria from various states in India have been reported in recent times (3-6). The present situation in India suggests a lack of immunization coverage and awareness, as well as the necessity for appropriate intervention programs and laboratory confirmation. Indian hospitals should aim to identify clinical and demographic characteristics of diphtheria cases and predictors of mortality at an early stage. The lack of attention to diphtheria over the last century resulted in knowledge gaps about its epidemiology, transmission, and control. Although diphtheria has been eliminated by many developed countries through effective immunization, it continues to exist in India and is the leading cause of morbidity and mortality in various states and border regions. Diphtheria is a re-emerging disease with a changing epidemiology. Therefore, there is a need to recognize the clinical profile, morbidity, and mortality pattern of diphtheria, as well as the immunization status against it. As for the changing epidemiology of diphtheria, studies have reported clinical profiles, and varying patterns remain scanty in North India. This study thus aimed to investigate the clinical and epidemiological profile of patients with diphtheria at the Infectious Disease Hospital, New Delhi, India.

2. Materials and Methods

The present study was a cross-sectional investigation conducted at the Infectious Disease Hospital, New Delhi, India. After obtaining the required consent from patients or their guardians, a total of 94 patients with a confirmed diagnosis of diphtheria at the age of 1-30 years from May 2020 to October 2020 were included in this study. The WHO case definition of diphtheria is an illness characterized by laryngitis, pharyngitis, or tonsillitis, as well as an adherent membrane of the tonsil, pharynx, and/or nose (7). Laboratory criteria for diagnosis were the isolation of *Corynebacterium diphtheriae* from a clinical specimen or a fourfold greater rise in serum antibody (only if both serum samples were obtained before the administration of diphtheria toxoid or antitoxin). A confirmed case was a case that is laboratory-confirmed or was epidemiologically linked to a laboratory-confirmed

case. A questionnaire was used to elicit details such as socio-demographic characteristics, clinical history, examination findings, and vaccination history. The case fatality rate was also calculated. The outcomes were classified as good and unfavorable. Ethical approval was obtained from the Institutional Ethics Committee, North DMC Medical College, Delhi, India (IEC ref No. 639). Permission from the Medical Superintendent of Infectious Disease Hospital, Delhi, was obtained for conducting the study and accessing patients' data. All information collected during the study was kept confidential. No personally identifying information was disclosed. Written informed consent was also obtained from adult patients and the guardians of younger patients. Data were entered into Microsoft Excel (version 2016; Microsoft, Redmond, WA, USA) and analyzed using STATA (version 13; Stata Corp., LLC, 4905 Lakeway Drive, College Station, TX 77845, USA). Descriptive statistics were applied to analyze sociodemographic details, clinical symptoms of patients, and complications. The Chi-squared test was applied to find differences, and a *P*-value of <0.05 was considered significant.

3. Results

The mean±SD age of the study participants was 9±4.4 years, with the minimum and maximum age being 1 and 28 years, respectively (Table 1). The throat swab was positive in 79 (84%), the nasal swab was positive in 24 (26%), and the contact history was positive in 6 (6.4%) participants. Complete diphtheria vaccination doses were taken by 6 (6.4%), partial vaccination was taken by 82 (87%), and vaccination status was unknown in 6 (6.4%) participants (Table 2). The median (IQR) duration of illness was 14 (12-19) days (Figure 1). Among the study participants, 82 (87%) recovered and were discharged, whereas 12 (13%) expired. The case fatality rate was 13%. No complications were developed in 65 (69%) participants (Table 3). Longer duration of illness, delayed presentation, and complications were associated with poor clinical outcomes (Table 4).

4. Discussion

In the present study, female participants (56.0%) were more than males (44.0%). In a similar study conducted in Karnataka by Harwalkar KK (8), 52.6% of the participants were male and 47.3% were female. In addition, in this study, almost half of the cases (54.0%) were 6-10 years old, while only 4.0% were 19-30. In a similar study conducted in Lucknow by Singh SN (9), almost half of the participants (49.5%) were 1-5 years old, followed by 43% in the age group of 5-10 years old. Harwalkar KK (8) study found that diphtheria was more common in the age group of 6-10 years.

Table 1 Baseline Characteristics Of The Study Participants. (n=94)

S. no.	Socio-Demographic Profile	N(%)
1.	Gender	
	Male	41(44)
	Female	53(56)
2.	Age group (in years)	
	1 - 5	14(15)
	6- 10	51(54)
	11- 18	25(27)
	19 - 30	4(4)
3.	Residence	
	City	37(40)
	Town	21(22)
	Village	36(38)
4.	Education of father	
	Illiterate	32(34)
	Primary	42(45)
	High secondary	18(19)
	Graduate and above	2(2)
5.	Occupation of the father	
	Unemployed	3(3)
	Unskilled/ Semiskilled	65(69)
	Skilled	2(2)
	Clerk, Shop owner, farmer	24(26)
6.	Education of mother	
	Illiterate	63(67)
	Primary	21(22)
	High secondary	10(11)
7.	Occupation of the mother	
	House wife	72(76)
	Unskilled/ Semiskilled	12(13)
	Skilled	10(11)
8.	Religion	
	Hindu	50(53)
	Muslim	44(47)
9.	Socioeconomic class	
	Upper	0
	Upper Middle	2(2)
	Lower Middle	15(16)
	Lower	77(82)
10.	Type of family	
	Nuclear	75(80)
	Joint	19(20)

Table 2: Clinical Presentation Among The Patients With Diphtheria. (N=94)

S. no.	Clinical symptoms	N(%)
1.	Clinical symptoms*	
	Fever	94(100)
	Throat pain	90(96)
2.	Membrane	94(100)
	Laryngeal involvement	
3.	Yes	10(11)
	No	84(89)
4.	Pharyngeal/tonsillar involvement	
	Yes	14(15)
5.	No	80(85)
	Cervical lymphadenopathy	
6.	Yes	21(23)
	No	73(77)
7.	Stridor	
	Yes	8(8)
8.	No	86(92)
	Membrane	
9.	Thin	4(4)
	Thick	90(96)
10.	Membrane colour	
	White	89(95)
11.	Grey	11(5)
	Membrane removal	
12.	Removal on striping	35(37)
	Non-removal	59(63)
13.	Illness before admission	
	< 5 days	11(12)
	5- 10 days	67(71)
14.	>10 days	16(17)

*multiple options


**Figure 1:** Mean duration (in days) of illness among the study participants. (n=94)

Table 3 Complications Among The Patients With Diphtheria. (N=94)

S. no.	Complications	N(%)
1.	Neuropathy	
	Palatal	11(12)
	Palatal + polyneuropathy	21(22)
	Carditis	
	Tachyarrhythmia	6(6)
	Bradyarrhythmia	12(13)
	Acute renal failure	2(2)
	Thrombocytopenia/Bleeding disorders	5(89)
	Respiratory failure/Airway compromise	13(14)
	ECG abnormalities	
No complications	65(69)	

Table 4: Comparison Of Outcome With Selected Variables. (N=94)

S.No.	Variables	Survived N=82	Expired N=12	OR(95%CI)	P-value
1.	Gender				
	Male	33(40)	8(67)	1.0	0.139
Female	49(60)	4(33)	2.1(0.3-4.3)		
2.	Age group				
	≤ 10	57(69)	8(67)	1.0	0.098
>10	25(31)	4(33)	3.2(0.9-2.1)		
3.	Duration of illness				
	≤ 10 days	45(55)	2(17)	1.0	0.021
>10 days	37(45)	10(83)	2.6(1.4-3.8)		
4.	Complications				
	No complications	64(78)	1(8)	1.0	0.03
Complications	18(22)	11(92)	3.4(1.8-4.5)		
5.	Illness before admission				
	< 10 days	76(80)	2(17)	1.0	0.01
>10 days	6(20)	10(83)	2.1(1.6-3.9)		

Chi square tests, p value of > 0.05 is significant

In terms of clinical manifestations, all our study participants (100%) had fever and the presence of membranes, and the majority (96%) had throat pain. Similar findings were found in a study conducted in Maharashtra by Meshram RM (10), in which all patients presented with fever and membrane in the throat, and the majority (95.74%) had throat pain. The present study also found that only 11% had laryngeal involvement and 15% had pharyngeal/tonsillar involvement. In another study conducted in Maharashtra by Meshram RM (10), the majority of participants (80.85%) had enlarged/congested tonsils. Furthermore, out of all the patients in our study, about one-fourth (23%) had cervical lymphadenopathy, and only 4% had a thin membrane. Another similar study by Singh SN (9) revealed that about two-thirds (63.9%) of study participants had neck swelling, and one-fourth (76.4%) had laryngeal membrane involvement. Moreover, in the present study, the majority of participants (71%) had an illness 5-10 days before admission, while 12% of them had the illness less than 5 days before admission. Another study by Singh SN (9) reported that the majority (83.9%) of participants presented with an illness they had

for less than 14 days. Considering vaccination, only 6.4% of the participants in this study had taken complete diphtheria vaccination doses, and the majority (87.0%) of participants had had partial vaccination. In a similar study by Harwalkar KK (8), more than half of the participants (59.5%) were partially immunized. In another study conducted in Maharashtra by Meshram RM (10), it was found that only 4.25% of participants were completely immunized, while more than half (57.45%) were partially immunized, and about two-thirds (38.30%) were not immunized at all. In addition, in the present study, the majority of the participants (87%) were recovered and discharged, and 13% of them expired. However, in a similar case study conducted in Maharashtra by Meshram RM (10), about one-fourth (21.2%) of the participants expired. In another study by Singh SN (9), about half of the participants (48.0%) expired. In the present study, about two-thirds of the participants (69.0%) had developed no complications. Among those who developed a complication, the most common complication (69.0%) was respiratory failure/airway compromise, followed by one-fourth (22.0%) having palatal palsy and polyneuropathy. Only 6% of the

participants in the present study had tachyarrhythmia and bradyarrhythmia. The most common complication (51.6%) reported was airway compromise, while only 7.2% and 14.0% had developed tachyarrhythmia and bradyarrhythmia, respectively. A similar study by Harwalkar KK (8) revealed that palatal palsy was the most common complication (10.7%), followed by myocarditis (8.6%) and asymptomatic bradycardia (4.3%). In a similar case study conducted in Maharashtra by Meshram RM (10), myocarditis was the most common complication (42.5%), followed by palatal palsy (14.89%), and only 4.25% developed acute renal failure and shock. In this study, longer duration of illness, delayed presentation, and complications were significantly associated with poor clinical outcomes, while they were not significantly associated with age group or gender. Another similar study conducted in Singapore by Ang LW (11) revealed that the seroprevalence in males (93.7%) was not significantly different from that of females (90.8%). There was also no difference in seroprevalence according to gender distribution, as reported by Meshram and Patil (10). Moreover, Ang LW (11) revealed that the prevalence of diphtheria declined significantly with age. In this study, the majority of diphtheria cases were 6-10 years old and had a lower socio-economic status. In addition, most of the patients had a partial vaccination against diphtheria, and around one-third of them had complications related to diphtheria. The case fatality rate was also high. Longer duration of illness, delayed presentation, and complications were associated with poor clinical outcomes. Therefore, special immunization campaigns for areas with low immunization coverage and the early identification and treatment of suspected cases will be key preventive strategies for controlling the deadly disease of diphtheria.

Novelty of the Study

This study reports findings from an exclusive Infectious Diseases Hospital, New Delhi, India. The study reported the case fatality rate of a vaccine-preventable disease, diphtheria, and assessed in detail the determinants of the clinical outcomes.

Acknowledgment

None

Authors' Contribution

BK, SKS, VGC, GM, and SJ conceived the idea and planned the study. VGC, BK, SKS, and JJ performed data extraction and tabulation. VGC, BK, SKS, SJ, GM, and JJ prepared the manuscript. All the authors read and approved the final manuscript.

Ethics

The study was conducted after receiving approval from the Institutional Ethics Committee. The confidentiality of study participants is maintained.

Conflict of Interest

None.

Grant Support

None.

References

- Gupta A, Gujral M, Singh A, Chawla K. Diphtheria. resurgence in India: A case study from south Karnataka. *J Family Med Prim Care*. 2020;9:5776-8. DOI: 10.4103/jfmpe.jfmpe_480_20
- Shaun AT, Keegan LT, Moss WJ, Chaisson LH, Macher E, Azman AS et al. Clinical and Epidemiological Aspects of Diphtheria: A Systematic Review and Pooled Analysis. *CID*. 2020;71:89-96. DOI: 10.1093/cid/ciz808.
- Maheriya KM, Pathak GH, Chauhan AV, Mehariya MK, Agrawal PC. Clinical and epidemiological profile of diphtheria in tertiary care hospital. *Gujarat medical J*. 2014;89(2):105-8. DOI: 10.18203/2349-3291.ijcp20182572
- Sardar JC, Saren AB, Haldar D, Chatterjee K, Biswas S, Chatterjee T et al. Obstinate diphtheria need innovation in immunization. *Int J Contemp Pediatr*. 2016;3(3):902-9. DOI: 10.18203/2349-3291.ijcp20162363
- Parande MV, Parande AM, Lakkannavar SL, Kholkute SD, Roy S. Diphtheria outbreak in rural North Karnataka, India. *JMM Case Reports*. 2014;1(3):14-9. DOI: 10.4103/ijmm.IJMM_17_48.
- Meera M, Rajarao M. Diphtheria in Andhra Pradesh-a clinical-epidemiological study. *Int J Infect Dis*. 2014;19:74-8. DOI: 10.1016/j.ijid.2013.10.017
- World Health Organization. WHO Vaccine-Preventable Diseases Surveillance Standards. Geneva, Switzerland: WHO.2018. WHO_SurveillanceVaccinePreventable_04_Diphtheria_R2.pdf.
- Harwalkar KK, Kadegaon B. Clinical profile of children with diphtheria admitted to the tertiary care centre. *Indian J Child Health*. 2019;6(10):563-65. DOI: 10.32677/IJCH.2019.v06.i10.012
- Singh SN, Singh A, Chandra S. Clinical profile and prediction of poor outcome of hospitalized diphtheria cases in children from Lucknow region of North India. *Clinic Epidemiol Glob Health*. 2014;2:75-9. DOI: 10.18203/2349-3291.ijcp20182572
- Meshram RM, Patil A. Clinical profile and outcome of diphtheria in central India: a retrospective observational study. *Int J Contemp Pediatr*. 2018 Jul;5(4):1600-1605. DOI: 10.18203/2349-3291.ijcp20182572Ang
- LW, James L, Goh KT. Prevalence of diphtheria and tetanus antibodies among adults in Singapore: a national serological study to identify most susceptible population groups. *J Pub Health*. 2015;38(1):99-105. DOI: 10.1093/pubmed/fdv011