

**Review Article****The Classification of Medicinal Plants used in Traditional Persian Medicine for the Treatment of Liver Disease based on Phytochemical Properties**Fatemeh Rabizadeh<sup>1\*</sup> and Maryam Sadat Mirian<sup>2</sup><sup>1</sup>Farzanegan Campus, Semnan University, Semnan, Iran<sup>2</sup>Department of Plant and Animal Biology, Faculty of Biological Science and Technology, University of Isfahan, Isfahan, Iran**Article History**

Received: 31 October 2023  
Accepted: 19 December 2023  
© 2012 Iranian Society of  
Medicinal Plants.  
All rights reserved.

**Keywords**

Liver  
Medicinal plants  
Phytochemicals  
Secondary metabolites  
Traditional Persian medicine

**\*Corresponding author**

f.rabizade@semnan.ac.ir

**ABSTRACT**

Chronic and acute liver diseases are considered a global issue and their medical treatments are commonly challenging to manage. Traditional medicines have used natural products for thousands of years to prevent and treat various diseases. Recent studies have revealed that the pharmacological impacts of herbs are primarily determined by their phytochemical constituents. Therefore, understanding plant chemistry is crucial for the therapeutic use of medicinal plants. In this review, we first introduced some medicinal plants that have the potential to be beneficial for treating liver diseases and disorders, based on Traditional Persian Medicine (TPM) textbooks. Subsequently, we investigated the secondary metabolites of these medicinal plants by analyzing pharmacological research collected from electronic databases. We also discussed their scientific and family names. According to TPM textbooks, 77 medical plants have been identified for the treatment of liver defects, belonging to 43 different families. Their secondary metabolites were studied through data obtained from electronic databases such as Google Scholar, PubMed, Science Direct, and Web of Science. These findings suggest that natural plant extracts hold promise for the prevention and treatment of liver diseases.

**INTRODUCTION**

Since ancient times, natural products such as plants, microorganisms, animals, and marine organisms have been used in medicines to attenuate injuries and treat diseases [1]. Early humans had an enormous challenge using natural products as medicines due to the consumption of poisonous plants, which resulted in diarrhea, vomiting, coma, or even death. This way made early humans expand their knowledge of edible plants and natural medicines. Traditional medicines have used natural products for the prevention and treatment of diseases for thousands of years [2]. The application of medicinal plant extracts for medical purposes originated and continues in traditional Indian, Egyptian, and Chinese medicine [3]. Several medicinal plants have been commercialized for their beneficial impacts on health, which are linked to various biological functions associated with their traditional applications [4]. Massage therapy and aromatherapy with herbal oils have been shown to have beneficial results and

promote immunological and physiological conditions in many ancient communities [3]. Recent studies have shown that the tremendous pharmacological impacts of herbs are fundamentally dependent on their phytochemical constituents, so that the study of plant chemistry is the foundation of the therapeutic uses of medicinal plants. Plants produce two large groups of compounds, primary and secondary metabolites. The secondary metabolites are small molecular products obtained from primary metabolites or play a role as intermediates in their biosynthetic pathways. These molecules show no noticeable effect on the essential cellular functions and are non-essential for the growth and reproduction of the organism, but play bio-ecological roles in the process of organism adaption to environmental niche [5]. Moreover, secondary metabolites have several biological effects, including antifungal, antibiotic, and antiviral properties. This makes them effective at protecting plants from pathogens and serving as vital UV-absorbing compounds that prevent severe leaf

damage from light. Secondary metabolites are classified into several large classes based on their chemical structure. Phenolics, terpenes, steroids, alkaloids, glucosides, amines, and flavonoids are some classes.

Recently, research works have shown that the secondary metabolites have positive beneficial effects on human health, agriculture production, and cosmetic products, contributing significantly to the economy [6]. It was proved that the secondary metabolites are responsible for the medicinal effect of plants. The importance of secondary plant metabolites in traditional medicine and folk uses cannot be overstated. These products of medical plants have been shown to alleviate a wide range of diseases. In modern medicine, secondary metabolites have been used as fundamental compounds to produce medications that treat different diseases, including various cancers and migraines.

Currently, there is no approved remedy for some infections, disorders, and diseases and besides, vaccination is limited to some viral infections. Additionally, medicines available on the market are often costly and may cause side effects. As a result, naturally based pharmacotherapy may be a proper alternative to treating diseases.

The liver is a vital organ in the human body that performs various functions such as detoxification, metabolism, and storage of nutrients. However, the liver can be affected by multiple diseases, such as hepatitis, cirrhosis, and liver cancer. Chronic and acute liver diseases are considered a global issue and their medical treatments are commonly challenging to manage and have restricted effectiveness [7]. For example, liver cancer is the most frequently detected form of cancer and has shown a significant rise in mortality rates worldwide. Surgery plays a crucial role in treating liver cancer; however, numerous patients with advanced stages or metastasis are not eligible candidates for surgical procedures [8]. Despite chemotherapy's ability to directly target cancer cells, it often has limitations such as resistance and side effects. Hence, there is a strong need for an effective strategy for liver cancer treatment that is non-toxic. Moreover, diabetes is a metabolic disease caused by reduced biological effects of glucose or insufficient insulin secretion in the body. This disease affects glucose metabolism and can lead to liver abnormalities and insulin resistance [9]. Various factors, including chemicals, drugs, and viruses, have

been reported to lead to serious liver necrosis, which can be difficult to manage with medical therapies. Therefore, it is crucial to find compounds that can effectively treat hepatic failure. Then, finding a new remedy that can safely and successfully prevent or treat liver diseases is a top priority. A wide variety of herbs, plant extracts, and plant-isolated compounds have been investigated for their beneficial effects on liver diseases.

Traditional medicine and folk remedies have utilized secondary plant metabolites for centuries to treat liver ailments. Recent studies have shown that these compounds possess biological properties such as antioxidant and anti-inflammatory effects that can protect the liver from damage. Furthermore, some secondary metabolites have been found to have hepatoprotective effects, which can prevent liver diseases. Therefore, understanding the role of secondary metabolites in liver health and the environmental factors that affect their production is crucial for developing effective treatments for liver diseases.

Nowadays, despite different opinions among medical professionals and complementary medicine practitioners, there is a rising trend to use alternative and complementary medicine which is widely permitted by current legal regulations across the world [10]. There are precious medical works by Iranian sages mentioned in traditional Persian medicine (TPM) books that may be useful for alternative treatments of some diseases.

TPM is an entire system of medicine that dates back to ancient Persia. Studies have shown that TPM has a rich history of using medicinal plants and herbs for treating various diseases. A number of these herbs have been proven to possess anti-inflammatory, antioxidant, and anti-cancer properties by laboratory research. Traditional knowledge of natural medicine is recorded in historical manuscripts. Some of the most popular and influential ancient Persian medical references were Razi or Rhazes *Continens Liber* (Kitab- al-Havi) by Rhazes and *Canon of Medicine* (Al-Qanun-fi-al-Tibb) by Avicenna.

The present review focused on effective herbs for liver diseases and disorders based on ancient Iranian books and studied bioactive constituents isolated from medicinal plants based on the scientific literature.


## METHODS

This review was designed to gather information regarding herbal medications for the liver in TPM. To search for TPM references, Makhzan-ol-Advieh, The Canon of Medicine, and Taghvim al-Abdan fi Tadbir al-Ensan were used. In the next step, the classification of plants has been performed based on their scientific names and phytochemical compositions. The required data have been collected from electronic databases such as Google Scholar, PubMed, Science Direct, and Web of Science. The main findings are summarized in the table.


## RESULTS

In the current review, a total of 77 medical plant species belonging to 42 families have been reported

to treat liver diseases in TPM textbooks specifically. Table 1 shows the bioactive metabolites of medicinal plants in the current study with their scientific and family names. Among them, Apiaceae (7 species), Asteraceae (6 species), and Lamiaceae (4 species) were the dominant families (Figure 1). Figure 2 shows the percentage of the present medicinal plants containing bioactive metabolites. Terpenoids were observed in 68% of medicinal plants with considerable effects on treating liver diseases. Almost 52% of the plants studied in the current review contained phenolic components. Moreover, flavonoids and glycosides were observed in 20% and 16% of medicinal plants, respectively.



**Fig. 1** Percentage of the medicinal plant species from various families studied in the current review.



**Fig. 2** Percentage of medicinal plants containing secondary metabolites studied in the current review.

## DISCUSSION

These days, there has been a rising trend in the usage of alternative medicine. Accumulating evidence has illustrated that early treatment with traditional medicine has proven effective in preventing the progression of mild and moderate diseases into severe and critical conditions [90]. Moreover, alternative medicine has also been successful in providing an effective therapeutic strategy for patients with extreme illnesses and enhancing the ability of the body's resistance to remove pathogenic factors [91]. The biggest challenge in traditional medicine is finding proper remedies from those which are unsafe or ineffective.

**Table 1** Secondary metabolites of medicinal plants are used to treat kidney diseases according to traditional Persian medicine.

No.	Scientific name	Family name	Organ	Glycoside	Steroids	Tannins	Flavanoids	Phenolic	Terpenoid	Saponins	Alkaloid	References
1	<i>Ajuga chamaepitys</i> (L.) Schreb.	Lamiaceae	Aerial parts	( $\alpha$ -1,6-galactosyl sucrose)				iridoid	Eucalyptol $\alpha$ -Thujone $\beta$ -Thujone Camphor Endoborneol			[11]
2	<i>Ajuga iva</i> (L.) Schreb.	Lamiaceae	aerial parts					Linalool Methyl salicylate	$\alpha$ -pinene camphene p-cymene 1,8-cineol 1-octen-3-yl acetate Camphor Limonene			[12]
3	<i>Aloe vera</i> (L.) Burm.f.	Asphodelaceae	Leaf	glucomannan	Squalene			Anthraquinone Phytol				[13]
4	<i>Althaea sp.</i>	Malvaceae	Flower and root						Squalene			[14]
5	<i>Anacyclus pyrethrum</i> (L.) Link	Asteraceae	Root		Stigmasterol/ Gamma-Sitosterol			Benzaldehyde, 2-hydroxy-6-methyl	7-Tetradecenal, (Z) Squalene Bicyclo[7.2.0]undec-4-ene, 4,11,11-trimethyl-8-methylene-, [1R-(1R*,4Z,9S*)]-			[15]
6	<i>Apium graveolens</i> L.	Apiaceae	Leaf						Geraniol			[16]
7	<i>Aquilaria sinensis</i> Mer	Thymelaeaceae							caryophyllene oxide			[17]

8	<i>Aristolochia longa</i> L.	Aristolochiaceae			$\alpha$ -Thujone Spathulenol Valerena Hexanone Artemisia ketone	$\gamma$ -Selinene Zonarene 4,7(11)-diene	[18]
9	<i>Artemisia maritima</i> L.	Asteraceae	Fresh, authentic essential oil		$\alpha$ -Terpinene $\gamma$ -Terpinene $\beta$ -Thujone	Santolina alcohol	[19]
10	<i>Artemisia sieversiana</i> Ehrh. ex Willd.	Asteraceae	Stems and leaves		b-Myrcene Carene p-Cymene Borneol Terpineol-4 a-Terpineol a-Copaene Caryophyllene E-b-Farnesene Germacrene D E, E-a-Farnesene d-Cadinene g-Eudesmol T-Cadinol a-Cadinol a-Bisabolol	Selina-4,11-diene	[20]
11	<i>Arum italicum</i> Mill.	Araceae		guaiacylglycerol- $\beta$ -coniferyl	Myrcene Nerol Caryophyllene		[21]
12	<i>Arum maculatum</i> L.	Araceae	Roots		Carveol p-Cymene $\alpha$ -Pinene $\beta$ -Pinene Terpinolene		[22]

13	<i>Asarum europaeum</i> L.	Aristolochiaceae	Whole plants			$\alpha$ -Asarone, $\beta$ -Asarone	$\alpha$ -pinene camphene $\beta$ -pinene myrcene $\delta$ -3-carene p-cymene limonene (E)- $\beta$ -ocimene terpinolene linalool borneol terpinen-4-ol methyl thymol $\alpha$ -copaene	[23]
14	<i>Asparagus adscendens</i> Roxb.	Asparagaceae	Root			Spirostanosides		[24]
15	<i>Asparagus officinalis</i> L.	Asparagaceae	Aerial parts/Fruits/Roots		Quercetin Apigenin Baicalein Kaempferol	Caffeic acid		[25]
16	<i>Asparagus racemosus</i> Willd.	Asparagaceae	Aerial parts, Fruits and Roots		kaempferol		Sarsapogenin	[26]
17	<i>Brassica juncea</i> (L.) Czern.	Brassicaceae	Seeds and leaves	$\alpha$ -Methyl-D-mannopyranoside $\beta$ -D-Glucopyranoside $\alpha$ -D-Galactopyranoside 3,4-Altrosan	$\gamma$ -Sitosterol/ Stigmasterol $\beta$ -Sitosterol/ Stigmastan-3,5-diene			[27]
18	<i>Calamintha incana</i> (Sm.) Boiss.	Lamiaceae	Aerial parts			Thymol	Limonene p-Cymene 2-Hydroxypiperitone Caryophyllene oxide	[28]

19	<i>Capparis decidua</i> (Forssk.) Edgew.	Capparidaceae	Fruits	butyl isothiocyanate	$\beta$ -Sitosterol		[29]
20	<i>Capparis spinosa</i> L.	Capparidaceae	Roots and leaves	capparisine		furfural, bis(5-formylfurfuryl) ether	Alkaloids [30]
21	<i>Carica papaya</i> L.	Caricaceae	Leaf		Campesterol Stigmasterol	Squalene $\alpha$ -Terpineol (E)-Geraniol	[31]
22	<i>Carum bulbocastanum</i> L.	Apiaceae	Fruit			$\alpha$ -Pinene Camphene $\beta$ -Pinene Limonene Sabinene 1,8-Cineole $\gamma$ -Terpinene Linalool Camphor Borneol Carvone $\beta$ -Caryophyllene triterpene	[32]
23	<i>Carum carvi</i> L.	Apiaceae	Fruit			Eugenol $\alpha$ -Pinene $\beta$ -Pinene $\beta$ -Myrcene p-Cymene D-Limonene 1,8-Cineol Linalool -Terpineol $\gamma$ -Terpinene p-Cymen-8-ol Dihydrocarvone cis-Carveol Cuminaldehyde (R)-Carvone Dihydrocarveol $\alpha$ -Terpinene-7-al	[33]

24	<i>Carum copticum</i> L.	Apiaceae	Fruit		Thymol	Myrcene Limonene $\alpha$ -Thujene $\alpha$ -Pinene b-Pinene Sabinene $\alpha$ -Phyllanderene p-Cymene b-Phyllanderene $\gamma$ -terpinen Terpinene- 4 - ol Carvacrol cymene	[34, 35]
25	<i>Celastrus paniculatus</i> Willd.	Celastraceae	Seed	beta. - Sitosterol acetate Stigmast- 4-en-3- one		Maymysine Linalool Cubenol trans-beta-Copaene Muurolene Sclareolide	[36]
26	<i>Cichorium intybus</i> L.	Asteraceae	Root		lactucin, 8- deoxylactu cin	Lactupicri $\alpha$ -amyrin	[37]
27	<i>Cichorium pumilum</i> Jacq.	Asteraceae	Flower and aerial parts		Anthocyani ns	Lactucin Lactucopicrin	[38]
28	<i>Cinnamomu m bejolghota</i> (Buch. - Ham.) Sweet	Lauraceae	leaf			a-pinene camphene P-pinene	[39]



29	<i>Cornus mas</i> L.	Cornaceae	Fruits	<p>cyandin 3-O-rutinoside pelargonidin 3-O-rutinoside quercetin 3-O-robinobioside</p>	<p>pelargonidin 3-O</p>	<p>2-(4-Methylcyclohex-3-en-1-yl) propan-2-ol <math>\alpha</math>-terpineol 4,7,7-Trimethylbicyclo [2.2.1] heptan-3-ol/ borneol; 0.29 2-[(2R,5S)-5-Ethenyl-5-methyloxolan-2-yl] propan-2-ol/<i>cis</i>-Linalool Oxide (furanoid)*f; 2-[(2S,5S)-5-Ethenyl-5-methyloxolan-2-yl] propan-2-ol/<i>trans</i>-Linalool oxide(furanoid), (3S,6S)-6-Ethenyl-2,2,6-trimethyloxan-3-ol/; &lt;0.42 <i>cis</i>-pyranoid Linalool oxide Limonene</p>	[40]
30	<i>Cucumis colocynthis</i> (L.) Schrad.	Cucurbitaceae	Fruits		Isosaponarin		[41]
31	<i>Cucumis melo</i> L.	Cucurbitaceae	Fruits	<p>Amentoflavone Hydroxytyrosol Apigenin-7-glycoside</p>		<p>Gallic acid Protocatechuic acid Chlorogenic acid Isovanillic acid</p>	[42]
32	<i>Equisetum arvense</i> L.	Equisetaceae	Aerial parts		Hexahydrofarnesyl acetone Thymol		[43]

33	<i>Eugenia Jambolana</i> Lam.	Myrtaceae	Fruits	Cyanidin Cyanidin-3- glucoside Cyanidin- 3,5- diglucoside Cyanidin-3- diglucoside- 5-glucoside Malvidin	Galloyl - glucose ester	Dimethyl- dihydromy ricetindigl ucoside Isorhamnet in	Caffeic acid Gallic acid	a-Pinene β-Pinene a-Phellandrene Terpinolene γ-Terpinene Myrtenal Bornyl acetate	[44]
34	<i>Eupatorium cannabinum</i> L.	Asteraceae						Camphene Myrcene o-cymene p-cymene Terpinolene Linalool Nonanal thujone Camphor borneol terpinen-4-terpineol nerol methyl thymol geraniol caryophyllene	[45]
35	<i>Feronia elephantum</i> Correa	Rutaceae	Leaf /bark				Tryptamine , N-[4- hydroxyhy drocinnam oyl]-		[46]

36	<i>Foeniculum vulgare</i> Mill	Apiaceae	Aerial parts		Linalool Fenchol (z)-p-Terpeneol Nerolidol Isophytol Fenchone Camphor Carvone $\alpha$ -thujene $\alpha$ -pinene sabinene 3-pinene Myrcene $\alpha$ -phellandrene limonene isomer Palustrol	[47]
37	<i>Fumaria officinalis</i> L.	Fumariaceae			<i>p</i> -Coumaric acid Isoquercitrin Ferulic acid Rutin	[48]
38	<i>Fumaria parviflora</i> Lam.	Fumariaceae		$\alpha$ -D-diglucoside	Salicylic acid	[49]

39	<i>Gentiana lutea</i> L.	Gentianaceae	Root and Aerial parts		Vanillin	Hydroxy 2-butanone Linalool n-Menthone Isomenthone 4-Terpineol Carvone Thymol (E)-Caryophyllene (E)- $\alpha$ -Ionone -Gujunene trans- $\alpha$ -Bergamotene Aromadendrene Geranyl acetone $\gamma$ -Himachalene	[50]
40	<i>Glycyrrhiza glabra</i> L.	Fabaceae	Root	Pectin	Coumarins	Glycyrrhizin	[51]
41	<i>Hippophae rhamnoides</i> L.	Elaeagnaceae				Limonene cis-Ocimene	[52]
42	<i>Hyacinthus orientalis</i> L.	Liliaceae	Flowers	Anthocyanin 3,5-diglucosides	Delphinidin		[53]
43	<i>Hypericum barbatum</i> L.	Hypericaceae				$\alpha$ -Pinene Camphene $\beta$ -Pinene Limonene $\gamma$ -Terpinene Linalool 4-Terpineol Geraniol	[54]
44	<i>Hypericum coris</i> L.	Hypericaceae	Leaves and flowers			Linalool Geraniol $\alpha$ -copaene $\beta$ -caryophyllene $\alpha$ -humulene	[55]

45	<i>Hypericum perforatum</i> L.	Hypericaceae				a-Pinene Myrcene p-Cymene Selinene Cadinene Bisabolol β-Phellandrene γ-Terpinene Aromadendrene	[56]	
46	<i>Iris florentina</i> L.	Iridaceae		-		-	[57]	
47	<i>Laurus nobilis</i> L.	Lauraceae	Leaf			ATHujene αPinene Camphene Sabinene BPinene Myrcene α-Phellandrene δ-3-Carene α-Terpinene p-Cymene Limonene Cis-Ocimene γ-Terpinene Terpinolene Linalool	[58]	
48	<i>Lepidium draba</i> L.	Brassicaceae		Verbascoside		Chlorogenic acid	[59]	
49	<i>Linaria vulgaris</i> Mill.	Scrophulariaceae		Pectolinarin Linarin Linariin Pectolinarigenin Hispidulin	p-Coumaric acid p-Methoxybenzoic acid Glucosyringic acid Ferulic acid	Antirrhinose 6-O-trans-p-Coumaroylantirrhinose Procumbide 5-O-Allosylantirrhinose Aucubin Cycloart-25-ene-3b-,24-1-diol Stigmastadienone	Vasicine tricyclic quinazoline 7-Hydroxyvasicine Linarinic acid Choline Linavuline	[60]

50	<i>Malus orientalis</i> Ugl.	Rosaceae		Phloridzin		Gallic acid caffeic acid syringic acid chlorogenic acid	[61]
51	<i>Malva parviflora</i> L.	Malvaceae	leaves	Rutin	Luteolin Quercetin		[62]
52	<i>Malva sylvestris</i> L.	Malvaceae			Quercetin		[63]
53	<i>Mandragora officinarum</i> L.	Solanaceae				Hyoscyamine Cuscohygine Apoatropine 3a-Tigloyloxytr opane Belladonine	[64]
54	<i>Nymphaea alba</i> L.	Nymphaeaceae			Rutin Quercetin Naringenin Naringin Luteolin Apigenin Orientin	Caffeic acid Cinnamic acid derivative Catechin	[65]
55	<i>Petroselinum crispum</i> (Mill.) Fuss	Apiaceae	Fruit and leaves		Carvacrol	cymene $\alpha$ -Thujene $\alpha$ -Pinene $\beta$ -Pinene $\beta$ -Myrcene 3-Carene p-Cymene Sabinene $\gamma$ -Terpinene $\alpha$ -Terpinolene 4-Thujanol Thymol $\beta$ -carotene p-cymenene	[66]

56	<i>Pinus cembra</i> L.	Pinaceae	Limonene Myrcene $\alpha$ -Pinene $\beta$ -pinene $\gamma$ -terpinene $\beta$ -phellandrene $\alpha$ -cadinene $\gamma$ -Cadinene	[67]
57	<i>Pinus halepensis</i> Mill.	Pinaceae	Limonene Myrcene $\alpha$ - $\beta$ pinene fl-Pinene Terpinene Sabinene $\alpha$ -phellandrene fl-Phellandrene <i>p</i> -cymene	[68]
58	<i>Pinus pinea</i> L.	Pinaceae	Pinocarvone $\beta$ -Pinene $\alpha$ -longipinene Limonene + $\beta$ -phellandrene <i>p</i> -cymene Bornyl <i>acetate</i> Myrtenol verbenone	[69]
59	<i>Piper cubeba</i> L.	Piperaceae	Cadinene Linalool Sabinene	[70]

60	<i>Pistacia chia</i> Desf.	Anacardiaceae	Fresh bark and leaves		$\alpha$ -campholenaldehyde (E)-anethole	$\alpha$ -pinene Camphene Sabinene $\beta$ -pinene $\beta$ -myrcene p-cymene Limonene (Z)- $\beta$ -ocimene E)- $\beta$ -ocimene $\alpha$ -terpinolene linalool perillene cis-verbenol trans-pinocarveol trans-verbenol $\beta$ -pinene epoxide myrtenal $\alpha$ -terpineol verbenone trans-carveoldihydrocarveol linalyl acetate	[71, 72]
61	<i>Pistacia lentiscus</i> L.	Anacardiaceae	Fresh bark and leaves	Oligomers flavonoids		Limonene $\beta$ -pinene caryophyllene Terpinene-4-ol	[73]
62	<i>Pistacia vera</i> L.	Anacardiaceae	fresh commercial samples of shelled pistachio		Pinocarveol Vanillin Tyrosol (4-hydroxyphenylethanol)	[74]	
63	<i>Portulaca oleracea</i> L.	Portulacaceae	Seed			Squalene	[75]
64	<i>Prunus avium</i> L.	Rosaceae	Fruit		4-vinylphenol-eugenol	Thymol Carvacrol	[76]



65	<i>Rumex acetosa</i> L.	Polygonaceae	Leaf		Quercetin 3-O-rutinoside Quercetin 3- $\beta$ -D-glucoside <i>Quercetin derivative</i> Kaempferol 3-O-glucoside Luteolin 7-O-glucoside	Catechin Gallic acid Caftaric acid Caffeic acid		[77]
66	<i>Salix acmophylla</i> Boiss.	Salicaceae		Styrene	Salicin	Coumarin		[78]
67	<i>Salix aegyptiaca</i> Fors.	Salicaceae					Terpineol Cedrol	Alkaloid [79]
68	<i>Salix excelsa</i> S.G.Gmel.	Salicaceae					tetramethyl-2-hexadecene-1-ol	[80]
69	<i>Sempervivum arboreum</i> (L.) Webb & Berthel.	Crassulaceae	Leaf				3,3',4',5,7-pentahydroxy-6-methoxyflavanonol-3-O-glucoside	[81]
70	<i>Seseli tortuosum</i> L.	Apiaceae	Fruit			coumarin	Myrcene $\alpha$ -pinene $\beta$ -pinene limonene	[82]
71	<i>Solanum melongena</i> L.	Solanaceae	Fruit	Ethyl isoallochololate	2-Hydroxyethylphosphine		Neophytadiene Farnesyl acetone 4-Oxononanedioic acid Clionasterol Citronellyl formate 13-Tetradecenal Solanesol	Hexanal (Dimethylamino)acetone Methoxyethylamine Boscalid [83]

72	<i>Stachys officinalis</i> (L.) Trevis	Lamiaceae	Leaf and flower		Thujene Pinene Sabinene Myrcene Limonene Linalool trans-pinocarveol myrtenal verbenone cubebene copaene phytol	[84]
73	<i>Syzygium aromaticum</i> (L.) Merr. & L.M.Perry	Myrtaceae		Eugenol Thymol	Limonene $\alpha$ -Pinene <i>p</i> -cymene Cadinene Linalool	[85]
74	<i>Viola odorata</i> L.	Violaceae		Methyl eugenol	Pinocarvone Camphor Benzaldehyde Geranylacetone Bornyl <i>acetate</i> Borneol Limonene $\beta$ -Pinene $\gamma$ -Terpinene myrcene <i>p</i> -Cymene $\beta$ -Farnesene	[86]
75	<i>Vitis vinifera</i> L.	Vitaceae			Farnesene	[87]
76	<i>Zingiber officinale</i> Rosc.	Zingiberaceae	Root/Rhizome	2-Butanone, 4-(4-hydroxy-3-methoxyphenyl)-	$\zeta$ -Elemene/1H-Cyclopenta [1,3] cyclopropa[1,2]/Copaene benzene, octahydro-7-methyl-3-methylene-4-(1-methylethyl)-, [3aS-(3a $\alpha$ ,3b $\alpha$ ,4 $\alpha$ ,7 $\alpha$ ,7aS*)]	[88]
77	<i>Ziziphus jujuba</i> Mill.	Rhamnaceae		Eugenol Isoeugenol	Carvacrol Cadinene	[89]

It is necessary to assess and standardize all traditional drugs and methods for use in modern medicine according to contemporary pharmaceutical and medical standards. Over the past years, the properties of some medicinal plants mentioned in this study were investigated for liver dysfunction or diseases *in vitro* or *in vivo*. For this purpose, there is a need to have a clear view of traditional knowledge compared to current concepts.

The current review discusses the phytochemicals of medicinal plants of TPM that have been proven to be robustly effective in treating patients suffering from liver diseases. Several studies have investigated the potential hepatoprotective effects of plant extracts against acute liver injuries induced by carbon tetrachloride (CCl<sub>4</sub>), oxymetholone and thioacetamide, which activate hepatocyte damage in key markers. The effect of medicinal plants has been studied through evaluation of liver function markers such as aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), and total bilirubin were performed, along with oxidative stress parameters including malondialdehyde (MDA), glutathione (GSH), superoxide dismutase (SOD), and glutathione peroxidase as well as inflammatory mediators such as serum cytokines (IL-1 $\beta$ , IL-10, IL-6), tumor necrosis factor (TNF)- $\alpha$ , and nitric oxide.

The methanol extract from *Ajuga iva* has high concentrations of polyphenolics and flavonoids and has been found to inhibit key digestive enzymes linked to type 2 diabetes. It is particularly effective against  $\alpha$ -glucosidase and has a significant inhibition against  $\alpha$ -amylase, along with high antioxidant activity, while it is non-toxic. The results suggest that the phenolic compounds in the extract may be responsible for its antioxidant and antidiabetic activities [92]. *Aloe vera*, with various medicinal properties, has been used in ancient medicine to treat fever, burns, and wounds. Previous studies suggest that *A. vera* has anti-diabetic effects with protection against high fat and fructose diet-induced oxidative stress, dyslipidemia, and liver dysfunction. It also improves albumin levels and antioxidant enzyme activities in treated mice. The liver tissues of treated mice showed normal hepatocytes [93]. The ethyl alcohol-water extract of *Ziziphus jujuba* leaves, another plant with hepatoprotective properties, was found to have a pharmacological effect and significantly alleviate liver damage induced by CCl<sub>4</sub>

in mice. The results have provided evidence that *Z. jujube* not only offers maximum conjugation with dangerous free radicals and diminishes their toxic properties but also suppresses the inflammatory responses of a CCl<sub>4</sub>-induced liver injury. Flavonoids were identified as the active ingredients responsible for the biological and pharmacological activities toward hepatoprotection [94, 95]. *Capparis spinosa* L., also known as caper, is a plant traditionally used for medicinal purposes in various parts of the world. The fresh leaf and bud powders of *C. spinosa* contain multiple phytochemicals, including rutin, quercetin, and kaempferol. Daily administration of *C. spinosa* leaf or bud powder normalized biochemical parameters such as blood glucose, insulin, and lipid levels in rats with diabetes mellitus. The hepatoprotective properties of *C. spinosa* may be attributed to flavonoids like quercetin and rutin [4]. Another study found that the administration of *Piper cubeba* ethanolic extract protected against CCl<sub>4</sub>-induced hepatic damage in rats by downregulating proinflammatory cytokines and upregulating IL-10. *P. cubeba* ethanolic extract also prevented drug-induced increases in hepatic enzymes, reduced lipid peroxidation, and restored antioxidant enzyme activity along with prevention of CCl<sub>4</sub>-induced hepatic damage based on histopathological studies [96]. The potential protective effect of the eugenol-rich fraction of *Syzygium aromaticum* (clove) was investigated on liver cirrhosis induced by thioacetamide. The results showed that *S. aromaticum* had a protective effect on liver cirrhosis, as it inhibited hepatic cell proliferation and decreased oxidative stress. Eugenol, a major component of *S. aromaticum*, may be metabolized to dieugenol, which inhibits lipid peroxidation. This suggests that eugenol may have antioxidant properties that protect against liver cirrhosis [97]. *Anacyclus pyrethrum* L. with potential medicinal properties is commonly used in traditional North African and Indian medicine. The study investigated the anti-diabetic effects of the aqueous extract of *A. pyrethrum* roots in both normal and streptozotocin-induced diabetic rats by administration of a daily dose of 250 mg/kg extract after 21 days. The results showed that the *A. pyrethrum* extract exhibited significant antihyperglycemic activity in diabetic rats. Phytochemical screening of the extract revealed the presence of various compounds, including tannins, saponins, alkaloids, amino acids, steroids, and

terpenoids, which may contribute to the potential therapeutic properties of the extract [98]. The neuroprotective and hepatoprotective potentials of *Anagallis arvensis* were evaluated in rat models of interstitial cystitis and hepatotoxicity. The study found that pretreatment with *A. arvensis* significantly decreased the levels of liver markers and lipid profile due to its antioxidant phytochemicals [99]. The study investigated the antioxidant effects of a methanolic extract of *Apium graveolens* in a rat model of arthritis-induced liver oxidative stress. The rats were orally given different doses of *A. graveolens* extract for 24 days after inducing arthritis. The study found that *A. graveolens* treatment significantly reduced the levels of superoxide anion, total peroxide, and oxidative stress index in the liver. Additionally, the activity of glutathione peroxidase and SOD, which are antioxidant enzymes, significantly increased in the liver of arthritic rats treated with *A. graveolens* extract [100].

Furthermore, the hepatoprotective activity of an aqueous extract of *Artemisia absinthium* L. was investigated in a mouse model of liver injury induced by a single  $\text{CCl}_4$  administration or injection of endotoxin (lipopolysaccharide). The results of histopathological parameters showed that pretreatment with *A. absinthium* extract significantly prevented the increase in serum levels of hepatic enzymes in experimental mice-induced liver injury. Additionally, the extract reduced lipid peroxidation in the liver tissue and restored the activities of defense antioxidant enzymes SOD and GPX to normal levels. Histopathological examination showed a reduction in hepatocellular necrosis and reduced inflammatory cell infiltration. Phytochemical analysis revealed the presence of sesquiterpene lactones, flavonoids, phenolic acids, and tannins in *A. absinthium* [101]. In another study, the potential antidiabetic properties of ethanolic extract of *Asparagus adscendens* root were studied *in vivo* and *in vitro* models. They found that *A. adscendens* inhibited the activity of carbohydrate metabolizing enzymes, which are involved in the breakdown and utilization of carbohydrates in the body. The ethanolic extract also stimulated insulin release, suggesting that it may enhance the body's ability to regulate glucose metabolism and enhance glucose uptake, potentially aiding in the management of diabetes. Additionally, in an animal model of diabetes, *A. adscendens* decreased fasting blood

glucose levels and increased serum insulin and  $\alpha$ -amylase levels [102]. This study investigated the potential protective effects of *Carica Papaya* L. Seed extract against liver damage induced by  $\text{CCl}_4$  in rats. The study found that *C. Papaya* contained various antioxidants and minerals. The results showed that *C. papaya* extract treatment reduced oxidative stress, inflammation, fibrosis, and apoptosis induced by  $\text{CCl}_4$ . Liver and kidney function was also improved with *C. Papaya* seed extract treatment [103]. The impact of *Cornus mas* L. fruit extract was studied on liver function in nona-lcoholic fatty liver disease. Fifty patients were randomly assigned to receive either fruit extract for 12 weeks and levels of certain liver function markers were measured before and after the intervention. They found that cytokeratin 18 levels decreased significantly in the group treated with fruit extract compared to the control group [104]. Another study aimed to investigate the hepatoprotective effects of *Citrullus colocynthis* fruit extract on rats with hepatotoxicity induced by paracetamol. The results showed that rats treated with paracetamol had significantly elevated levels of liver function markers, but pretreatment with *C. colocynthis* fruit extract decreased these levels. The histopathological analysis also showed that fruit extract preserved the normal cellular architecture of the liver, suggesting that *C. colocynthis* has significant hepatoprotective and antioxidant activity [105]. The hepatoprotective effects of *Feronia elephantum* correa were evaluated against thioacetamide-induced liver necrosis in diabetic rats. The results showed that *F. elephantum* significantly reduced mortality and improved liver function parameters without affecting liver weight, volume, or serum glucose levels. The results concluded that *F. elephantum* could be useful for preventing liver complications in diabetes. The protective effects of *F. elephantum* against liver necrosis were attributed to its antioxidant activity, particularly from the flavonoids orientin and vitexin present in the extract [106]. Research work has been performed on the antidiabetic potential of *Fumaria officinalis*, a plant traditionally used to treat hypertension, hepatitis, and diabetes. Administration of the aqueous and methanolic extracts exhibited a significant hypoglycemic effect in alloxan-induced diabetic rats compared to normo-glycaemic rats. The extract also improved liver and kidney function tests and reduced damage to cells in the kidney and liver of diabetic rats

[107]. The potential hepatoprotective effects of methanol extracts from the aerial parts and roots of *Gentiana cruciata* L. were studied against liver injury induced by CCl<sub>4</sub> in rats. The extracts were found to contain high concentrations of sweroside, swertiamarin, and gentiopicrin. Pretreatment with *G. cruciata* dose-dependently and significantly reduced the levels of serum transaminases, ALP, and total bilirubin, while increasing the total protein level compared to the group treated with CCl<sub>4</sub> alone. Microscopic examination of the liver showed minimal CCl<sub>4</sub>-induced lesions and toxic manifestations in rats pretreated with extracts at 400 mg per kg body weight [108]. The effect of *Glycyrrhiza glabra* extracts was investigated on acute liver injuries induced by CCl<sub>4</sub>. Aqueous and ethanol extracts of *G. glabra* were used to administer orally to rats that were intraperitoneally injected with CCl<sub>4</sub>. The extracts significantly inhibited the activities of AST and ALT and increased the activity of SOD in both serum and liver tissue. Phytochemical analysis showed the presence of flavonoids and polysaccharides in the extracts, which may be responsible for their hepatoprotective activity [109]. In another study, the hepatoprotective effect of an aqueous extract of *Glycyrrhiza glabra* roots was evaluated in rabbit models with acute liver injury induced by CCl<sub>4</sub>. The results showed a significant reduction in hepatic enzyme levels, serum bilirubin, and improvement in serum protein levels in animals treated with the extract. The liver tissue also showed restoration of its architecture, absence of necrosis, and mild fatty infiltration [110]. Another study evaluated the hepatoprotective activity of *Hippophae rhamnoides* L. leaf extract on CCl<sub>4</sub>-induced liver injury in male albino rats. The extract significantly protected the animals from liver injury and enhanced antioxidant activity, suggesting it could be developed as a nutraceutical or food supplement against liver diseases [111].

Another study investigated the hepatoprotective potential of *Iris florentina* L. methanolic extract with flavonoids and phenols on paracetamol-induced liver injury in rats. The extract significantly improved serum biomarkers and restored hepatic injury, indicating hepatoprotective potential [112]. A study found that laurel leaf extract *Laurus nobilis* has potential as a natural remedy for managing liver damage induced by CCl<sub>4</sub> in male Wistar rats. Rats treated with the extract had significantly lower liver

damage indicators and less severe liver damage. The extract also increased levels of antioxidant enzymes, suggesting a protective effect against oxidative stress [113]. Another research investigated the protective effects of *Lepidium draba* extract on oxymetholone-induced hepato-renal toxicity in rats. The extract improved hepatic and renal biochemical parameters, reduced inflammatory cytokines and nitric oxide levels, and increased antioxidant enzyme activity. The high antioxidant and anti-inflammatory properties of *L. draba* are attributed to its phenolic and flavonoid components [114]. The antidiabetic activity of *Malva parviflora* L. leaf extract and its nano-formulation was measured in rats. The extract and its nano-formulation improved biochemical parameters, decreased glucose levels, increased insulin production, and improved the lipid profile of liver and kidney functions in diabetic rats, showing high antioxidant action and antimicrobial activity [115]. The aqueous methanolic extract of *M. parviflora* was studied to evaluate hepatoprotective activity in mice intoxicated with paracetamol. The extract significantly reduced liver enzymes and total bilirubin levels and was supported by histopathological investigation and detection of hepatoprotective constituents, suggesting that *M. parviflora* could be used as a natural remedy for liver damage. [116] *Nardostachys jatamansi* (D. Don) DC., an herb used in traditional Indian medicine, has been found to have hepatoprotective activity in rats. A 50% ethanolic extract of the herb's rhizomes significantly reduced liver damage caused by the toxic compound thioacetamide, as evidenced by lowered levels of serum enzymes and increased survival rates. This study supports the traditional use of *N. jatamansi* as a component of some hepatotoxic preparations used in Unani medicine. [117]. The potential hepatoprotective effects of *Nymphaea alba* L. leaf extract were measured on CCl<sub>4</sub>-induced hepatotoxicity in rats. The extract was found to significantly improve liver function, oxidative stress parameters, and TNF- $\alpha$ , as well as ameliorating histopathological features of the liver and decreasing caspase-3 expression, implying that *N. alba* leaf extract may be a therapeutic alternative for hepatic disorders. [118]. The study investigated the hepatoprotective effects of three plant extracts, *Pistacia lentiscus*, *Phillyrea latifolia* L., and *Nicotiana glauca* Graham, which are used in Jordanian folk medicine for the treatment of jaundice.

The extract effects were tested on liver function and serum bilirubin levels of rats with CCl<sub>4</sub>-induced hepatotoxicity. The non-boiled aqueous extract of *N. glauca* leaves reduced total serum bilirubin levels, while the boiled aqueous extract of *P. latifolia* reduced bilirubin and ALP levels without affecting ALT and AST activities. Both boiled and non-boiled aqueous extracts of *P. lentiscus* showed significant antihepatotoxic activity by reducing the activity of all three enzymes and bilirubin levels. The non-boiled extract was more effective than the boiled extract and may be a potential treatment for hepatic jaundice in humans [119]. The study aimed to investigate the potential hepatoprotective effects of *Pinus eldarica* extract on acetaminophen-induced liver injury in rats. The administration of *P. eldarica* extract significantly attenuated the increase in serum levels of ALT, AST, and ALP caused by acetaminophen. Additionally, *P. eldarica* administration prevented extensive necrosis and lymphocytic inflammation caused by acetaminophen, suggesting that *P. eldarica* has potential therapeutic effects on acetaminophen-induced liver toxicity in rats [120]. The potential protective effects of *Pistacia lentiscus* var. chia extracts were measured on CCl<sub>4</sub>-induced liver damage in rats. The results showed that *P. lentiscus* var. chia extracts have a strong inhibitory effect against lipid peroxidation in rat livers, with a decrease in levels of AST, ALT, and MDA. The pre-treatment *P. lentiscus* var. chia extracts reduced GSH depletion caused by CCl<sub>4</sub>, yielding GSH levels comparable to that observed in untreated rats, showing that *P. lentiscus* var. extracts protect liver cells from CCl<sub>4</sub>-induced oxidative damage [120]. The effects of *Portulaca oleracea* extract were investigated on acute alcoholic liver injury in 60 male Wistar rats. Results showed that *Portulaca oleracea* extract reduced serum levels of certain enzymes and triglycerides, increased antioxidant capacity, decreased inflammation, and improved lipid metabolism disorder induced by ethanol after seven days [122]. Another study evaluated the effect of feeding *Rumex patientia* L. seeds on serum glucose and lipid profile in streptozotocin-diabetic rats. Diabetic rats treated with *R. patientia* showed a significant reduction in serum glucose levels at the 2<sup>nd</sup> and 4<sup>th</sup> weeks compared to untreated diabetics. *R. patientia* also reduced lipid peroxidation in hepatic tissue, suggesting that this extract could improve glucose and lipid profiles, partly due to its attenuation

of lipid peroxidation in hepatic tissue [123]. The effects of *Saccharum officinarum* L. juice were studied on liver injury caused by the tuberculosis drug Isoniazid in mice. The group treated with Isoniazid and *S. officinarum* juice had decreased levels of liver enzymes. The group treated with juice also showed significant recovery in liver structure [124]. Another study investigated the protective effects of *S. officinarum* juice on paracetamol-induced liver damage in rats. Paracetamol caused liver damage in rats, as evidenced by increased enzyme levels, decreased antioxidant levels, and increased liver weight and volume. Treatment with *S. officinarum* juice reduced serum ALT, AST, ALP, and bilirubin, along with an increase in the antioxidant parameter MDA level [125]. The aqueous methanolic extract of *Viola odorata* was tested in mice with paracetamol-induced liver injury, which caused necrosis and inflammation. *V. odorata* restored elevated levels of serum hepatic enzymes and total bilirubin, and histopathological studies showed attenuation of hepatocellular necrosis and inflammation. High-performance liquid chromatography analysis revealed the presence of hepatoprotective flavonoids isorhamnetin and luteolin in the extract [126]. Another study examined the effects of aqueous and hydro-alcoholic extracts of *V. odorata* on liver function in diabetic rats. Results showed that the extracts reduced Kupffer cells, inflammation, and congestion in the liver tissue. Additionally, the extracts decreased the level of liver enzymes and serum glucose levels in diabetic rats [127]. The study evaluated the effects of total triterpenoids and total flavonoids from *Vitis vinifera* L on immunological liver injury induced by Bacille-Calmette-Guerin (BCG) and lipopolysaccharide (LPS) mice and found that various doses of triterpenoids and flavonoids reduced liver injury, decreased BCG/LPS-induced elevated liver index and spleen index, decreased hepatic nitric oxide and MDA content, increased liver homogenate ALT and AST levels, and restored hepatic SOD activity. The results suggest that the presence of triterpenoids and flavonoids in *V. vinifera* may have properties for the treatment of liver injury [128]. The wide variety of the medical plants of the current study have been used in traditional North African, Indian, Unani, and Uighur medicine as well as in Turkish and Jordanian folk medicine. Overall, these findings suggest that natural plant extracts have

potential therapeutic applications in the prevention and treatment of liver diseases. Further research is needed to understand the mechanisms underlying their hepatoprotective effects fully and to develop effective therapies based on these natural compounds.

## Declarations

### Ethics Approval and Consent to Participate

This article does not contain any studies with human participants or animals performed by any of the authors. The authors declare no conflicts of interest.

### Consent for Publication

All authors consent for publication of this article.

### Competing Interests

The authors declare no competing interests.

### Availability of Data and Material

The datasets generated during and/or analyzed during the current study are available from the corresponding author upon reasonable request.

### Funding

It had no financial support and was done with personal expenses.

### Authors' Contributions

The authors have contributed equally to the parts of the article.

## ACKNOWLEDGMENT

The authors would like to thank Mr. Okhovat and Ms. Eftekhari for their support and helpful comments on this article.

## REFERENCES

1. Yuan H., Ma Q., Ye L., Piao G. The Traditional Medicine and Modern Medicine from Natural Products. *Molecules*. 2016;21(5).
2. Rabizadeh, F., Mirian, M.S., Doosti, R., Kiani-Anbouhi, R. and Eftekhari, E. Phytochemical Classification of Medicinal Plants Used in the Treatment of Kidney Disease Based on Traditional Persian Medicine. *Evidence-Based Complementary and Alternative Medicine*, 2022, Article ID 8022599, 13 page
3. Hamedí A., Zarshenas M.M., Sohrabpour M., Zargaran A. Herbal medicinal oils in traditional Persian medicine. *Pharmaceutical biology*. 2013;51(9):1208-18.
4. Mollica A., Zengin G., Locatelli M., Stefanucci A., et al. Anti-diabetic and anti-hyperlipidemic properties of *Capparis spinosa* L.: In vivo and in vitro evaluation of its nutraceutical potential. *Journal of Functional Foods*. 2017;35:32-42.
5. Muria-Gonzalez M.J., Chooi Y.H., Breen S., Solomon P.S. The past, present and future of secondary metabolite research in the D othideomycetes. *Molecular plant pathology*. 2015;16(1):92-107.
6. Pang Z., Chen J., Wang T., Gao C., et al. Linking plant secondary metabolites and plant microbiomes: a review. *Frontiers in Plant Science*. 2021;12:621276.
7. Lee C.-H., Park S.-W., Kim Y.S., Kang S.S., et al. Protective mechanism of glycyrrhizin on acute liver injury induced by carbon tetrachloride in mice. *Biological and Pharmaceutical Bulletin*. 2007;30(10):1898-904.
8. Gao X., Wang Y., Li Y., Wang Y., et al. Hugianpian, a traditional chinese medicine, inhibits liver cancer growth in vitro and in vivo by inducing autophagy and cell cycle arrest. *Biomedicine & Pharmacotherapy*. 2019;120:109469.
9. Zhang L., Su S., Zhu Y., Guo J., et al. Mulberry leaf active components alleviate type 2 diabetes and its liver and kidney injury in db/db mice through insulin receptor and TGF- $\beta$ /Smads signaling pathway. *Biomedicine & Pharmacotherapy*. 2019;112:108675.
10. Nimrouzi M., Mahbodi A., Jaladat A.-M., Sadeghfard A., et al. Hijamat in traditional Persian medicine: risks and benefits. *Journal of evidence-based complementary & alternative medicine*. 2014;19(2):128-36.
11. Yilmaz B. Chemical constituents of *Ajuga chamaepitys* (L.) Schreb growing in Turkey by GC-MS method. *Int J Pharmacol*. 2019;6(3):108-12.
12. Khemkham A., Belhadj S., Meddour R., Kenmoku H., et al. HS-SPME-GC/MS analysis of 3lamiaceae plants: *Ajuga iva* (L.) Schreb., *Salvia verbenacal*. and *Thymus algeriensisboiss.* & Reut. *Journal of Fundamental and Applied Sciences*. 2020;12(2):700-11.
13. Lakshmi P., Rajalakshmi P. Identification of phyto components and its biological activities of aloe vera through the gas chromatography-mass spectrometry. *International research journal of pharmacy*. 2011;2(5):247-9.
14. Valiei M., Shafaghat A., Salimi F. Chemical composition and antimicrobial activity of the flower and root hexane extracts of *Althaea officinalis* in Northwest Iran. *Journal of medicinal plants research*. 2011;5(32):6972-6.
15. Canlı K., Yetgin A., Akata I., Altuner E.M. Antimicrobial activity and chemical composition screening of *Anacyclus pyrethrum* root. *Indian Journal of Pharmaceutical Education and Research*. 2017.
16. Awad H., Awda J.M., Abd-Alssirag M., Allaalfalahi D. GC-mass analysis of (*Apium graveolens*) leaf extracts obtained with aqueous and methanol extraction and study its antimicrobial activity. *Asian Journal of Microbiology, Biotechnology & Environmental Sciences*. 2019;21.
17. Bylka W., Szauffer-Hajdrych M., Matławska I., Goślińska O. Antimicrobial activity of isocytisoid and

- extracts of *Aquilegia vulgaris* L. Letters in applied microbiology. 2004;39(1):93-7.
18. Dhouioui M., Boulila A., Chaabane H., Zina M.S., et al. Seasonal changes in essential oil composition of *Aristolochia longa* L. ssp. *paucinervis* Batt.(Aristolochiaceae) roots and its antimicrobial activity. Industrial Crops and Products. 2016;83:301-6.
  19. Shah A.J., Gilani A.-H., Abbas K., Rasheed M., et al. Studies on the chemical composition and possible mechanisms underlying the antispasmodic and bronchodilatory activities of the essential oil of *Artemisia maritima* L. Archives of pharmacal research. 2011;34:1227-38.
  20. Zhigzhitzhapova S., Popov D., Pintaeva E.T., Radnaeva L., et al. Essential oil from *Artemisia sieversiana* Willd. and development of related oil-in-water emulsions. Pharmaceutical Chemistry Journal. 2017;51:388-90.
  21. Gibernau M., Favre C., Talou T., Raynaud C. Floral odor of *Arum italicum*. Aroideana. 2004;27:142-7.
  22. Kochmarov V., Marinov L., Kozuharova E., Hristova-Avakumova N., et al. Exploration of collagenase, cyclooxygenases, angiogenesis and free radical processes as the putative pharmacological targets of *Arum maculatum* L. Biotechnology & Biotechnological Equipment. 2020;34(1):126-34.
  23. Wilczewska A.Z., Ulman M., Chilmonczyk Z., Maj J., et al. Comparison of volatile constituents of *Acorus calamus* and *Asarum europaeum* obtained by different techniques. Journal of Essential Oil Research. 2008;20(5):390-5.
  24. Jadhav A., Bhutani K. Steroidal saponins from the roots of *Asparagus adscendens* Roxb and *Asparagus racemosus* Willd. 2006.
  25. Zhang H., Birch J., Pei J., Mohamed Ahmed I.A., et al. Identification of six phytochemical compounds from *Asparagus officinalis* L. root cultivars from New Zealand and China using UAE-SPE-UPLC-MS/MS: effects of extracts on H<sub>2</sub>O<sub>2</sub>-induced oxidative stress. Nutrients. 2019;11(1):107.
  26. SR J S.K. Screening of phytochemical and GC-MS analysis of some bioactive constituents of *Asparagus racemosus*. Screening. 2014;6(2):428-32.
  27. Sharma A., Rai P., Prasad S. GC-MS detection and determination of major volatile compounds in *Brassica juncea* L. leaves and seeds. Microchemical Journal. 2018;138:488-93.
  28. Popović-Djordjević J., Cengiz M., Ozer M.S., Sarikurkcu C. Calamintha incana: essential oil composition and biological activity. Industrial Crops and Products. 2019;128:162-6.
  29. Devki G.S., Sisodia R. Screening of the potential phytochemicals from the *capparis decidua* fruit extract using GC-MS. 2020.
  30. Altameme H.J.M. GC-MS and FTIR analysis Phytocomponents on different parts of *Capparis spinosa* L.(Capparidaceae) in Iraq. Journal of Chemical and Pharmaceutical Sciences. 2016;9(4):3269-82.
  31. Lieb V.M., Esquivel P., Castillo E.C., Carle R., et al. GC-MS profiling, descriptive sensory analysis, and consumer acceptance of Costa Rican papaya (*Carica papaya* L.) fruit purees. Food chemistry. 2018;248:238-46.
  32. Kapoor I.P.S., Singh B., Singh G., De Heluani C.S., et al. Chemistry and antioxidant activity of essential oil and oleoresins of black caraway (*Carum bulbocastanum*) fruits: Part 69. Journal of the Science of Food and Agriculture. 2010;90(3):385-90.
  33. Fang R., Jiang C.H., Wang X.Y., Zhang H.M., et al. Insecticidal activity of essential oil of *Carum carvi* fruits from China and its main components against two grain storage insects. Molecules. 2010;15(12):9391-402.
  34. Kazemi M. GC/MS analyses for detection and identification of antioxidant constituents of *Carum copticum* essential oil. Thai Journal of Agricultural Science. 2014;47(3):141-5.
  35. Sahaf B.Z., Moharrampour S., Meshkatsadat M.H. Chemical constituents and fumigant toxicity of essential oil from *Carum copticum* against two stored product beetles. Insect Science. 2007;14(3):213-8.
  36. Sahu L., Joshi P., Rout O., Sahu A. Phytochemical evaluation of *Celastrus paniculatus* seed oil extracted by a method used by 'Uraan' tribe of Chhattisgarh. Journal of Ayurveda and Integrated Medical Sciences. 2020;5(02):57-65.
  37. Singh R., Chahal K.K. *Cichorium intybus* from India: GC-MS profiling, phenolic content and in vitro antioxidant capacity of sequential soxhlet extracted roasted roots. Brazilian Archives of Biology and Technology. 2019;62.
  38. Al-Akhras M.-A.H., Aljarrah K., Al-Khateeb H., Jaradat A., et al. Introducing *Cichorium pumilum* as a potential therapeutic agent against drug-induced benign breast tumor in rats. Electromagnetic Biology and Medicine. 2012;31(4):299-309.
  39. Baruah A., Nath S.C., Hazarika A.K., Sarma T.C. Essential Oils of the Leaf, Stem Bark and Panicle of *Cinnamomum bejolghota* (Buch.-Ham.) Sweet. Journal of essential oil research. 1997;9(2):243-5.
  40. Bakirtzi C., Tsatalas P., Spanakis M., Kokkalou E. GC-MS analysis of volatile constituents of *Cornus mas* fruits and pulp. Journal of Essential Oil Bearing Plants. 2013;16(2):183-200.
  41. Singh S., Devi B. Estimation of phytoconstituents from *Citrullus colocynthis* (L.) schrad roots extract by GC-MS spectroscopy. International Journal of Science and Research. 2016;7.
  42. Najjaa H., Neffati M., Zouari S., Ammar E. Essential oil composition and antibacterial activity of different extracts of *Allium roseum* L., a North African endemic species. Comptes Rendus Chimie. 2007;10(9):820-6.



43. Boeing T., Tafarelo Moreno K.G., Gasparotto Junior A., Mota da Silva L., et al. Phytochemistry and pharmacology of the genus *Equisetum* (Equisetaceae): A narrative review of the species with therapeutic potential for kidney diseases. *Evidence-Based Complementary and Alternative Medicine*. 2021;2021.
44. Sharma R.J., Gupta R.C., Bansal A.K., Singh I.P. Metabolite Fingerprinting of *Eugenia jambolana* Fruit Pulp Extracts using NMR, HPLC-PDA-MS, GC-MS, MALDI-TOF-MS and ESI-MS/MS Spectrometry. *Natural Product Communications*. 2015;10(6):1934578X1501000644.
45. Judzentiene A. Chemical composition of leaf and inflorescence essential oils of *Eupatorium cannabinum* L. from Eastern Lithuania. *Journal of Essential Oil Research*. 2007;19(5):403-6.
46. Muthulakshmi A., Mohan V. GC-MS analysis of bioactive components of *Feronia elephantum* Correa (Rutaceae). *Journal of Applied Pharmaceutical Science*. 2012(Issue):69-74.
47. Afifi S.M., El-Mahis A., Heiss A.G., Farag M.A. Gas chromatography–mass spectrometry-based classification of 12 fennel (*Foeniculum vulgare* Miller) varieties based on their aroma profiles and estragole levels as analyzed using chemometric tools. *ACS omega*. 2021;6(8):5775-85.
48. Păltinean R., Mocan A., Vlase L., Gheldiu A.M., et al. Evaluation of Polyphenolic Content, Antioxidant and Diuretic Activities of Six *Fumaria* Species. *Molecules*. 2017;22(4).
49. Jameel M., Ali A., Ali M. New phytoconstituents from the aerial parts of *Fumaria parviflora* Lam. *Journal of advanced pharmaceutical technology & research*. 2014;5(2):64-9.
50. Mustafa A.M., Caprioli G., Maggi F., Vittori S., et al. Comparative analysis of the volatile profiles from wild, cultivated, and commercial roots of *Gentiana lutea* L. by headspace solid phase microextraction (HS–SPME) coupled to gas chromatography mass spectrometry (GC–MS). *Food analytical methods*. 2016;9:311-21.
51. Akhtar R., Shahzad A. Alginate encapsulation in *Glycyrrhiza glabra* L. with phyto-chemical profiling of root extracts of in vitro converted plants using GC-MS analysis. *Asian Pacific journal of tropical biomedicine*. 2017;7(10):855-61.
52. Singh S., Sharma P.C. Gas chromatography–mass spectrometry (GC–MS) profiling reveals substantial metabolome diversity in seabuckthorn (*Hippophae rhamnoides* L.) berries originating from different geographical regions in the Indian Himalayas. *Phytochemical Analysis*. 2022;33(2):214-25.
53. Hosokawa K., Fukunaga Y. Production of essential oils by flowers of *Hyacinthus orientalis* L. regenerated in vitro. *Plant Cell Reports*. 1995;14:575-9.
54. Smelcerovic A., Spitteller M., Ligon A.P., Smelcerovic Z., et al. Essential oil composition of *Hypericum* L. species from Southeastern Serbia and their chemotaxonomy. *Biochemical Systematics and Ecology*. 2007;35(2):99-113.
55. Schwob I., Bessiere J., Dherbomez M., Viano J. Composition and antimicrobial activity of the essential oil of *Hypericum coris*. *Fitoterapia*. 2002;73(6):511-3.
56. Morshedloo M.R., Ebadi A., Maggi F., Fattahi R., et al. Chemical characterization of the essential oil compositions from Iranian populations of *Hypericum perforatum* L. *Industrial Crops and Products*. 2015;76:565-73.
57. Kara N., Baydar H. Scent components in essential oil, resinoids and absolute of *Iris (Iris florentina* L.). *Anadolu Tarim Bilimleri Dergisi*. 2014;29(1):70.
58. Peris I., Blázquez M.A. Comparative GC-MS analysis of bay leaf (*Laurus nobilis* L.) essential oils in commercial samples. *International journal of food properties*. 2015;18(4):757-62.
59. Hussein H.M. DETERMINATION OF PHYTOCHEMICAL COMPOSITION AND TEN ELEMENTS CONTENT (CD, CA, CR, CO, FE, PB, MG, MN, NI AND ZN) OF *CARDARIA DRABA* BY GC-MS, FT-IR AND AAS TECHNIQUES. *International Journal of Pharma and Bio Sciences*. 2016;7(3):1009-17.
60. Cheriet T., Mancini I., Seghiri R., Benayache F., et al. Chemical constituents and biological activities of the genus *Linaria* (Scrophulariaceae). *Natural product research*. 2015;29(17):1589-613.
61. Rehman R.N.U., You Y., Yang C., Khan A.R., et al. Characterization of phenolic compounds and active anthocyanin degradation in crabapple (*Malus orientalis*) flowers. *Horticulture, Environment, and Biotechnology*. 2017;58:324-33.
62. Al-Qarawi K.K., Al-Obaidi H.M.R. Detection of the active compounds in the leaves of the Common mallow plant *Malva parviflora* L. using GC-MS and HPLC technology. *Kufa Journal For Agricultural Sciences*. 2018;10(4).
63. Jabbari H., Shendabadizad R. GC-MS analysis of essential oils of *Humulus lupulus*, *Malva Sylvestris* and thymus plants in water solvent. *Journal of Advanced Pharmacy Education & Research*. 2020;10.
64. Mou K.M., Parvin M.N., Dash P.R. *International Journal of Pharmacognosy and Pharmaceutical Research*.
65. Cudalbeanu M., Furdui B., Cârâc G., Barbu V., et al. Antifungal, antitumoral and antioxidant potential of the danube delta *nymphaea alba* extracts. *Antibiotics*. 2019;9(1):7.
66. Intirach J., Junkum A., Lumjuan N., Chaithong U., et al. Antimosquito property of *Petroselinum crispum* (Umbellifereae) against the pyrethroid resistant and susceptible strains of *Aedes aegypti* (Diptera: Culicidae). *Environmental Science and Pollution Research*. 2016;23:23994-4008.

67. Apetrei L.C., Spac A., Brebu M., Tuchilus C., et al. Composition, antioxidant and antimicrobial activity of the essential oils of a full grown tree of *Pinus cembra* L. from the Calimani mountains (Romania). *Journal of the Serbian Chemical Society*. 2013;78(1):27-37.
68. Aloui F., Baraket M., Jedidi S., Hosni K., et al. Chemical composition, anti-radical and antibacterial activities of essential oils from needles of *Pinus halepensis* Mill., *P. pinaster* Aiton., and *P. pinea* L. *Journal of Essential Oil Bearing Plants*. 2021;24(3):453-60.
69. Tumen I., Hafizoglu H., Kilic A., Dönmez I.E., et al. Yields and constituents of essential oil from cones of *Pinaceae* spp. natively grown in Turkey. *Molecules*. 2010;15(8):5797-806.
70. Andriana Y., Xuan T.D., Quy T.N., Tran H.-D., et al. Biological activities and chemical constituents of essential oils from *Piper cubeba* Bojer and *Piper nigrum* L. *Molecules*. 2019;24(10):1876.
71. Assimopoulou A., Papageorgiou V. GC-MS analysis of penta-and tetra-cyclic triterpenes from resins of *Pistacia* species. Part I. *Pistacia lentiscus* var. Chia. *Biomedical Chromatography*. 2005;19(4):285-311.
72. Assimopoulou A., Papageorgiou V. GC-MS analysis of penta-and tetra-cyclic triterpenes from resins of *Pistacia* species. Part II. *Pistacia terebinthus* var. Chia. *Biomedical Chromatography*. 2005;19(8):586-605.
73. Piccolella S., Nocera P., Carillo P., Woodrow P., et al. An apolar *Pistacia lentiscus* L. leaf extract: GC-MS metabolic profiling and evaluation of cytotoxicity and apoptosis inducing effects on SH-SY5Y and SK-N-BE (2) C cell lines. *Food and Chemical Toxicology*. 2016;95:64-74.
74. Saitta M., La Torre G.L., Potortì A.G., Di Bella G., et al. Polyphenols of pistachio (*Pistacia vera* L.) oil samples and geographical differentiation by principal component analysis. *Journal of the American Oil Chemists' Society*. 2014;91:1595-603.
75. Durgawale, T.P., Khanwelkar, C.C. and Pratik, P.D., 2018. Phytochemical analysis of *Portulaca oleracea* and *Portulaca Quadrifida* extracts using Gas Chromatography and Mass Spectrophotometry. *Asian J Pharm Clin Res*.
76. Tsao R., Zhou T. Interaction of monoterpenoids, methyl jasmonate, and Ca<sup>2+</sup> in controlling postharvest brown rot of sweet cherry. *HortScience*. 2000;35(7):1304-7.
77. Feduraev P., Skrypnik L., Nebreeva S., Dzhobadze G., et al. Variability of phenolic compound accumulation and antioxidant activity in wild plants of some *Rumex* species (Polygonaceae). *Antioxidants*. 2022;11(2):311.
78. Allaway Z., Sosa A. Chemical study in leaf and fruit of some species for *Populus* and *Salix* in Diwanayah governorate using gas chromatography-mass spectrometry (GS-MS). *Plant Archives*. 2019;19:102-11.
79. Shamspur T., Sheikhshoae I., Afzali D., Mostafavi A., et al. Chemical Compositions of *Salix aegyptiaca*. L. Obtained by Simultaneous Hydrodistillation and Extraction. *Journal of Essential Oil Bearing Plants*. 2011;14(5):543-8.
80. Javidnia K., Shokrollahi A., Kazemi M., Shahabipour S. Volatile composition of the essential oil *Salix excelsa* from Iran. *Research in Pharmaceutical Sciences*. 2012;7(5):765.
81. Affes S., Ben Younes A., Frikha D., Allouche N., et al. ESI-MS/MS analysis of phenolic compounds from *Aeonium arboreum* leaf extracts and evaluation of their antioxidant and antimicrobial activities. *Molecules*. 2021;26(14):4338.
82. Cabral C., Lemos M., Cavaleiro C., Cruz M., et al. Essential oil of *Seseli tortuosum* L. from Portugal: safety and anti-inflammatory potential evaluation. *Arabian Journal of Medicinal and Aromatic Plants*. 2015;1(1):31-43.
83. Hanifah A., Maharijaya A., Putri S.P., Laviña W.A., et al. Untargeted metabolomics analysis of eggplant (*Solanum melongena* L.) fruit and its correlation to fruit morphologies. *Metabolites*. 2018;8(3):49.
84. Giuliani C., Pellegrino R.M., Selvaggi R., Tani C., et al. Secretory structures and essential oil composition in *Stachys officinalis* (L.) Trevisan subsp. *officinalis* (Lamiaceae) from Italy. *Natural product research*. 2017;31(9):1006-13.
85. Teles A.M., Silva-Silva J.V., Fernandes J.M.P., Abreu-Silva A.L., et al. GC-MS characterization of antibacterial, antioxidant, and antitrypanosomal activity of *Syzygium aromaticum* essential oil and eugenol. *Evidence-Based Complementary and Alternative Medicine*. 2021;2021:1-12.
86. Jasim S.F., Baqer N.N., Alraheem E. Detection of phytochemical constituent in flowers of *Viola odorata* by gas chromatography-mass spectrometry. *Asian Journal of Pharmaceutical and Clinical Research*. 2018;11(5):262-9.
87. Petretto G.L., Mercenaro L., Urgeghe P.P., Fadda C., et al. Grape and wine composition in *Vitis vinifera* L. Cv. Cannonau explored by GC-MS and sensory analysis. *Foods*. 2021;10(1):101.
88. Chinonye I., Oze R., Lynda O., Nkwoada A., et al. Phytochemical and Gc/Ms Analysis of The Rhizome of *Zingiber officinale* Plant Grown In Eastern Part Of Nigeria. *African Journal of Biology and Medical Research*. 2016;1(1):43-54.
89. Song L., Zheng J., Zhang L., Yan S., et al. Phytochemical profiling and fingerprint analysis of Chinese Jujube (*Ziziphus jujuba* Mill.) Leaves of 66 cultivars from Xinjiang Province. *Molecules*. 2019;24(24):4528.
90. Huang K., Zhang P., Zhang Z., Youn J.Y., et al. Traditional Chinese Medicine (TCM) in the treatment of COVID-19 and other viral infections: Efficacies and mechanisms. *Pharmacology & therapeutics*. 2021;225:107843.

91. Ren J.L., Zhang A.H., Wang X.J. Traditional Chinese medicine for COVID-19 treatment. *Pharmacological research*. 2020;155:104743.
92. Fettach S., Mrabti H.N., Sayah K., Bouyahya A., et al. Phenolic content, acute toxicity of *Ajuga iva* extracts and assessment of their antioxidant and carbohydrate digestive enzyme inhibitory effects. *South African Journal of Botany*. 2019;125:381-5.
93. Abubakar A.M., Dibal N.I., Attah M.O.O., Chiroma S.M. Exploring the antioxidant effects of *Aloe vera*: Potential role in controlling liver function and lipid profile in high fat and fructose diet (HFFD) fed mice. *Pharmacological Research - Modern Chinese Medicine*. 2022;4:100150.
94. Bai L., Cui X., Cheng N., Cao W., et al. Hepatoprotective standardized EtOH–water extract of the leaves of *Ziziphus jujuba*. *Food & function*. 2017;8(2):816-22.
95. Shen X., Tang Y., Yang R., Yu L., et al. The protective effect of *Ziziphus jujuba* fruit on carbon tetrachloride-induced hepatic injury in mice by anti-oxidative activities. *Journal of Ethnopharmacology*. 2009;122(3):555-60.
96. AlSaid M., Mothana R., Raish M., Al-Sohaibani M., et al. Evaluation of the Effectiveness of *Piper cubeba* Extract in the Amelioration of CCl<sub>4</sub>-Induced Liver Injuries and Oxidative Damage in the Rodent Model. *BioMed Research International*. 2015;2015:359358.
97. Ali S., Prasad R., Mahmood A., Routray I., et al. Eugenol-rich Fraction of *Syzygium aromaticum* (Clove) Reverses Biochemical and Histopathological Changes in Liver Cirrhosis and Inhibits Hepatic Cell Proliferation. *Journal of cancer prevention*. 2014;19(4):288-300.
98. Selles C., Medjdoub H., Dib M.E.A., Zerriouh M., et al. Anti-diabetic activity of aqueous root extract of *Anacyclus pyrethrum* L. in streptozotocin-induced-diabetic rats. *Journal of medicinal plants research*. 2012;6(16):3193-8.
99. Shabbir U., Anjum I., Naveed Mushtaq M., Nasir Hayat Malik M., et al. Uroprotective and Hepatoprotective Potential of *Anagallis arvensis* against the Experimental Animal Model. *Journal of Tropical Medicine*. 2022;2022.
100. Sukketsiri W., Chonpathompikunlert P., Tanasawet S., Choosri N., et al. Effects of *Apium graveolens* extract on the oxidative stress in the liver of adjuvant-induced arthritic rats. *Preventive Nutrition and Food Science*. 2016;21(2):79.
101. Amat N., Upur H., Blažeković B. In vivo hepatoprotective activity of the aqueous extract of *Artemisia absinthium* L. against chemically and immunologically induced liver injuries in mice. *Journal of ethnopharmacology*. 2010;131(2):478-84.
102. Sunday R.M., Obuotor E.M., Kumar A. Antidiabetic Effect of *Asparagus adscendens* Roxb. in RIN-5F Cells, HepG2 Cells, and Wistar Rats. *Canadian Journal of Biotechnology*. 2019;3(1):132.
103. Shaban N.Z., El-Kot S.M., Awad O.M., Hafez A.M., et al. The antioxidant and anti-inflammatory effects of *Carica Papaya* Linn. seeds extract on CCl<sub>4</sub>-induced liver injury in male rats. *BMC Complementary Medicine and Therapies*. 2021;21(1):1-15.
104. Sangsefidi Z.S., Yarhosseini F., Hosseinzadeh M., Ranjbar A., et al. The effect of (*Cornus mas* L.) fruit extract on liver function among patients with non-alcoholic fatty liver: A double-blind randomized clinical trial. *Phytotherapy Research*. 2021;35(9):5259-68.
105. Vakiloddin S., Fuloria N., Fuloria S., Dhanaraj S.A., et al. Evidences of hepatoprotective and antioxidant effect of *Citrullus colocynthis* fruits in paracetamol induced hepatotoxicity. *Pak J Pharm Sci*. 2015;28(3):951-7.
106. Sharma P., Bodhankar S.L., Thakurdesai P.A. Protective effect of aqueous extract of *Feronia elephantum correa* leaves on thioacetamide induced liver necrosis in diabetic rats. *Asian Pacific journal of tropical biomedicine*. 2012;2(9):691-5.
107. Fatima S., Akhtar M.F., Ashraf K.M., Sharif A., et al. Antioxidant and alpha amylase inhibitory activities of *Fumaria officinalis* and its antidiabetic potential against alloxan induced diabetes. *Cellular and Molecular Biology*. 2019;65(2):50-7.
108. Mihailović V., Katanić J., Mišić D., Stanković V., et al. Hepatoprotective effects of secoiridoid-rich extracts from *Gentiana cruciata* L. against carbon tetrachloride induced liver damage in rats. *Food & function*. 2014;5(8):1795-803.
109. Abd-Al-Sattar Sadiq Layl L. Hepatoprotective effect of *Glycyrrhiza glabra* L. extracts against carbon tetrachloride-induced acute liver damage in rats. *Extracts Against Carbon Tetrachloride-Induced Acute Liver Damage In Rats (June 30, 2016) TJPRC: International Journal Of Veterinary Science, Medicine & Research (TJPRC: IJVSMR) Vol. 2016;1:1-8.*
110. Al-Razuqi R., Al-Jawad F., Al-Hussaini J., Al-Jeboori A. Hepatoprotective effect of *Glycyrrhiza glabra* in carbon tetrachloride-induced model of acute liver injury. *J Phys Pharm Adv*. 2012;2(7):259-63.
111. Geetha S., Jayamurthy P., Pal K., Pandey S., et al. Hepatoprotective effects of sea buckthorn (*Hippophae rhamnoides* L.) against carbon tetrachloride induced liver injury in rats. *Journal of the Science of Food and Agriculture*. 2008;88(9):1592-7.
112. Nawaz M.A., Aleem A., Hussain S.A., Manzoor M., et al. Hepatoprotective effect of methanolic extract of *Iris florentina* L. on paracetamol-induced liver toxicity in rats. *Pakistan Journal of Pharmaceutical Sciences*. 2022;35(5).
113. Gasparyan G., Tiratsuyan S., Kazaryan S., Vardapetyan H. Effect of *Laurus nobilis* extract on the functioning of liver against CCl<sub>4</sub> induced toxicity. *Journal of Experimental Biology and Agricultural Sciences*. 2015;3(2):174-83.

114. Feng J., Gao H., Yang L., Xie Y., et al. Renoprotective and hepatoprotective activity of *Lepidium draba* L. extracts on oxymetholone-induced oxidative stress in rat. *Journal of Food Biochemistry*. 2022;46(9):e14250.
115. Mohammed D.M., Elsayed N., Abou Baker D.H., Ahmed K.A., et al. Bioactivity and antidiabetic properties of *Malva parviflora* L. leaves extract and its nano-formulation in streptozotocin-induced diabetic rats. *Heliyon*. 2022;8(12):e12027.
116. Mallhi T.H., Abbas K., Ali M., Qadir M.I., et al. Hepatoprotective activity of methanolic extract of *Malva parviflora* against paracetamol-induced hepatotoxicity in mice. *Bangladesh Journal of Pharmacology*. 2014;9(3):342-6.
117. Ali S., Ansari K.A., Jafry M., Kabeer H., et al. *Nardostachys jatamansi* protects against liver damage induced by thioacetamide in rats. *Journal of ethnopharmacology*. 2000;71(3):359-63.
118. Bakr R.O., El-Naa M.M., Zaghloul S.S., Omar M.M. Profile of bioactive compounds in *Nymphaea alba* L. leaves growing in Egypt: Hepatoprotective, antioxidant and anti-inflammatory activity. *BMC Complementary and Alternative Medicine*. 2017;17(1):1-13.
119. Janakat S., Al-Merie H. Evaluation of hepatoprotective effect of *Pistacia lentiscus*, *Phillyrea latifolia* and *Nicotiana glauca*. *Journal of ethnopharmacology*. 2002;83(1-2):135-8.
120. Asadian S., Moallem S.A., Moshiri M., Mansouri M., et al. Evacuation the Effect of Single-and Multi-dose Administration of Ethanolic Extract of *Pinus eldarica* Pollen Against Acetaminophen-induced Rat Liver Injury. *International Journal of Basic Science in Medicine*. 2020;5(4):160-5.
121. Mavridis S., Gortzi O., Lalas S., Paraschos S., et al. Hepatoprotective effect of *Pistacia lentiscus* var. Chia total extract against carbon tetrachloride-induced liver damage in rats. *Planta Medica*. 2008;74(09):PA339.
122. Qiao J.-Y., Li H.-W., Liu F.-G., Li Y.-C., et al. Effects of *Portulaca oleracea* extract on acute alcoholic liver injury of rats. *Molecules*. 2019;24(16):2887.
123. Sedaghat R., Roghani M., Ahmadi M., Ahmadi F. Antihyperglycemic and antihyperlipidemic effect of *Rumex patientia* seed preparation in streptozotocin-diabetic rats. *Pathophysiology*. 2011;18(2):111-5.
124. Khan S.W., Tahir M., Lone K.P., Munir B., et al. Protective effect of *Saccharum officinarum* L.(sugar cane) juice on isoniazid induced hepatotoxicity in male albino mice. *Journal of Ayub Medical College Abbottabad*. 2015;27(2):346-50.
125. Patel B., Patel J., Raval B. Hepatoprotective activity of *Saccharum officinarum* L. against paracetamol induced hepatotoxicity in rats. *IJSPR*, 2010a. 2010;4(1):102-8.
126. Qadir M.I., Ali M., Saleem M., Hanif M. Hepatoprotective activity of aqueous methanolic extract of *Viola odorata* against paracetamol-induced liver injury in mice. *Bangladesh Journal of Pharmacology*. 2014;9(2):198-202.
127. Azari Z., Kherullahi Z., Mohammadghasemi F., Aghajany Nasab M., et al. Effect of the aqueous and hydro-alcoholic extracts of *Viola odorata* L. on biochemical and histologic liver parameters in diabetic Wistar rats. *Anatomical Sciences Journal*. 2020;17(1):21-32.
128. Liu T., Zhao J., Ma L., Ding Y., et al. Hepatoprotective effects of total triterpenoids and total flavonoids from *Vitis vinifera* L against immunological liver injury in mice. *Evidence-Based Complementary and Alternative Medicine*. 2012;2012.