



Are insecticidal seed treatments effective on overwintered adults of the Sunn pest, *Eurygaster integriceps* (Hemiptera: Scutelleridae)?

Aziz Sheikhiharjan^{id} & Ali Mohammadipour^{id}

Agricultural Entomology Research Department, Iranian Research Institute of Plant Protection, Tehran, Iran

✉ asheikhi48@gmail.com

^{id} <https://orcid.org/0000-0003-2462-3512>

✉ ali.mohammadipour@gmail.com

^{id} <https://orcid.org/0000-0002-8437-5242>

Abstract. Application of insecticides against overwintered adults of Sunn pest has limitations. To find the new measurement against the insect, two insecticides imidacloprid, and thiamethoxam, were tested as seed treatments against overwintered adults of Sunn pest in semi-field and field conditions in Alborz province. This project was carried out in a split-plot design with planting dates at 3 levels (at the beginning of wheat cultivation in 2016, Nov. 7, mid- and late-dates in 2016, Nov. 20, and Dec. 4) as the main plots and seed treatments at 5 levels, including imidacloprid 700WS with two concentrations of 1 and 0.5 g/kg of wheat seed and thiamethoxam 350FS with two concentrations of 0.8 and 0.4 g/kg and control (without seed treatment) as the sub-plots. The comparison of the mean efficiency of treatments at different cultivation dates in the semi-field conditions showed that the lowest (9.1%) and the highest (71.66%) efficiencies were observed at the beginning and late cultivation dates, respectively. In seed treatments, thiamethoxam at high concentration had the maximum efficacy (84.4%) 48 days after the last planting date, and the minimum efficacy (5.55%) was related to thiamethoxam at low concentration 116 days after. The results of field conditions showed that the highest (15%) and lowest (1.25%) efficacies of seed treatments were 97 days after the last cultivation date. Thiamethoxam seed treatment at high concentration had the highest efficiency (21.6%) among the treatments at the first sampling date. On other sampling dates, the efficiency of all treatments was less than 10%. The seed treatment with imidacloprid and thiamethoxam did not have acceptable effectiveness against overwintered adults and nymphs of Sunn pest. Also, the change in the cultivation date of seed-treated wheat cannot be effective in reducing the overwintered adult population of Sunn pest.

Keywords: imidacloprid, thiamethoxam, insecticide, chemical control, pest management

Citation: Sheikhiharjan, A. & Mohammadipour, A. (2024) Are insecticide seed treatments effective on overwintered adults of the Sunn pest, *Eurygaster integriceps* (Hemiptera: Scutelleridae)? *J. Entomol. Soc. Iran* 44 (2), 181–187.

Article History

Received:
23 November 2023

Accepted:
07 February 2024

Subject Editor:
Jahangir Khajecali

Introduction

Wheat is one of the oldest and most widely used crops in the world. This strategic crop has a very high nutritional value for humans, especially in terms of energy supply (carbohydrates). Wheat is cultivated extensively in different parts of the world. The cultivation area of irrigated and dry wheat in Iran is 5.4 million hectares, with a production of 13.3 million tons and an average irrigated yield of 4.3 tons per hectare (Ahmadi *et al.*, 2018).

Sunn pest, *Eurygaster integriceps* (Hemiptera: Scutelleridae), is the most important wheat pest in Iran. The infected areas range from southeastern Iran to southeastern Europe and North Africa. Sunn pest has one generation per year. Overwintering adults of Sunn pest spend the summer, autumn, and winter under the weeds of mountain slopes. When the average temperature reaches about 14 degrees Celsius, they migrate from mountains to fields. The migration period takes 7–30 days, 1–3 times. Overwintering adults travel 20–30 km. After settling in barley and wheat fields, they feed on leaves, stems, and final buds. In this stage, crop loss is quantitative. Then they mate and lay eggs. After one to 10 days, the nymphs hatch. The nymphs usually feed on the clusters. The crop loss of the new generation is in both quantity and quality, but quality damage to grain is very important because the product of wheat fields contains more than 2% of damaged grain, which is not recommended for bakery use (Khanjani, 2004; Javadipouya *et al.*, 2023). In Iran's weather conditions, the earliest and latest times of emigration from the mountains are in February and May. Based on phenological wheat fields, emigration starts at

Corresponding author Aziz Sheikhiharjan (Email: asheikhi48@gmail.com)



© 2024 by Author(s), Published by the Entomological Society of Iran

This Work is Licensed under Creative Commons Attribution-Non Commercial 4.0 International Public License.

the beginning of the pawing period and ends at cluster formation, and most of the emigration occurs when the wheat is in the pawing state (Rajabi, 2000). The emigration time of Sunn pest depends on the topography, altitude, rainfall, and the weather conditions of the area each year. The cold weather can cut off flying Sunn pests from mountains to fields (Rajabi, 2000).

Chemical control by seed treatment is not only effective against sucking pests but also reduces viral plant diseases including Maize dwarf mosaic virus (MDMV) and Barley yellow dwarf virus (BYDV). Seed treatment with thiamethoxam controlled *Rhopalosiphum padi* and decreased BYDV up to 50%. (Gourmet *et al.*, 1996a and 1996b). Preliminary studies have shown that neonicotinoids are effective on overwintering adults of Sunn pests (Jafarpour *et al.*, 2011).

Due to the lack of sufficient information about the migration behavior of overwintered adults on the fields, there is no optimized strategy for chemical control, and the chemical application on the overwintered adult stage often has less efficacy than on the nymphal stage. The seed treatment can be used in hot spots where Sunn pest infestation happens in those fields every year. It can increase the efficacy of the chemical control against Sunn pests and decrease the total treated area. Therefore, this study aimed to determine the effectiveness of chemical control by seed treatment methods with two insecticides, imidacloprid, and thiamethoxam, against overwintered adults of Sunn pest in laboratory and field conditions.

Materials and methods

Insecticides

Insecticides tested were imidacloprid (Gaucho® 700WS, Bayer Crop Science Inc., Germany) at two concentrations of 0.7 and 0.35 g a.i./kg of wheat seeds and thiamethoxam (Cruiser® 350FS, Syngenta Crop Protection, Greensboro, NC, USA) at two concentrations of 0.28 and 0.14 g a.i./kg of seeds that were used as a seed treatment (Gourmet *et al.*, 1996a; Way & van Emden, 2000).

Insect Collection

Overwintered adults of Sunn pest that had completed the diapause stage and were capable of mating and oviposition were collected from Ghara Aghaj Mountains at 2300 to 2200 meters above sea level from under Milk vetch and *Artemisia* plants in February and March 2016. It is located near Varamin, Tehran province. The collected insects were released into special plastic containers 15×30 and 20 cm high and then transferred to the laboratory and stored in the refrigerator at 3 ± 1 °C. They were immediately used to perform biometric tests on the same day or in less than 24 hours.

Laboratory Tests

The wheat cultivar Pishtaz was used for all experiments. On each of the planting dates, 6 pots were planted for each treatment, and 5 seeds were sowed in each pot. The bioavailability and efficacy of treatments were conducted against overwintered adults of Sunn pest at 48, 90, and 116 days after treatment. In each test, the number of overwintering adults was released in each pot, and the mortality was recorded 48 hours after exposure. The evaluation dates were the same as the field study.

Field Tests

Pishtaz cultivar was planted in experimental plots. The trial was performed in Alborz province under normal field conditions. The field location was at northern latitudes 35°48'38.9" and eastern longitudes 50°57'52.3". Winter wheat, Pishgam, was used for this study. The experiment was set in a split-plot design in a randomized complete block design. The main plot included planting dates in three levels with a 15-day interval: (1) beginning of planting date: 2016. Nov. 6; (2) middle of planting date: 2016. Nov. 19; (3) late planting date: 2016. Dec. 3. The sub-plots comprised insecticide seed treatment at 5 levels, including seed treatment by Gaucho® 700 WS at two concentrations of 1, 0.5 g/kg of wheat seeds, Cruiser 350 FS at two concentrations of 0.8, 0.4 g/kg of seeds, and control (without seed treatment).

Intervals between planting dates were at least 15 days. The block distance was considered to be 3 meters. The experiment was conducted with 15 treatments and three repetitions, and the size of the sub-plots was 50 square meters, which included 9 rows of 10 meters. Irrigation of each subplot was done separately to prevent water from entering from one subplot into the other.

The evaluation of seed treatments was conducted twice, at 25 and 61 Zadoks indexes, which are cereal growth staging scales. Cages with dimensions of 1.5 h× 0.5 w× 0.5 l meters were used in each plot, and 30 overwintered adults of Sunn pest were introduced in each cage. Mortality was recorded 2–4 days after release. The

first experiment was synchronized with the migration of the overwintered adults to fields at Zadoks 25, and in the second test at Zadoks 61 (flouring stage), 30 overwintering females and 10 males were released in each cage. In the second test, the cages did not move until harvesting time.

Statistical Analyses.

Data on mortality within treatments in laboratory conditions were analyzed using PROC GLM. Differences between treatments at each location were tested using the Fisher Least Significant Difference (LSD) Test (one-way ANOVA). The difference was considered significant at $P < 0.05$. The data were analyzed by the SAS version for Windows (SAS Institute, 2009).

Results

Laboratory Tests

Evaluation of the efficiency of seed treatment in autumn against the overwintered adults of Sunn pest with imidacloprid and thiamethoxam in pots showed that the interaction of planting dates and seed treatment was significant 48 and 90 days after the last planting date, while it was not significant 116 days later. (Table 1).

A comparison of the mean efficiency of seed treatments on different planting dates showed that the lowest efficiency of the treatments was related to the first planting date (9.1%) at 116 days after planting, and the highest efficiency was related to the second (61.66%) and third (71.66%) planting dates at 48 days after the last planting date. Thiamethoxam (84.4%) at a high concentration (0.8 g/kg seed) had statistically significant differences with the other treatments (Table 2).

Field Tests

Data analysis in the field conditions revealed that at all sampling times (97, 116, and 121 days after the last planting date), the interaction between planting dates and seed treatments was significant at the 5% level. Also, the efficacy of seed treatments with insecticides was significantly different at the 5% level. Thiamethoxam treatment (0.8 g/kg seed) had the highest efficiency (21.6%) at 97 days after the last planting date, and on the following sampling dates, the efficiency of all treatments was often less than 15% (Tables 3 and 4).

At 150 days after the last date, all seeds treated by neonicotinoid insecticides were not able to reduce the crop losses of Sunn pest in the economic injury level, all clusters of wheat were white in the cage, and the kernel rate was more than 70% (Table 5).

Table 1. The effect of sowing dates on the efficacy of the insecticide seed treatment on the overwintered Sunn pest on different days after the last planting date and the mean comparison based on LSD at laboratory conditions.

Sowing date in 2016	Efficiency (%) of planting date at days after last planting date		
	+48	+90	+116
1 st planting date (November 6)	17.5±7.7b	18.33±8.14a	9.1±3.5a
2 nd planting date (November 19)	61.66±8.1a	31.6±9.5a	18.33±7.42a
3 rd planting date (December 3)	71.66±8.2a	22.5±5.6a	15.83±4.16a
LSD (alpha 5%)	17.1	19.1	19

The means in a column followed by the same small letters are not significantly different.

Table 2. The mean effectiveness of the insecticide seed treatments against the overwintered Sunn pest on different days after the last planting date and the comparison based on LSD at the laboratory conditions.

Seed treatments (g/kg of seeds)	Efficiency (%) of the insecticides at different days after last planting date		
	+48	+90	+116
Gaicho [®] 700 WS(1)	41.11±10.46bc	11.11±3.09b	10±3.72b
Gaicho [®] 700 WS(0.8)	28.88±7.15c	7.77±2.22b	10±5.2b
Cruiser [®] 350 FS (0.8)	84.44±6.8a	61.11±8.57a	32.22±7.59a
Cruiser [®] 350 FS (0.4)	46.66±14.5b	16.66±6.45b	5.55±3.37b
LSD (alpha 5%)	13.77	15.12	16.41

The means in a column followed by the same small letters are not significantly different.

Discussion

The efficiency comparison of seed treatments in laboratory tests (pots) and field conditions showed that seed treatment in the pots has higher efficacy than in the field. Such that thiamethoxam at a high concentration had 84.44% efficacy in the pot and 21.6% efficacy in the field in the first sampling time, while the efficacy of imidacloprid at a high concentration was 41.11% and 5%, respectively. Seed treatments with imidacloprid and thiamethoxam insecticides did not have an acceptable efficacy (<10%) on overwintered adults of Sunn pest in the

subsequent evaluations in the field. However, seed treatment with imidacloprid and thiamethoxam is an effective chemical control method against wheat aphids and wheat bulb fly (Way & van Emden, 2000).

Seed treatment is the most important pest management technique. The rate of crop loss by wireworms, ground beetles (*Zabrus tenebrionis* Goege), and aphids in seed treatment methods is lower in comparison to conventional control methods (Adam & Noal, 2014). Growers often use neonicotinoids as seed treatments for controlling early-season pests on crops. It is used mostly in some crops, such as soybeans, corn, cotton, and wheat. The usage of insecticide seed treatment has significantly increased in the world since 2003. In the United States, insecticide seed treatment is applied on 40 million hectares (Douglas & Tooker, 2015).

Potato seed treatment with imidacloprid at a concentration of 30 g a.i./kg tubers can protect the plant from sucking pests for 30–38 days. Also, raising the insecticide concentration (60 g a.i./kg seed) can increase the duration of bio effectiveness (Nault *et al.*, 2004). In the present study, the results of wheat seed treatment by neonicotinoids against overwintered adults of Sunn pest showed that the efficacy was 84–28% at 48 days after planting in laboratory conditions and 7.77–61.6% at 90 days after planting, respectively, whereas in field conditions, the maximum efficacy of insecticide seed treatment was 21.6% at the migration time of overwintered adults to the fields (at 97 days after planting), and in the subsequent month it decreased to 9.2%. Based on our results and others, we found that the bio efficacy of insecticide seed treatment could not be persistent until 90 days after planting. The low percentage of efficiency of insecticide seed treatments would be due to the difference in feeding behavior of overwintered adults of Sunn pest and aphids (Rajabi, 2001; Rajabi, 2007).

Table 3. The effect of sowing dates on the efficacy of the insecticide seed treatment in planting dates against the overwintered adults of Sunn pest and the mean comparison based on LSD in field conditions.

Sowing date in 2016	Efficiency (%) of planting date at different days after last planting date		
	+97	+117	+121
1 st planting date (November 6)	1.25±1.11c	5.11±1.9a	7.73±1.18a
2 nd planting date (November 19)	10±3.3b	6.08±1.47a	5.7±1.2a
3 rd planting date (December 3)	15±3.19a	4±1.76a	7.17±1.8a
LSD (α 5%)	4.05	2.71	3.47

Table 4. The mean effectiveness of the seed treatments against the overwintered adults of Sunn pest on different days after the last planting date and comparison based on LSD in a field condition.

Seed treatments (g/kg of seeds)	Efficacy (%) of the insecticides at different days after last planting date		
	+97	+117	+121
Gaucht [®] 700 WS (1)	5±1.39bc	7.25±2.4a	8.39±1.5a
Gaucht [®] 700 WS (0.8)	6.66±2.84b	4.91±1.49ab	2.38±1.17b
Cruiser [®] 350 FS (0.8)	21.6±4.3a	5.12±1.9ab	9.2±2.1a
Cruiser [®] 350 FS (0.4)	1.66±0.7c	2.98±2b	7.53±0.95a
LSD (α 5%)	4.3	4.05	3.89

Table 5. The mean white head, numeric and weight kernel rates (Sunn pest damaged grain) of wheat in insecticide seed treatments on Sunn pest in the cages used in field.

Damage indexes	Seed treatments					LSD
	Gaucht [®] 700 WS (1)	Gaucht [®] 700 WS (0.8)	Cruiser [®] 350 FS (0.8)	Cruiser [®] 350 FS (0.4)	Control	
White head rate %	2.33±2.33	12.43±8.56	8.27±4.4	20.5±6.35	58.33±1.6	15.48
Numeric kernal rate %	77.17±23	82.23±17.66	69±31	100	100	61.86
Weight kernal rate %	77.17±23	77±23	68.9±31	100	100	64.81

The overwintered adults of Sunn pest feed from more parts of the wheat plant in comparison with aphids, which mostly select foliage ends for feeding. It can reduce the effectiveness of the seed treatment against the overwintered adult Sunn pest. The main reason for the low efficiency of seed treatment against overwintered adults may be the low concentration of neonicotinoids and their metabolites in the plant and their non-uniform distribution in different parts of the wheat plant. Since tested neonicotinoids are systemic, they can be absorbed through the roots and reach all parts of plant green tissues, nectarines, and pollen grains (Laurent & Rathahao, 2003; Schmuck *et al.*, 2001). The residue of thiamethoxam and its metabolites in different growth stages of sunflower plants showed that their concentrations were low in the flowering stage (2.7 ng/cm²), compared to the early growth stages (15.89 ng/cm²) (Bredeson & Lundgren, 2015). When imidacloprid is used as a seed treatment, most of its residue is found in the soil, so 23% of imidacloprid can be found in the soil at 97 days after planting (Westwood *et al.*, 1998). The highest amount of imidacloprid was observed in sugar beet plants at 64 days after

planting (5.3%), whereas at 97 days it reached 4.5% of the total residue on the seed (Westwood *et al.*, 1998). Imidacloprid may be metabolized for half or slightly more in the plant. Seed treatments of sugar beet (900 µg per seed) showed that the concentration of imidacloprid and its metabolites (olefin) in the sugar beet leaves was 15.2 µg/g at 21 days after planting, but it was reduced to 0.3-0.5 µg/g at 64 and 97 days after planting (Westwood *et al.*, 1998). The residue decrease of imidacloprid until the flowering stage was observed when it was used as chemigation (130 µg/L) in canola fields in the autumn (Reetz *et al.*, 2015). Plants uptake 5–6% of the total neonicotinoid residue in soil and seeds. The neonicotinoid concentration in soil reduces to 60% and 23% on 21 and 97 days after planting (Douglas & Tooker, 2015). The results of seed treatment with imidacloprid and thiamethoxam against overwintered adults of Sunn pest showed that the bio efficiency of this chemical control method is low. It can be due to the reduced concentration of imidacloprid and thiamethoxam and their metabolites in wheat plants at the emigration time of Sunn pest to wheat and barley fields. Seeds treated with neonicotinoids have been planted for many years. The concentration of neonicotinoid in the soil increased until 4-5 years, but it fixed from the sixth year onwards, so the mean concentration of clothianidin in the soil was 7 ng/g in 10 years after planting the treated seeds (Westwood *et al.*, 1998; Xu *et al.*, 2016).

In conclusion, seed treatment with neonicotinoids has a lower usage rate of insecticide per hectare than other chemical control methods. On the other hand, the results showed that the wheat seed treatment with the insecticides imidacloprid and thiamethoxam cannot control the wintered adult Sunn pest on winter planting dates, but it needs to be tested on spring planting dates of wheat. We hope that in the future, the new technology of formulations will increase insecticide efficacy against pests and reduce their environmental side effects. However, at present, the use of insecticide seed treatment on a large scale on overwintered adult Sunn pest does not have effectiveness and is not recommended.

Acknowledgments

We thank Dr. S. Shahrokhi and Dr. M. Amirmoafi for their financial and scientific assistance in project implementation at the research station for plant protection in Karaj, Alborz province. We also thank Ms. S. Gholami for providing cages and Ms. Nasreen Khosravani for the English-language editing of the article.

Funding

The research leading to these results received funding from the Iranian Research Institute of Plant Protection, Agricultural Research, Education and Extension Organization (AREEO) under Grant Agreement No: 2-16-16-95137.

REFERENCES

- Adam, N. M. & Noal, P. (2014) The development of a novel insecticide seed treatment for use in northern and central Europe for protection against soil dwelling and foliar pests in cereal crops. *Proceedings Crop Protection in Northern Britain*. 25 – 26 February, 2014, Scotland, 119-124. <https://www.cabidigitallibrary.org/doi/full/10.5555/20143314548>
- Ahmadi, K., Ebadzadeh, H., Abdshah, H., Kazemian, A. & Rafie, M. (2018) Agricultural Census of 2017–2018. Tehran, Iran: Ministry of Agriculture Jihad, Deputy of Planning and Economics, Information and Communication Technology Center. <https://www.jkgc.ir/fa/Page-346>
- Bredeson, M. M. & Lundgren, J. G. (2015) Thiamethoxam Seed Treatments Have No Impact on Pest Numbers or Yield in Cultivated Sunflowers. *Journal of Economic Entomology*, 108: 2665–2671. <https://doi.org/10.1093/jee/tov249>
- Douglas, M. R. & Tooker, J. F. (2015) Large-scale deployment of seed treatments has driven rapid increase in use of neonicotinoid insecticides and preemptive pest management in U.S. Field crops. *Environmental Science & Technology*, 49: 5088–5097. <https://doi.org/10.1021/es506141g>
- Gourmet, C., Kolb, F. L., Smyth, C. A. & Pedersen, W. L. (1996a) Use of imidacloprid insecticide to control barley yellow dwarf virus (BYDV) in oat and wheat. *Plant Disease Journal*, 80: 136-141. <https://www.proquest.com/openview/e22213fb2de49bce29c4fcc2632f4e80/1?pqorigsite=gscholar&cbl=18750&diss=y>
- Gourmet, C., Hewings, A. D., Kolb, F. L. & Smyth, C. A. (1996b) Effect of imidacloprid on nonflight movement of *Robopalosiphum Padi* and the subsequent spread of barley yellow dwarf virus. *Plant Disease Journal*, 78: 1098-1101. https://www.apsnet.org/publications/plantdisease/backissues/Documents/1994Articles/PlantDisease78n11_1098.PDF
- Jafarpour, A. A., Garjan, A. S., Imani, S., Mahjoub, S. M. & Mahmoudvand, M. (2011) Toxic effects of neonicotinoid insecticides on nymphs and adults of the Sunn pest, *Eurygaster integriceps* Puton (Hemiptera: Scutelleridae). *Acta Entomologica Sinica*, 54(8): 938-94. <http://www.insect.org.cn/EN/Y2011/V54/I8/938>

- Javadipouya, H., Valizadegan, O. & Sheikharjan, A.** (2023) 'Susceptibility of Sunn pest in overwintered adults and new generation to eight insecticide groups of IRAC', *Journal of Entomological Society of Iran*, 43(3): 247-257. doi: 10.61186/jesi.43.3.5
- Khanjani, M.** (2004) Field crop pests in Iran. Bu-Ali Sina University, (195), pp.1-719.
- Laurent, F. M. & Rathahao, E.** (2003). Distribution of [14C] imidacloprid in sunflowers (*Helianthus annuus* L.) following seed treatment. *Journal of Agricultural and Food Chemistry*, 51: 8005–8010. DOI: [10.1021/jf034310n](https://doi.org/10.1021/jf034310n)
- Nault, B. A., Taylor, A. G., Urwiler, M. Rabaey, T. & Hutchison, W. D.** (2004) Neonicotinoid seed treatments for managing potato leafhopper infestations in snap bean. *Crop Protection*, 23: 147–154. <https://doi.org/10.1016/j.cropro.2003.08.002>
- Rajabi, Gh.** (2000) Ecology of Cereal Sunn pests in Iran. Agricultural Research, Education, Extension and Organization Publication, Tehran, Iran.
- Rajabi, Gh.** (2001) Investigation on the downward migration of hibernating Sunn pest individuals from the altitudes to the cereal field in Varamin region. *Journal of Applied Entomology and Phytopathology*, 68 (1, 2): 107- 122.
- Rajabi, Gh.** (2007). Sunn pest Management Based on its Outbreaks' Key Factor Analysis in Iran. Agricultural Education Publications, Tehran, Iran. 163pp.
- Reetz, J. E., Schulz, W., Seitz, W., Spittler, M., Zühlke, S., Armbruster, W. & Wallner, K.** (2015) Uptake of neonicotinoid insecticides by water-foraging honeybees (Hymenoptera: Apidae) through guttation fluid of winter oilseed rape. *Journal of Economic Entomology*, 109(1): 31-40. <https://doi.org/10.1093/jee/fov287>
- SAS Institute Inc.** (1999) SAS/STAT User's Guide, Ver. 7. SAS Institute Inc.
- Schmuck, R., Schöning, R., Stork, A. & Schramel, O.** (2001) Risk posed to honeybees (*Apis mellifera* L., Hymenoptera) by an imidacloprid seed dressing of sunflowers. *Pest Management Science*, 57(3): 225-238. DOI: [10.1002/ps.270](https://doi.org/10.1002/ps.270)
- Way, M. & van Emden, H.** (2000) Integrated pest management in practice — pathways towards successful application. *Crop Protection*, 19: 81–103. [https://doi.org/10.1016/S0261-2194\(99\)00098-8](https://doi.org/10.1016/S0261-2194(99)00098-8)
- Westwood, F., Bean, K. M., Dewar, A. M., Bromilow, R. H. & Chamberlain, K.** (1998) Movement and persistence of [14C] imidacloprid in sugar-beet plants following application to pelleted sugar-beet seed. *Pesticide Science* 52(2), 97-103. [https://doi.org/10.1002/\(SICI\)1096-9063\(199802\)52:2<97::AID-PS687>3.0.CO;2-%23](https://doi.org/10.1002/(SICI)1096-9063(199802)52:2<97::AID-PS687>3.0.CO;2-%23)
- Xu, T., Dyer, D. G., McConnell, L. L., Bondarenko, S., Allen, R. & Heinemann, O.** (2016) Clothianidin in agricultural soils and uptake into corn pollen and canola nectar after multiyear seed treatment applications. *Environmental Toxicology and Chemistry*, 35(2): 311-321. DOI: [10.1002/etc.3281](https://doi.org/10.1002/etc.3281)

آیا روش بذر مال با مشره کش روی سن بالغ زمستان گذران (*Eurygaster integriceps* (Hemiptera: Scutelleridae) موثر است؟

عزیز شیخی گرجان^{id} و علی محمدی پور^{id}

بخش تحقیقات حشره شناسی کشاورزی، موسسه تحقیقات گیاه پزشکی کشور، تهران، ایران

✉ asheikhi48@gmail.com

✉ ali.mohammadipour@gmail.com

^{id} <https://orcid.org/0000-0003-2462-3512>

^{id} <https://orcid.org/0000-0002-8437-5242>

تاریخچه مقاله

دریافت: ۱۴۰۲/۰۹/۰۲ | پذیرش: ۱۴۰۲/۱۱/۱۸ | دبیر تخصصی: جهانگیر خواجه‌علی

هکیده

کنترل شیمیایی علیه سن مادر به روش محلول پاشی با محدودیت‌هایی همراه است. استفاده از روش ضدعفونی بذر گندم با غلظت مناسب در مناطقی که احتمال ریزش سن مادر در آن مزارع بیشتر است می‌تواند کارایی کنترل شیمیایی علیه سن مادر را بهبود بخشد. به همین خاطر برای تعیین میزان کارایی کنترل شیمیایی سن گندم به روش ضدعفونی بذر از دو حشره کش ایمیداکلوپراید (گائوچو) و تیمتوکسام (کرویزر) به روش ضدعفونی بذر علیه سن بالغ زمستان گذران (سن مادر) استفاده شد. آزمایش در استان البرز در شرایط مزرعه و گلخانه اجرا گردید. این پروژه به صورت اسپلیت پلات در قالب طرح بلوک‌های کامل تصادفی اجرا شد که فاکتور اصلی، تاریخ کشت در سه سطح (همزمان با شروع کشت گندم تاریخ ۱۳۹۵/۰۸/۱۶، اواسط تاریخ کشت گندم ۱۳۹۵/۰۸/۲۹، اواخر تاریخ کشت گندم ۱۳۹۵/۰۹/۱۳) و فاکتور فرعی آن تیمارهای ضدعفونی بذر در ۵ سطح شامل ضدعفونی بذر با حشره کش گائوچو 700WS با دو غلظت ۰/۵ گرم بر کیلوگرم بذر گندم و حشره کش کرویزر 350FS با دو غلظت ۰/۸ و ۰/۴ گرم بر کیلوگرم بذر گندم و شاهد (بدون ضدعفونی بذر گندم) بود. مقایسه میانگین درصد کارایی تیمارها در تاریخ‌های کشت مختلف در گلدان نشان داد که کمترین و بیشترین درصد کارایی تیمارها به ترتیب مربوط به تاریخ کشت اول (۹/۱٪) و آخر (۲۱/۶۶٪) است و در حشره کش تیمتوکسام با غلظت بالا، بیشترین کارایی (۸۴/۴٪) در ۴۸ روز و کمترین کارایی (۵/۵٪) در غلظت پایین آن حشره کش در ۱۱۶ روز بعد از آخرین تاریخ کاشت حاصل شد. بررسی داده‌ها در شرایط مزرعه نشان داد که بیشترین (۱۵/۱۶٪) و کمترین (۱/۲۵٪) درصد کارایی تیمارهای ضدعفونی بذر در ۹۷ روز بعد از آخرین تاریخ کشت بود. همچنین مقایسه تیمارهای ضدعفونی بذر از نظر درصد کارایی نشان داد که تیمار تیمتوکسام در غلظت بالا و در اولین نوبت نمونه برداری بیشترین کارایی را در میان تیمارها داشت (۲۱/۶٪) و در بقیه تاریخ‌های نمونه برداری درصد کارایی همه تیمارها کمتر از ۱۰ درصد بود. در مجموع، ضدعفونی بذر گندم با حشره کش‌های ایمیداکلوپراید و تیمتوکسام روی سن مادر و پوره کارایی قابل قبولی نداشت.

کلمات کلیدی: ایمیداکلوپراید، تیمتوکسام، حشره کش، کنترل شیمیایی، مدیریت آفت

نویسنده مسئول: عزیز شیخی گرجان (پست الکترونیک: asheikhi48@gmail.com)

Citation: Sheikhigarjan, A. & Mohammadipour, A. (2024) Are insecticide seed treatments effective on overwintered adults of the Sunn pest, *Eurygaster integriceps* (Hemiptera: Scutelleridae)? *J. Entomol. Soc. Iran* 44 (2), 181–187. <https://doi.org/10.61186/jesi.44.2.7>