

## A new sexual morph for *Neosetophoma iranianum*, and a key to *Neosetophoma* species

M. Mehrabi 

Department of Horticultural Science and Landscape Design, Shirvan Faculty of Agriculture, University of Bojnord, Bojnord, Iran

B. Asgari

Department of Botany, Iranian Research Institute of Plant Protection, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran

**Abstract:** *Neosetophoma iranianum* has so far only been described in its coelomycetous asexual morph. In this study, the sexual morph for this species was recovered for the first time, from dead branches of *Lonicera caprifolium*, in North Khorasan Province, Iran. The connection between the asexual and sexual morphs of the species is confirmed by phylogenetic analysis based on ITS rDNA sequence data, and morphological traits. The sexual and asexual morphs of *N. iranianum* are described and illustrated here, and compared with closely related species. A dichotomous key for the identification of all species assigned to the genus *Neosetophoma* is also provided.

**Keywords:** Biodiversity, *Phaeosphaeriaceae*, Phylogenetics, Ribosomal DNA, Saprobe, Taxonomy.

### INTRODUCTION

*Neosetophoma* Gruyter, Aveskamp & Verkley, a member of the family *Phaeosphaeriaceae* (*Pleosporales*), was introduced by de Gruyter et al. (2010) and typified by *N. samarorum* (Desm.) Gruyter, Aveskamp & Verkley based on *Phoma samarorum* Desm. This genus currently contains 28 species (Index Fungorum; <http://www.indexfungorum.org/Names/Names.asp>) which are saprobes, pathogens and endophytes on woody and herbaceous plants or in soil (Quaedvlieg et al. 2013, Phookamsak et al. 2014, Liu et al. 2015, Karunarathna et al. 2017, Wanasinghe et al. 2018, Marin-Felix et al. 2019, Hyde et al. 2020, Crous et al. 2021, Zhang et al. 2024). *Neosetophoma* species are mostly asexual morphs characterized by globose to irregular pycnidial conidiomata with papillate ostioles, and yellowish conidia that are attenuate at one end (de Gruyter et al. 2010, Marin-Felix et al. 2019).

*Neosetophoma Garethjonesii* Tibpromma, E.B.G. Jones & K.D. Hyde was reported as the first sexual morph of this genus (Tibpromma et al. 2017). Later, other species with sexual morphs, i.e. *N. camporesii* Q. Tian & K.D. Hyde, *N. guiyangensis* J.F. Zhang, J.K. Liu, K.D. Hyde & Z.Y. Liu, *N. miscanthi* A. Karun., C.H. Kuo & K.D. Hyde, *N. poaceicola* Goonas., Thambug. & K.D. Hyde, *N. shoemakeri* Senwana, Wanas., Bulgakov, E.B.G. Jones & K.D. Hyde, *N. trachycarpi* S.N. Zhang, K.D. Hyde & Jian K. Liu, and *N. xingrenensis* J.F. Zhang, J.K. Liu, K.D. Hyde & Z.Y. Liu were added to this genus (Hyde et al. 2018, 2019, 2020, Thambugala et al. 2017), from which only *N. shoemakeri* produces the asexual morph (Hyde et al. 2018). The sexual morph of *Neosetophoma* species is characterized by globose to subglobose, solitary, gregarious or scattered, unilocular, ostiolate ascospores; a carbonaceous peridium consisting of few layers of dark brown to brown pseudoparenchymatous cells of *textura angularis*; hamathecium comprising numerous, filiform, hyaline, pseudoparaphyses; bitunicate, fissitunicate, cylindrical, 8-spored asci; 0–3-seriately arranged ascospores, fusiform, smooth-walled, hyaline to brown, 1–3-septate, without gelatinous sheath (Thambugala et al. 2017, Hyde et al. 2018, 2019, 2020).

In an investigation of the biodiversity of microfungi in the North Khorasan Province of Iran, we found an unknown sexual morph on dead branches of *Lonicera caprifolium* L. Further morphological and molecular studies enabled the recognition of *Neosetophoma iranianum* Papizadeh, Amoozegar, Wijayaw., Shahz. Faz. & K.D. Hyde (Karunarathna et al. 2017), for which no sexual morph has been reported yet. Therefore, a new sexual morph for *N. iranianum* is described here using morphological and molecular data, and *Lonicera caprifolium* is reported as a new host for this species. We also present an identification key to all described species of *Neosetophoma*.


### MATERIALS AND METHODS

#### Morphological characterization


Samples were collected from dead branches of *Lonicera caprifolium* in the Shirvan Faculty of Agriculture in the North Khorasan Province of Iran

(37° 26' 11.36" N, 57° 49' 24.49"). Single ascospore cultures were obtained on potato dextrose agar (PDA, Merck, Germany), following the method of Senanayake et al. (2020). Macroscopic observations were carried out using a Nikon (SMZ-1B) stereo microscope. Colony growth and characteristics were determined on PDA and potato carrot agar (PCA; Domsch et al. 2007) incubated at 25 °C. Color names and codes used for descriptions were based on Rayner (1970). Microscopic slides were prepared in water.

Free-hand sections of ascomata were prepared and examined under Olympus CX21 light microscope. Photographs were taken using an AmScope digital camera (MD500). Herbarium specimen was preserved at the Fungus Reference Collection (IRAN...F) of Herbarium Ministerii Iranici Agriculturae "IRAN," Iranian Research Institute of Plant Protection (Tehran, Iran), while living culture at the Iranian Fungal Culture Collection (IRAN...C) of the "IRAN" Herbarium.



**Fig. 1.** RAxML tree based on the ITS sequence alignment of all *Neosetophoma* species. Maximum Likelihood bootstrap supports ( $\geq 50\%$ ) are shown above or below the branches. The branches with full statistical support (ML = 100%) are thickened. The scale bar shows the expected number of changes per site. The tree is rooted to *Poaeicola arundinis* MFLU 16-0158 and *P. bromi* MFLUCC 13-0739 (*Phaeosphaeriaceae*). T: ex-type strain.



**Fig. 2.** *Neosetophoma iranianum* IRAN 18640F (sexual morph). a. Ascomata on *Lonicera caprifolium* branch; b. Vertical section through an ascoma; c. Peridium; d. Pseudoparaphyses; e-g. Ascii; h. Apical apparatus of asci; i-l. Ascospores. Scale bars = (b) 100 μm; (c-g) 20 μm; (h) 10 μm; (i-l) 5 μm.

#### DNA extraction and sequencing


Fresh fungal mycelium was scraped from the surface of a PDA plate incubated at 25 °C for 7 d and transferred into a 1.5 mL centrifuge tube. Genomic DNA was extracted as described by Liu et al. (2000) with an initial step of grinding the mycelia in liquid nitrogen. The ITS region (ITS1-5.8S-ITS2) was amplified using primers ITS1/ITS4 (White et al. 1990). Amplification of D1–D2 of the 28S rDNA (LSU) was achieved using primers LROR/LR3 (Rehner and Samuels 1995). The PCR reaction (25

μL) contained 1 μL of each primer (10 pmol/μL, SinaClon Inc.), 1.0 μL of genomic DNA (30 ng/μL), 2.5 μL of 10× high yield PCR buffer (SinaClon, Iran), 0.3 μL of *Taq* polymerase (5 units/ μL, SinaClon, Iran), 1 μL of MgCl<sub>2</sub> (25 mM), 0.5 μL of dNTPs (10 mM), and 17.7 μL of sterile distilled water. PCR reactions were run on an Applied Biosystems 2720 Thermal cycler (Life Technologies, Singapore) with an initial denaturation step of 5 min at 95°C, followed by 35 cycles of 45 s at 94°C, 35 s at 58°C, 90 s at 72°C, and a final extension of 10 min at 72°C. PCR products were visualized in 1% agarose gel in 1×TBE



buffer. The PCR products were purified by Microsynth Company, Switzerland. The purified DNA samples were submitted for sequencing to a

capillary sequencing machine (ABI 3730XL, Applied Biosystem, Foster City, CA) of the same company.



**Fig. 3.** *Neosetophoma iranianum* IRAN 5290C (asexual morph). a. Colonies on PDA, incubated for two weeks at 25°C; b. Colonies on PCA incubated for two weeks at 25°C; c. Pycnidia produced on PCA; d. Pycnidium; e. Ostiole; f. Pycnidial wall; g, h. Conidiogenous cells; i. Conidia. Scale bars = (d) 100  $\mu$ m; (e, f) 20  $\mu$ m; (g-i) 10  $\mu$ m.

### Phylogenetic analysis

The chromatograms of the obtained sequences were analyzed using Finch TV version 1.40v (Geospiza). The sequences of a strain recovered in this study (IRAN 5290C) were aligned against the ex-type or reference strains of all described species of *Neosetophoma* (Table S1). The alignments were obtained using MAFFT v. 7 at the web server (<https://mafft.cbrc.jp/alignment/server/>) (Kuraku et

al. 2013, Katoh et al. 2019), using default settings. The alignment was edited manually where necessary with MEGA X (Kumar et al. 2018). Maximum Likelihood (ML) analysis including 1000 bootstrap replicates was performed in RAXMLGUI v. 1.5b1 (Silvestro and Michalak 2012), and the best scoring tree was selected among suboptimal trees from each replicate by comparing likelihood scores under the GTR+GAMMA substitution model. The tree was rooted with *Poaceicola arundinis* W.J. Li, Camporesi,

D.J. Bhat & K.D. Hyde (MFLU 16–0158) and *P. bromi* Wijayaw., W.J. Li, Camporesi, D.J. Bhat & K.D. Hyde (MFLUCC 13–0739), the *Phaeosphaeriaceae* (*Pleosporales*). Trees were visualized in FigTree version 1.4.4 (Rambaut 2012). The sequences generated in this study were deposited in GenBank (<http://www.ncbi.nlm.nih.gov/genbank>).

## RESULTS AND DISCUSSION

To elucidate the relationships of the newly discovered sexual morph, we conducted phylogenetic analyses using sequences of the ITS (483–563 bp) and partial LSU rDNA (526–857 bp) individually. Phylogenies of the LSU region (not shown) revealed that LSU sequences do not contain sufficient variation for distinguishing among species of *Neosetophoma*, and therefore this locus was excluded from our analyses. The ML analysis based on the ITS rDNA yielded a best-scoring tree (Fig. 1) with a final ML optimization likelihood value of  $-1917.529289$ . The matrix contained 154 distinct alignment patterns, with 4.98% of undetermined characters or gaps. Parameters for the GTR+GAMMA model were estimated as follows: base frequencies A = 0.225594, C = 0.243192, G = 0.228652, and T = 0.302562; substitution rates AC = 1.960977, AG = 5.029146, AT = 1.893325, CG = 0.611176, CT = 7.286687, and GT = 1.000000; gamma distribution shape parameter  $\alpha = 0.106773$ . Based on phylogenies of the ITS rDNA, our single ascospore culture (IRAN 5290C) was grouped with the ex-type strain of *Neosetophoma iraniana* (IBRC-M 30176), without known sexual morph, in a well-supported clade (98% BS). The ITS and LSU sequence comparison between the two strains, as recommended by Jeewon and Hyde (2016), also revealed that these strains were the same (no base pair difference). Moreover, the asexual morph produced in culture by strain IRAN 5290C aligns with the morphology described by Karunarathna et al. (2017). Therefore, the relevant description of *N. iraniana* by Karunarathna et al. (2017) is here expanded to include the new sexual morph.

## Taxonomy

*Neosetophoma iraniana* Papizadeh, Amoozegar, Wijayaw., Shahzadeh Fazeli & K.D. Hyde, *Mycosphere* 8(10): 1826 (2017)

**Sexual morph:** Ascospores 250–300  $\mu\text{m}$  high, 235–290  $\mu\text{m}$  diam ( $\bar{x} = 280 \times 265 \mu\text{m}$ ,  $n = 10$ ), solitary, scattered, immersed under epidermis, becoming erumpent through host surface, uniloculate, globose to subglobose, black, glabrous,

ostiolate. Peridium 36–63  $\mu\text{m}$  thick, consisting of several layers of dark brown to brown, pseudoparenchymatous cells of *textura angularis*, 3–8  $\mu\text{m}$  diam. Hamathecium comprising up to 200  $\mu\text{m}$  long, 2–3  $\mu\text{m}$  wide, numerous, filamentous, septate, pseudoparaphyses embedded in a gelatinous matrix. Asci 93–123  $\times$  9–11  $\mu\text{m}$  ( $\bar{x} = 112 \times 10.7 \mu\text{m}$ ,  $n = 20$ ), 8-spored, bitunicate, fissitunicate, cylindrical, short pedicellate, apically rounded with an ocular chamber. Ascospores 15–20  $\times$  6–10  $\mu\text{m}$  ( $\bar{x} = 17.3 \times 8.1 \mu\text{m}$ ,  $n = 30$ ), uniseriate, partially overlapping, fusiform to ellipsoidal, initially hyaline to subhyaline, becoming brown at maturity, 1–3-septate, mostly 3-septate, straight to slightly curved, deeply constricted at the septa, smooth-walled (Fig. 2, 3).

**Asexual morph** coelomycetous. Conidiomata (On PCA) 180–290  $\mu\text{m}$  high, 185–295  $\mu\text{m}$  diam ( $\bar{x} = 220 \times 210 \mu\text{m}$ ,  $n = 10$ ), pycnidial, solitary, dark brown, globose to subglobose, immersed to semi-immersed, unilocular, ostiolate, 15–20  $\mu\text{m}$  wide, with a 20–36 high  $\times$  30–44  $\mu\text{m}$  diam neck. Pycnidial wall 4–10  $\mu\text{m}$  thick, composed of 2–3 layers of light brown to brown, thick-walled cells of *textura angularis*, 3–9  $\mu\text{m}$  diam. Conidiophores reduced to conidiogenous cells. Conidiogenous cells 4–6  $\mu\text{m}$  high  $\times$  4–6  $\mu\text{m}$  wide, enteroblastic, phialidic, ampulliform, determinate, hyaline, smooth-walled. Conidia 5–8.5  $\times$  3–4  $\mu\text{m}$  ( $\bar{x} = 7.2 \times 3.7 \mu\text{m}$ ,  $n = 20$ ), ellipsoidal to tear-drop shape, sometimes attenuated at one end, (0–)1(–2) septate, continuous or constricted at the septa, initially hyaline to subhyaline, becoming light brown at maturity, smooth-walled, exuding as a black mass.

Cultural characteristics (25 °C, 4 weeks): Ascospores germinate on PDA within 24–48 h. Mycelium consists of 1.5–3.5  $\mu\text{m}$  wide, branched, septate, smooth, subhyaline hyphae. Colonies on PDA reach 75 mm diam, circular to irregular, fluffy, surface rough with hyphal tufts, edge entire, at first week white to buff (45), with sparse aerial mycelium, becoming pale grey after one month; reverse grayish sepia (106) in margin and black in center. Colonies on PCA reaching 70 mm diam, circular, velvety buff (45); the reverse of the same obverse colony color.

Specimen examined: IRAN, North Khorasan Province, Shirvan Faculty of Agriculture, 37° 26' 11.36" N, 57° 49' 24.49", on dead branches of *Lonicera caprifolium* L., 2 October 2024, M. Mehrabi, IRAN 18640F, IRAN 5290C.

Notes: *Neosetophoma iraniana* was introduced by Karunarathna et al. (2017) from soil in the Golestan Province of Iran. The asexual morph of our strain has a pycnidium shape, a conidium shape and septation, peridial structures, and

conidiogenous cells similar to those of *N. iranimum*. However, it produces larger pycnidia (180–290 × 186–296 µm vs 70–110 × 75–120 µm) and conidia (5–8.5 × 3–4 µm vs 4–6 × 2–4 µm) than the ex-type strain of *N. iranimum* (IBRC-M 30176). Nevertheless, there are some contradictions between the scale bars and pycnidial and conidial dimensions presented in the original description of *N. iranimum* by Karunarathna et al. (2017). Further examination of the ex-type strain of *N. iranimum* will help to better resolve the asexual morph of this species.

Phylogenetically, *N. iranimum* strains were placed in a large, moderately supported clade (84% BS) comprising ex-type strains of *N. aseptata*, *N. camporesii*, *N. clematidis*, *N. hnaniceana*, *N. lunariae*, *N. rosae*, *N. salicis* and *N. shoemakeri* (Fig. 1), from which only *N. camporesii*, lacks an asexual morph, and *N. shoemakeri* have sexual morphs. The sexual morph of *N. iranimum* is different from *N. camporesii* (Hyde et al. 2020) in asci size (62–90 × 7–10 µm) and ascospores size (20–32 × 3–5 µm), and from *N. shoemakeri* (Hyde et al. 2018) in ascomata size (170–230 × 170–200 µm) and ascospores size (15–27 × 3–6 µm). The asexual morph of *N. iranimum* can be distinguished from all the above-mentioned species by its septate conidia (aseptate in *N. aseptata* and *N. salicis*), shorter conidia (11–15 × 2–4 µm in *N. clematidis*, 8–9 × 3–4 µm in *N. hnaniceana*, 8–14 × 1.5–3 µm in *N. rosae*, 14–17 × 3 µm in *N. lunariae* and 7.5–10.5 × 2.5–3 µm in *N. shoemakeri*) and smaller pycnidia (120–180 × 120–160 µm in *N. shoemakeri*). Phookamsak et al. (2019) described *N. loniceriae* Phookamsak, Wanas. & K.D. Hyde as a coelomycetous fungus on dead hanging branches of *Lonicera maackii* in China. Morphologically, *N. loniceriae* is different from *N. iranimum* in conidiomata size (110–160 × 80–160 µm) and conidia size (9–12 × 4–5 µm).

### Key to the species of *Neosetophoma*

- |   |   |   |   |
|---|---|---|---|
| 1a. Fruiting bodies absent  | 2   | 5a. Ascomata 235–360 × 170–280 µm, with ostioles having a reddish tinge at the pores, ascospores 20–32 × 3–5 µm                             | <i>N. camporesii</i>                                      |
| 1b. Fruiting bodies present   | 3   | 5b. Ascomata with ostioles lacking a reddish tinge at the pores   | 6   |
| 2a. Colonies fluffy, with dense green spots in the center, hyphae 2–4.5 µm wide | <i>N. endophyticum</i> <sup>1</sup>                                   | 6a. Ascomata 250–300 × 235–290 µm, ascospores 15–20 × 6–10 µm; pycnidia 70–110(–290) × 75–120(–295) µm, conidia 4–6(–8.5) × 2–4 µm          | <i>N. iranimum</i>  |
| 2b. Colonies fluffy, later flocculent, hyphae 1.5–3.5 µm wide                   | <i>N. polygonata</i> <sup>1</sup> / <i>N. qimenensis</i> <sup>1</sup> | 6b. Ascomata 403–444 × 332–363 µm, ascospores 16–26 × 3–5 µm; asexual morph absent  | <i>N. garethjonesii</i>                                   |
| 3a. Sexual morph determined; coelomycetous asexual morph rarely present         | 4   | 7a. Ascospores mostly 3–5-septate   | 8   |
| 3b. Sexual morph undetermined; coelomycetous asexual morph always present       | 12  | 7b. Ascospores mostly 1–3-septate   | 10  |
| 4a. Ascomata more than 250 µm high  | 5   | 8a. Ascomata with darkened and yellow pigments around the ostioles  | <i>N. trachycarpi</i>                                     |
| 4b. Ascomata less than 250 µm high  | 7   | 8b. Ascomata without darkened area and yellow pigments around the ostioles  | 9   |
|   |   | 9a. Ascospores 20–25.5 × 3.5–4.5 µm, on woody plants  | <i>N. guiyangensis</i>                                    |
|   |   | 9b. Ascospores 18.5–22.5 × 3.5–5 µm, on grasses and woody plants  | <i>N. poaceicola</i>                                      |
|   |   | 10a. Asci sessile, 52 × 9 µm, ascospores 22 × 5 µm  | <i>N. xingrensis</i>                                      |
|   |   | 10b. Asci pedicellate   | 11  |
|   |   | 11a. Ascomata 170–230 × 170–200 µm, asci 84 × 9.5 µm, ascospores 15–27 × 3–6 µm; pycnidia 120–180 × 120–160 µm, conidia 7.5–10.5 × 2.5–3 µm | <i>N. shoemakeri</i>                                      |
|   |   | 11b. Ascomata 90–130 × 110–120 µm, asci 48 × 10.5 µm, ascospores 18–21 × 4.7–5.3 µm; asexual morph absent                                   | <i>N. miscanthi</i>                                       |
|   |   | 12a. Conidia aseptate   | 13  |
|   |   | 12b. Conidia septate  | 19  |
|   |   | 13a. Conidia 3–4 × 1–1.5 µm, on   | <i>Ficus elastica</i>                                     |
|   |   |   | <i>N. fici</i>  |
|   |   | 13b. Conidia wider than 1.5 µm, on other hosts  | 14  |
|   |   | 14a. Conidia longer than 8 µm   | 15  |
|   |   | 14b. Conidia shorter than 8 µm  | 16  |
|   |   | 15a. Conidia 8–9 × 2–3 µm, ellipsoidal to fusoid, on wood of <i>Buxus sempervirens</i>  | <i>N. buxi</i>  |
|   |   | 15b. Conidia 8–10 × 2–3 µm, allantoid, on twigs and branches of <i>Salix</i> sp.  | <i>N. salicis</i>   |
|   |   | 16a. Conidiogenous cells 2–3 × 1.5–2 µm, conidia 4–6 × 1.5–2.5, on <i>Rosa</i> sp.  | <i>N. rosigena</i>  |
|   |   | 16b. Conidiogenous cells longer than 3 µm, on other hosts   | 17  |
|   |   | 17a. Pycnidia 55–170 µm diam, conidiogenous cells 6–13 × 7–12 µm, conidia 4–7 × 1–2 µm, on <i>Ulmus</i> sp.                                 | <i>N. ulmi</i> (inval. nam.)                              |
|   |   | 17b. Pycnidia more than 170 µm diam   | 18  |
|   |   | 18a. Pycnidia 250–350 µm diam, conidiogenous cells 4–8 × 4–5 µm, conidia 4–5 × (1.5–)2 µm, on <i>Viburnum opulus</i>                        | <i>N. aseptata</i>  |
|   |   | 18b. Pycnidia 150–350 µm diam, conidiogenous cells 4–6 × 4–6 µm, conidia 6–8 × 2(–2.5) µm, on <i>Poaceae</i>                                | <i>N. cerealis</i> (= <i>N. phragmitis</i> ) <sup>2</sup> |
|   |   | 19a. Pycnidia 425–475 × 220–270 µm, conidia 11–15 × 2–4 µm, 3-septate   | <i>N. clematidis</i>                                      |
|   |   | 19b. Pycnidia less than 400 µm diam   | 20  |

- 20a. Conidia always 1-septate 21  
 20b. Conidia often multiseptate 22  
 21a. Conidia 7–10 × 3–4 µm, ellipsoidal to fusoid, on dead leaves and wood of *Buxus sempervirens*  
*N. hnaniceana*  
 21b. Conidia 5–10 × 2–3 µm, subcylindrical, on twigs of *Sambucus nigra* *N. sambuci*  
 22a. Conidia 5.5–8 × 1.5–2.5 µm, subcylindrical, 1–3-septate *N. rosarum*  
 22b. Conidia mostly longer than 8 µm 23  
 23a. Pycnidia more than 150 µm diam 24  
 23b. Pycnidia less than 150 µm diam 25  
 24a. Pycnidia 150–300 µm diam, conidia 10–22 × 2.5–3 µm, 1–4-septate *N. lunariae*  
 24b. Conidioma 250–280 µm diam, conidia 7–16 × 1–2 µm, 1–2-septate *N. samarorum*  
 25a. Pycnidia 100–120 µm diam, conidia 8–14 × 1.5–3 µm *N. rosae*  
 25b. Conidia wider than 3 µm 26  
 26a. Pycnidia 50–60 µm diam, conidia 6–11 × 3–4 µm, subcylindrical, fusiform, or ellipsoidal, 1–2-septate *N. italica*  
 26b. Pycnidia 80–160 µm diam, conidia 8.5–14 × 4–5 µm, ellipsoidal, 1–3-septate *N. lonicerae*

<sup>1</sup> It is difficult to distinguish these three species using morphological traits but they are phylogenetically distinct.

<sup>2</sup> Crous et al. (2020b) combined *Coniothyrium cerealis* E. Müll. into the genus *Neosetophoma* as *N. cerealis* (E. Müll.) Crous. They also, despite some variation regarding the conidial dimensions, synonymized *N. phragmitis* Crous, R.K. Schumacher & Y. Marin under *N. cerealis* based on phylogenies of the ITS rDNA. *Neosetophoma phragmitis* (Marin-Felix et al. 2019) has smaller conidia than *N. cerealis* (4–5 × 2 µm vs 6–8 × 2 µm). In this identification key, we follow Crous et al. (2020b) and treat *N. phragmitis* as a synonym of *N. cerealis*; nevertheless, based on the presented phylogeny in this study (Fig. 1) these species are phylogenetically distinct.

## ACKNOWLEDGMENTS

We are grateful to Dr. A.H. Pahlevani (Department of Botany, Iranian Research Institute of Plant Protection) for assistance in the identification of the host plants.

## REFERENCES

- Ahmadi, N., Arzanlou, M. and Narmani, A. 2021. Molecular phylogeny and morphology differentiate a new *Neosetophoma* species from Iran. *Nova Hedwigia*. 112(3-4):383-397
- Crous, P.W., Cowan, D.A., Maggs-Kölling, G., Yilmaz, N., Thangavel, R., et al. 2021. Fungal Planet description sheets: 1182–1283. *Persoonia*. 46:313-528
- Crous, P.W., Osieck, E.R., Shivas, R.G., Tan, Y.P., Bishop-Hurley, S.L., et al. 2023. Fungal Planet description sheets: 1478–1549. *Persoonia*. 50:158-310
- Crous, P.W., Wingfield, M.J., Chooi, Y.H., Gilchrist, C.L.M., Lacey, E., et al. 2020a. Fungal Planet description sheets: 1042–1111. *Persoonia*. 44:301-459
- Crous, P.W., Wingfield, M.J., Schumacher, R.K., Akulov, A., Bulgakov, T.S., et al. 2020b. New and Interesting Fungi. 3. *Fungal Systematics and Evolution*. 6:157-231
- De Gruyter, J., Woudenberg, J.H.C., Aveskamp, M.M., Verkley, G.J.M., Groenewald, J.Z. and Crous, P.W. 2010. Systematic reappraisal of species in *Phoma* section *Paraphoma*, *Pyrenochaeta* and *Pleurophoma*. *Mycologia*. 102:1066–1081.
- Domsch, K.H., Gams, W. and Anderson T-H. 2007. *Compendium of soil fungi*, 2nd edn. IHW Verlag, München 672 p.
- Gruyter, J. de., Woudenberg, J.H.C., Aveskamp, M.M., Verkley, G.J.M., Groenewald, J.Z. and Crous, P.W. 2010. Systematic reappraisal of species in *Phoma* section *Paraphoma*, *Pyrenochaeta* and *Pleurophoma*. *Mycologia*. 102(5):1066-1081
- Hernández-Restrepo, M., Schumacher, R.K., Wingfield, M.J., Ahmad, I., Cai, L., et al. 2016. *Fungal Systematics and Evolution: FUSE 2*. *Sydowia*. 68:193-230
- Hyde, K.D., Chaiwan, N., Norphanphoun, C., Boonmee, S., Camporesi, E., et al. 2018. *Mycosphere notes* 169–224. *Mycosphere*. 9(2):271-430
- Hyde, K.D., Dong, Y., Phookamsak, R., Jeewon, R., Bhat, D.J., et al. 2020. Fungal diversity notes 1151–1276: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity*. 100:5-277
- Hyde, K.D., Tennakoon, D.S., Jeewon, R., Bhat, D.J., Maharachchikumbura, S.S.N., et al. 2019. Fungal diversity notes 1036–1150: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity*. 96:1-242.
- Jeewon, R. and Hyde, K.D. 2016. Establishing species boundaries and new taxa among fungi: recommendations to resolve taxonomic ambiguities. *Mycosphere*. 7:1669-1677.
- Karunarathna, A., Papizadeh, M., Senanayake, I.C., Jeewon, R., Phookamsak, R., et al. 2017. Novel fungal species of *Phaeosphaeriaceae* with an asexual/sexual morph connection. *Mycosphere*. 8(10):1818-1834.
- Katoh, K., Rozewicki, J. and Yamada, K.D. 2019. MAFFT online service: Multiple sequence alignment, interactive sequence choice and

- visualization. Briefings in Bioinformatics. 20:1160-1166.
- Kumar, S., Stecher, G., Li, M., Knyaz, C. and Tamura, K. 2018. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution*. 35:1547-1549.
- Kuraku, S., Zmasek, C.M., Nishimura, O. and Katoh, K. 2013. aLeaves facilitates on-demand exploration of metazoan gene family trees on MAFFT sequence alignment server with enhanced interactivity. *Nucleic Acids Research*. 41:W22-W28.
- Li, W.J., Bhat, D.J., Camporesi, E., Tian, Q., Wijayawardene, N.N., et al. 2015. New asexual morph taxa in *Phaeosphaeriaceae*. *Mycosphere*. 6(6):681-708.
- Liu, D., Coloe, S., Baird, R. and Pedersen, J. 2000. Rapid mini-preparation of fungal DNA for PCR. *Journal of Clinical Microbiology*. 38:471
- Liu, J.K., Hyde, K.D., Jones, E.B.G., Ariyawansa, H.A., Bhat, D.J., et al. 2015. Fungal Diversity Notes 1–100: taxonomic and phylogenetic contributions to fungal species. *Fungal Diversity*. 72:1-197
- Marin-Felix, Y., Hernandez-Restrepo, M., Iturrieta-Gonzalez, I., García, D., Gene, J., et al. 2019. Genera of phytopathogenic fungi: GOPHY 3. *Studies in Mycology*. 94:1-124.
- Phookamsak, R., Hyde, K.D., Jeewon, R., Bhat, D.J., Jones, E.B.G., et al. 2019. Fungal diversity notes 929–1035: taxonomic and phylogenetic contributions on genera and species of fungi. *Fungal Diversity*. 95:1-273
- Phookamsak, R., Liu, J.K., McKenzie, E.H.C., Manamgoda, D.S., Ariyawansa, H.A., et al. 2014. Revision of *Phaeosphaeriaceae*. *Fungal Diversity*. 68:159–238.
- Quaedvlieg, W., Verkley, G., Shin, H.-D., Barreto, R., Alfenas, A., et al. 2013. Sizing up *Septoria*. *Studies in Mycology*. 75:307-390.
- Rambaut, A. 2012. FigTree version 1.4.0. Available from: <http://tree.bio.ed.ac.uk/software/figtree/> (accessed 15 May 2024).
- Rayner, R.W. 1970. A mycological colour chart. CMI and British Mycological Society, Kew
- Rehner, S.A. and Samuels, G.J. 1995. Molecular systematics of the *Hypocreales*: a teleomorph gene phylogeny and the status of their anamorphs. *Canadian Journal of Botany*. 73(Suppl. 1):S816-S823.
- Senanayake, I.C., Rathnayake, A.R., Marasinghe, D.S., Calabon, M.S., Gentekaki, E., et al. 2020. Morphological approaches in studying fungi: collection, examination, isolation, sporulation and preservation. *Mycosphere*. 11: 2678-2754.
- Silvestro, D. and Michalak, I. 2012. raxmlGUI: a graphical front-end for RAxML. *Organisms Diversity & Evolution*. 12:335-337.
- Thambugala, K.M., Wanasinghe, D.N., Phillips, A.J.L., Camporesi, E., Bulgakov, T. S., et al. 2017. Mycosphere notes 1–50: grass (*Poaceae*) inhabiting *Dothideomycetes*. *Mycosphere*. 8(4):697-796.
- Tibpromma, S., Hyde, K.D., Jeewon, R., Maharachchikumbura, S.S.N., Liu, J.K., et al. 2017. Fungal diversity notes 491–602: taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity*. 83:1-261.
- Vu, D., Groenewald, M., de Vries, M., Gehrman, T., Stielow, B., et al. 2019. Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in Mycology*. 92:135-154.
- Wanasinghe, D.N., Phukhamsakda, C., Hyde, K.D., Jeewon, R., Lee, H.B., et al. 2018. Fungal diversity notes 709–839: taxonomic and phylogenetic contributions to fungal taxa with an emphasis on fungi on Rosaceae. *Fungal Diversity*. 89:1-236
- White, T.J., Bruns, T., Lee, S. and Taylor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *In*: Innis, M.A., Gelfand, D.H., Sninsky J.J. and White T.J. (Eds.) *PCR protocols: a guide to methods and applications*. Academic Press, New York, pp. 315-322.
- Zhang, P.X., Han, Y.R., Chang, X.Y., Hu, J.X., Liu, Y.J., et al. 2024. Three new endophytic species of *Neosetophoma* in the roots of *Polygonatum cyrtonema*. *Mycosystema*. 43(5, no. 230358):1-13



## شکل جنسی جدید برای *Neosetophoma iranianaum*، همراه با یک کلید شناسایی برای گونه‌های *Neosetophoma*

مهدی محرابی<sup>۱</sup>، بیتا عسگری<sup>۲</sup>

۱- گروه علوم باغبانی و فضای سبز، دانشکده کشاورزی شیروان، دانشگاه بجنورد، بجنورد، ایران

۲- بخش تحقیقات رستنی‌ها، موسسه تحقیقات گیاه‌پزشکی کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران

**چکیده:** گونه *Neosetophoma iranianaum* تا کنون تنها در شکل غیرجنسی سلومیستی توصیف شده است. در مطالعه حاضر، شکل جنسی این گونه برای اولین بار از شاخه‌های مرده گیاه *Lonicera caprifolium* در استان خراسان شمالی گزارش می‌شود. ارتباط بین شکل غیرجنسی و جنسی این گونه از طریق واکاوی‌های تبارشناختی بر اساس ناحیه ITS از DNA ریبوزومی و ویژگی‌های ریخت‌شناختی تأیید شده است. در این مطالعه، تصاویر و توصیف‌های مربوط به شکل‌های جنسی و غیرجنسی گونه *Neosetophoma iranianaum* ارائه و با گونه‌های نزدیک مقایسه گردیده است. همچنین کلید شناسایی تمامی گونه‌های توصیف شده در جنس *Neosetophoma* ارائه شده است.

**کلمات کلیدی:** پوده‌زی، تاکسونومی، تبارشناسی، تنوع زیستی، DNA ریبوزومی، *Phaeosphaeriaceae*