

Expression and Purification of LigA Antigen, a Surface Lipoprotein in the Pathogenic *Leptospira interrogans* 1

2 Aida Chalesh¹, Pejvak Khaki^{2*}, Soheila Moradi Bidhendi², Majid Tebianian³, Morteza Taghizadeh
3 Tarnabi⁴

5

6 1 Department of Microbiology, North Tehran Branch, Islamic Azad University, Tehran, Iran. , ORCID:
7 <https://orcid.org/0009-0001-5850-0812>

8 2 Department of Microbiology, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and
9 Extension Organization (AREEO), Karaj, Iran, ORCID: <https://orcid.org/0000-0001-8839-1023>

10 3 Department of Immunology, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and
11 Extension Organization (AREEO), Karaj, Iran, ORCID: <https://orcid.org/0000-0002-9778-5048>

12 4 Department of Medical Vaccine, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and
13 Extension Organization (AREEO), Karaj, Iran, ORCID: <https://orcid.org/0000-0001-5456-4889>

16

17

18 Abstract

19 Although considerable progress has been made in leptospiral vaccine development, their use is limited
20 because of short-term and serovar-specific immunity. Thus far, many approaches have been used to identify
21 heterologous and cost-effective antigen(s) against the leptospirosis. Recent advances have identified
22 leptospiral immunoglobulin-like (Lig) proteins as promising candidates for vaccine development. Hence,
23 in this study, the recombinant LigA subunit consists of the ligA9, ligA10, ligA11, and ligA12 domains,
24 were selected as conserved regions of the LigA protein. Immunoinformatics approaches including I-
25 TASSER, ProSA, DiscoTope v2.0, and Molprobit were exploited to check the conformational accuracy.

26 Furthermore, 10 of the most efficient peptides for MHC-I and II grooves were predicted by the ElliPro
27 servers, NetMHCpan 4.1 EL and NetMHCIIpan 4.1 EL. The Ramachandran plot showed acceptable
28 conformations of the selected recombinant protein amino acid residues. The results showed that selected
29 epitopes elicit both humoral- and cell-mediated immune responses. Hence, the selected epitopes were
30 constructed in the pET41a⁺ plasmid and synthesized by General Biosystems. Recombinant plasmids were
31 transferred to *E. coli* Top10-DH5 α and BL21 StarTM (DE3) competent cells for cloning and expression,
32 respectively. Plasmid transformation and purification were confirmed using polymerase chain reaction and
33 enzymatic digestion. Recombinant LigA (r-LigA) was expressed in the presence of 0.5 M IPTG at 30^oC for
34 16 hours. The SDS-PAGE result revealed the production of 38-kDa protein, which accumulated mostly in
35 inclusion bodies and was purified using the urea method and dialysis. r-LigA protein dot blotting confirmed
36 a high degree of accuracy of immunogenicity. The present study revealed that r-LigA is a promising
37 candidate for developing diagnostic and subunit leptospirosis vaccines.

38 **Keywords:** Immuno-Informatic approach, Leptospirosis, LigA epitope, Recombinant protein

39 1. Introduction

40 Leptospirosis or Weil's disease is a zoonotic and a potentially serious bacterial disease caused by
41 pathogenic spirochetes of the genus *Leptospira*. It is transmitted to humans primarily through contact with
42 water or soil contaminated with the urine of infected animals, particularly rodents. Sporadic outbreaks have
43 occurred throughout the world, and it is particularly endemic in tropical and subtropical regions. Thus,
44 leptospirosis is an important public health concern in many countries (1).

45 One of the most effective strategies for combating leptospirosis is vaccination, which stimulates the immune
46 system by producing antibodies against the bacteria. The first successful vaccine against Leptospirosis was
47 reported in 1919, and it dramatically decreased the number of leptospirosis cases in Japan (2). Heat-killed
48 whole-cell vaccines (bacterin) against leptospirosis successfully protected coal miners in Kyushu, where
49 the disease was endemic in 1933. Live-attenuated and lipopolysaccharide (LPS) vaccine were another
50 alternative introduced vaccine. Major issues like lack of immunological memory, incomplete immunity,
51 reactogenicity, and serovar-restricted protection have led to a decrease in the use of these types of
52 vaccines. The drawbacks of these types of vaccines highlight the necessity of novel vaccine strategies (2-
53 4). In addition, with recent advances in molecular techniques and the availability of the complete *Leptospira*
54 genome sequence, the development of novel vaccines like recombinant proteins, using reverse
55 vaccinology has shown promising outcomes. Recent studies have highlighted the key role of outer
56 membrane proteins (OMPs) in pathogenesis through the communication between the leptospira and host
57 (5).

58 Highly immunogenic OMPs such as *Leptospira* subsurface lipoproteins (LipL32, LipL41), trans-membrane
59 outer membrane protein L1 (OmpL1), *leptospira* immunoglobulin-like protein (LigA, LigB, and LigC),
60 and leptospira OmpA-like protein (Loa22) as the main skeleton of recombinant vaccines have played a
61 major role in the development of new vaccines. Nevertheless, variation in serovar distribution in different
62 regions hinders the development of effective leptospira vaccines (4-6).

63 Structurally, LigA contains 13 domains of the bacterial immunoglobulin (Big) -like fold, and numerous
64 studies support its role in virulence. This protein promotes interactions with a range of specific host proteins
65 that mediate adhesion of *L. interrogans* to the extracellular matrix, inhibit hemostasis, and inactivate key
66 complement proteins to establish a systemic infection (7).

67 The development of vaccines against bacterial pathogens can be complex and time-consuming, and there
68 is always a risk of unforeseen complications or side effects. In some studies, *in silico* approaches were used
69 to identify potential epitopes on the LigA protein that could be used to develop vaccines or diagnostic tools
70 for leptospirosis. Moreover, by providing targeted protection against specific serovars of *Leptospira*, the
71 vaccine can reduce the incidence of leptospirosis and improve public health outcomes.

72 In terms of successful vaccine production, recombinant-protein production and purification present
73 challenges, particularly in terms of cost effectiveness, expression yield, and stability. On the other hand,
74 Lig proteins have been identified as suitable markers for the serodiagnosis of acute leptospirosis; thus, they
75 could be implemented in immunodiagnostic kits to address leptospirosis

76 (8). Accordingly, this study aimed to design and produce a recombinant LigA antigen in the prokaryotic
77 system to develop a recombinant leptospirosis vaccine.

78 **2. Materials and Methods**

79 This study was conducted from March 2021 to December 2021 at Razi Serum and Vaccine Research
80 Institute (Karaj, Iran).

81 **2.1 Recombinant LigA (r-LigA) immunogenic epitope sequence and properties prediction and** 82 **validations**

83 Previously designed conserved amino acid regions of LigA that stimulate the immune system were used in
84 this study (unpublished data). The sequence of the selected r-LigA protein consisted of amino acid positions

85 852-937 (ligA9: BID_2), 943-1029 (ligA10: Big_2), 1034-1119 (ligA11: Big_2), and 1125-1210 (ligA12:
86 Big_2).

87 The three-dimensional structure of the r-LigA protein was predicted by submitting the recombinant protein
88 sequence in I-TASSER (Iterative Threading ASSEMBly Refinement)
89 (<https://zhanglab.ccmb.med.umich.edu/I-TASSER/>) (9). Invasin protein of *Yersinia pseudotuberculosis* is
90 a well-characterized protein which belongs to immunoglobulin superfamily in structure and topology, and
91 its Big domain shares common themes to those observed in LigA (10). Hence, during threading for the
92 conformational annotation Invasin of *Yersinia pseudotuberculosis* protein (PDB ID: 1CWV) was used as
93 reference template for protein homology modeling (11). The structure with the minimum score in the
94 previous step was selected and downloaded in protein database (pdb) compatible file format, visualized by
95 Pymol, and used for further analysis. The prediction model quality was assessed using ProSA
96 (<https://prosa.services.came.sbg.ac.at>). The Ramachandran's plot was predicted using Molprobity
97 webserver (<http://molprobity.biochem.duke.edu/>).

98 **2.2 Linear and conformational B-Cell epitopes presence in r-LigA prediction**

99 To predict the linear (continuous) B-cell-epitopes present on r-LigA using the Ellipro module on the IEDB
100 server, the threshold was adjusted to 0.5. Conformational (discontinuous) B-cell epitopes were predicted
101 by another module, DiscoTope 3.0 (<https://services.healthtech.dtu.dk/services/DiscoTope-3.0/>), while the
102 input structure type set as default (Solved structure) and Calibrated score set as default (Moderate
103 confidence;0.90, recall up to ~50 %) (11).

104 **2.3 MHC class I and II peptides T-Cell epitopes presence in r-LigA prediction**

105 T-cell epitopes were evaluated by their binding capacity to major histocompatibility complex (MHC) I and
106 II peptides on the <http://tools.iedb.org/> server, and their locations were mapped on structural models. The
107 NetMHCpan 4.1 EL (recommended epitope predictor- 2023.09) method was used for MHC-I peptides. The
108 human-specific alleles HLA-A01:01, HLA-A02:03, HLA-A03:01, HLA-A11:01, HLA-A23:01, HLA-

109 B35:01, and HLA-B44:03 with a peptide length of 10 were selected (12). MHC-II peptides were predicted
110 using the NetMHCIIpan 4.1 EL (recommended epitope predictor- 2023.09) method. Human HLA-DR was
111 selected for prediction with alleles DRB1*01:01, DQA1*01:01/DQB1*02:01, and
112 DPA1*01:03/DPB1*04:01. A peptide length of 15 was selected (11).

113 **2.4 Cloning and expression of r-LigA**

114 r-LigA sequence synthesized between EcoRI and BamHI restriction sites of the pET41a⁺ plasmid (carrying
115 glutathione-S-transferase (GST-Tag), 6xHis, thrombin site, S-tag, enterokinase cleavage site and 8xHis tag
116 coding sequence as a *fusion partner* beside having the kanamycin resistance gene) by General Biosystems
117 (Anhui, USA). r-LigA – pET41a⁺ cloning vector transformed into the *Escherichia coli* Top10-DH5 α
118 competent cells prepared by the conventional MgCl₂ protocol. Transformations were performed using either
119 standard 42 °C heat shock protocol at 42°C and cultured on 2xYT agar (Merck, Germany) with the addition
120 of kanamycin (50 μ g mL⁻¹, K1377 sigma).

121 Random colonies were first screened by PCR using reverse T7 universal (Gene Link, Cat# 26-3000-13)
122 and forward T7 universal (Gene Link, Cat# 26-3000-13) primers. Bacterial amplified plasmids were
123 extracted using BioFact™ Plasmid Mini Prep Kit (Cat# PM105-100, South Korea) according to the
124 manufacturer's instructions, followed by restriction enzyme digestion with EcoRI and BamHI (Fermentas
125 Life Sciences, Thermo Fischer Scientific, Waltham, Massachusetts, USA) performed to confirm the
126 presence of the insert r-LigA gene in the pET41a⁺-LigA cassette before downstream transformation. To
127 achieve enhanced and better expression, extracted plasmids (500 ng) were transformed into *E. coli* BL21
128 Star™ (DE3) competent cells were screened using the heat shock method described above.

129 The induction-expression analysis of recombinant *E. coli* was conducted in the presence of 0.5 mM IPTG
130 in 2xYT broth at 30 °C for 16 hours with constant agitation. *E. coli* BL21 Star™ (DE3) without the plasmid
131 was used as a control strain.

132 **2.5 Purification of r-LigA and Blotting analysis of the r-LigA protein**

133 Transformed cells were harvested by centrifugation (5500 rpm for 5 min at 4°C), and the resulting bacterial
134 pellet was re-suspended in 10 mL of Phosphate Buffered Saline (PBS 1X), containing 10 µL of
135 Phenylmethanesulfonyl Fluoride (PMSF), and lysed with the aid of a sonicator (UP200St, Hielscher
136 Ultrasonics, Germany). Sonication was conducted 12 times by 1 min pulses with 1 min intervals in an ice
137 bath. The samples were then centrifuged at 15000 rpm for 15 min at 4°C, supernatants and pellets were
138 collected separately and then subjected to SDS-PAGE to compare their solubility and yield.

139 As r-LigA was expressed in the form of inclusion bodies, the protein was further purified under denaturation
140 conditions using serial concentrations of 1–8 M urea solutions in PBS buffer at 37°C. To this end, the pellets
141 were suspended in 5 mL of 1 M urea solution and shaken for 1 h, then it was centrifuged (5000rpm, 10
142 min), and supernatant discarded. This step was repeated with 2, 3, 4, and 5 M urea solutions. For 6–8 M
143 urea, the incubation time was increased overnight, and the supernatant was collected. In addition, after each
144 step, the efficiency of the purification was assessed using 10% SDS-PAGE. Fractions containing the
145 recombinant proteins were mixed and extensively dialyzed (MWCO 100 kDa, Thermo Fisher Scientific,
146 MA) against PBS (1:1000) overnight at 4°C, for 24 h. The Protein concentration was quantified using the
147 Bradford assay (13).

148 Purified r-LigA protein (approximately 200 ng) was used to charge the polyvinylidene fluoride membrane
149 (PVDF) (Hybond-P, Amersham Biosciences, UK) and allowed to dry. The membranes were then blocked
150 with 5% BSA in Tris-Buffered Saline-Tween 20 (TBST) for 40 min at RT. Consequently, the membrane
151 was washed in 1X TBST three times for 10 minutes, each with gentle rocking. The proteins were reacted
152 with conjugated anti-His tags antibodies (Thermo Fisher Scientific, USA) at RT for 1.5 hr. Finally, after
153 extensive washing with TBST once for 15 min and twice for 5 min, the reaction was visualized using 5-
154 thio-2-nitrobenzoic acid (TNB).

155 **3. Results**


156 **3.1 Recombinant LigA (r-LigA) immunogenic epitope structure and properties prediction and** 157 **validations**

158 The selected sequence of r-LigA consists of 359 amino acids (approximately 38 kDa). The 3D structure of
159 the protein was predicted using the I-TASSER server, which are represented in Table 1.

160 **Table 1.** Properties of modeled protein based on the ITASSER results

Parameters	Values
Number of amino acids	359
Molecular weight	37897.12= ~38 kDa
C-score	-0.62
Estimated TM-score	0.63±0.13
Estimated RMSD	8.0±4.4Å

161
162 The predicted three-dimensional structure of the r-LigA protein visualized using Chimera software
163 (v1.13.1, University of California, San Francisco, CA, USA) is depicted in Figure 1A. The majority of the
164 amino acid residues present in the protein may adopt β -sheet conformation. The overall quality of the
165 predicted structure of r-LigA using ProSA had a z-score of -6.37 (Figure 1B), which is well within the
166 range of native conformations, indicating that the predicted structure is reliable. According to MolProbity
167 scores, 92.2% (329/357) of all residues were Ramachandran-favored (i.e., without any steric clashes)
168 (Figure 1C).




170 **Figure 1.** Prediction and validation results of r-LigA protein. (A) Predicted 3D structure of r-LigA protein visualized by Pymol
171 software. (B) Overall model quality of the predicted structure given in a graphical form where X & Y axes represents residue
172 position & predicted

173 3.2 Linear and conformational B-Cell epitopes presence in r-LigA prediction

174 The graphical representation of amino acid scores as propensities for involvement in linear (continuous)
175 and conformational (discontinuous) B-cell epitopes is presented in Figure 2. According to the DiscoTopo

176 calibrated score with the default threshold (0.9), only 44 amino acids tended to be functional discontinuous
177 B-cell epitopes (Figure 2A and 2B). According to the Bepipred Linear Epitope Prediction 2.0 module; there
178 were 175 amino acid sequences above the threshold value (0.5) that are highlighted in yellow and
179 considered continuous B-cell epitope residues (Figure 2C).




180
181 **Figure 2. (A) Continuous epitope by DiscoTope: Residues with higher epitope**
182 **propensity are colored in a deeper red, while residues with lower epitope propensity**
183 **are colored in a deeper blue. (B) Residues interacting as continuous epitope**
184 **calculated by DiscoTope (C) Linear epitope: Amino acid residue with a score above**
185 **the threshold Bepipred Linear Epitope Prediction is considered to be part of an**
186 **epitope & coloured in yellow.**

187

188 The ElliPro prediction server identified 13 linear and three conformational potential peptide sequences as
 189 B-cell-epitopes (Figures 3 and 4). The linear B-cell epitopes range in length from 5 to 31 amino acids,
 190 whereas the three potential conformational B-cell epitopes comprise 57 and 87 amino acids, respectively.

191



192

193

194

195 **Figure 3.** Linear B cell epitopes predicted using ElliPro prediction server. Yellow highlight regions capable to stimulate B-cells.
 196 Peptide sequences, residue positions and predicted scores (in parenthesis) are given above each pictures.


197

198

199

200

4. Residues	No of residues	Score
-------------	----------------	-------



201

202

203

204

Figure 4. Conformational B cell epitopes predicted using Ellipro prediction server. Yellow highlight regions capable to stimulate B cells.

205 **4.1 MHC class I and II peptides T-Cell epitopes presence in r-LigA prediction**

206 Based on the IEDB analysis prediction results, there were various amino acid residues in the shape of the
 207 MHC-I and MHC-II grooves, and their most efficient peptides are presented with their scores in Tables 2
 208 and 3, respectively.

209 **Table 2.** MHC-I binding peptides predicted by the NetMHCpan EL 4.1.

No	Peptide Sequence	Allele	Position	Score
1	IVLNPTSSHK	HLA-A*03:01	5-14	0.892
2	ATYDSIKSNR	HLA-A*11:01	247-256	0.877
3	IVLNPTSSHK	HLA-A*11:01	5-14	0.758
4	FGDSEFTATY	HLA-A*01:01	240-249	0.747
5	SIEVTPNFF	HLA-B*44:03	90-99	0.739
6	ATYDSIKSNR	HLA-A*03:01	247-256	0.738
7	TTALSVGSSK	HLA-A*11:01	321-330	0.732
8	VLSEGLTLQL	HLA-A*02:03	186-195	0.713
9	DSMSASTTLY	HLA-A*01:01	164-173	0.689
10	LTDKGSAQQF	HLA-A*01:01	272-261	0.892

210

211

212 **Table 3.** MHC-II binding peptides predicted by the IEDB recommended 2.22 method.

No	Peptide Sequence	Allele	Position	Adjusted Rank *	Method Used
1	KSNRAWIFVNDEKLV	HLA-DQA1*01:01/DQB1*02:01	262-276	0.77	NetMHCIIpan
2	NRAWIFVNDEKLVNI	HLA-DQA1*01:01/DQB1*02:01	264-278	0.98	NetMHCIIpan
3	SNRAWIFVNDEKLVN	HLA-DQA1*01:01/DQB1*02:01	263-277	1.1	NetMHCIIpan
4	TTLYVTSVAVLIDIEV	HLA-DQA1*01:01/DQB1*02:01	175-189	1.1	NetMHCIIpan
5	DNTFSLAGSATAIDD	HLA-DRB1*01:01	144-158	1.3	Consensus (comb.lib./smm/nn)
6	IDNTFSLAGSATAID	HLA-DRB1*01:01	143-157	1.3	Consensus (comb.lib./smm/nn)
7	NTFSLAGSATAIDDG	HLA-DRB1*01:01	145-159	1.3	Consensus (comb.lib./smm/nn)
8	TLYVTSVAVLIDIEVK	HLA-DQA1*01:01/DQB1*02:01	176-190	1.4	NetMHCIIpan
9	IKSNRAWIFVNDEKLV	HLA-DQA1*01:01/DQB1*02:01	261-275	1.6	NetMHCIIpan
10	SIDNTFSLAGSATAI	HLA-DRB1*01:01	142-156	1.8	Consensus (comb.lib./smm/nn)


213 *Lower adjusted rank indicates a better binding.

214

215

216 **4.2 Cloning and expression of r-LigA protein**

217 Transformation of the r-LigA pET41a⁺ plasmid into the *E. coli* Top10-DH5 α (cloning vector) and *E. coli*
218 BL21 StarTM (DE3) (expression vector) produced several colonies in the presence of ampicillin. This was
219 confirmed by amplification of a 2000-bp segment in PCR of the target gene using T7 universal primers on
220 multiple random clones (Figure 5A). As shown in Figure 5B, EcoRI and BamHI digestion of the purified r-
221 LigA- pET41a⁺ plasmid yielded 5.9-kb and 1.1-kb fragments that can be attributed to the plasmid and r-
222 *LigA*, respectively.



223 **Figure 5. Transformation confirmation electrophoresis on 1% agarose gel. (M: Marker, NC:**
224 **Negative control)**


225 To achieve high-level expression of the peptide plasmid transformed into the *E. coli* BL21 StarTM (DE3)
226 induced with 0.5 mM IPTG at 30^oC for 16 hours which led to the expression of recombinant r-LigA protein

227 with approximately 71 KDa confirmed on SDS-PAGE. *E. coli* BL21 Star™ (DE3) without the plasmid was
228 subjected to the same experiment and did not express the protein under any conditions.

229 4.3 Purification and Blotting analysis of the r-LigA protein

230 Analysis of the cell lysate post-sonication revealed that the r-LigA r protein expressed using the pET41a+
231 vector in the *E. coli* BL21 system mostly accumulates as an insoluble form in the pellet, probably in
232 inclusion bodies. The molecular weight of r-LigA is approximately 38 KDa, whereas the different fused
233 partner proteins of the pET41a+ plasmid are approximately 32.29 KDa. Thus it is speculated that the
234 molecular weight of the expressed protein of this construct to be approximately 71 KDa. There was an
235 obvious band at 71 KDa in the purified protein after washing in serial concentrations of urea in PBS (1 to
236 5M) followed by solubilization (6 to 8M) at 37 °C. The outcome of this procedure on SDS-PAGE revealed
237 that r-LigA was successfully induced. Expressed and purified. The protein content gradually increased and
238 then decreased. The highest level of r-LigA gained in 6-M urea (Figure 6A).


239 To ensure high confidence, the Dot blot-binding assay was performed, and its result confirmed the
240 interaction of r-LigA with conjugated anti-His tags antibodies (Figure 6B).



241 **Figure 5.** Transformation confirmation electrophoresis on 1% agarose gel. (M: Marker, NC: Negative control)

242

243



244

245 **Figure 6.** SDS-PAGE of expressed r-LigA in *E. coli* (BL21 Star™ (DE3)) under different concentration of Urea after dialyze
246 (Marker: PM2500)

247 The quantity of the yield of purified recombinant protein was estimated to be 834 $\mu\text{g}/\text{mL}$ in Bradford assay.

248

249 5. Discussion

250 The global distribution of leptospirosis and its association with autoimmune disease have motivated
251 researchers to develop effective and safe vaccines. Commercial Leptospiral vaccines are available in many
252 countries for use in dogs, pigs, and livestock, most of which have limited efficacy because of a lack of
253 serovar specificity and their adverse effects, such as pain and fever (14, 15).

254 In fact, most surface proteins of pathogens are involved in virulence and even host immune response
255 stimulation, making them good targets for vaccine development (2). Matsunaga *et al.* (2003) expressed
256 surface-exposed proteins belonging to the bacterial immunoglobulin superfamily by pathogenic *Leptospira*
257 species and reported a new family of Big domain proteins called immunoglobulin-like proteins (Lig) in

258 pathogenic *Leptospira* (10). LigA, as a member of the Leptospiral outer membrane proteins, possess highly
259 conserved domains, which make it a proper candidate in antigen-based vaccine development (2, 16).

260 Kozumi *et al.* determined that recombinant full-length LigA and truncated LigB proteins can induce
261 immunoprotection against Leptospiral lethal infection in C3H/HeJ mice (17). Silva *et al.* expressed
262 carboxy-terminal region unique to the LigA protein (LigANI), LigBNI, and LigBrep from *Leptospira*
263 *interrogans* serovar Copenhageni in *E. coli* BL21(DE3)Star as recombinant proteins and in the hamster
264 infection model found that their potent activity of immunoprotection against mortality (18).

265 As it has been suggested that the presence of at least three Big domains (especially ligA11, and ligA12
266 specific Big) is critical, a construct of r-LigA protein carrying ligA9, 10, 11, and 12 domains inserted in
267 pET41a⁺ plasmid transformed into the *E. coli* Top10-DH5 α and *E. coli* StarTM (DE3) cells in this study
268 (19).

269 I-TASSER is an automated for protein tertiary structure prediction in which uses PSSpred for second
270 structure prediction, LOMETS for template detection, replica-exchange Monte Carlo simulation for
271 fragment structure assembly, SPICKER for model selection by clustering structure decoys, fragment-
272 guided molecular dynamics simulation (FG-MD) or ModRefiner for atomic-level structure refinement and
273 COACH for structure-based biology function annotation (9, 20). The I-TASSER server calculates the
274 confidence score (C-score), which estimates the quality of the predicted models based on the significance
275 of threading template alignments and the convergence parameters of the structure assembly simulations,
276 revealing high confidence in prediction. Normally, C-score values lie in between [-5 to 2], where a value
277 denotes a model with a higher confidence and correct topology (9). Then, a C-score value of -0.62 indicates
278 good topology of the modeled structure. The topological similarity of protein structures is reported as TM-
279 score in I-TASSER. TM-score <0.17 indicates a random similarity, while TM-score > 0.5 indicates a model
280 of correct topology (9). Based on these results, the TM-score 0.63 \pm 0.13 confirms a proper topology.

281 As a structural validation, the statistical distribution of the combinations of the backbone dihedral angles ϕ
282 and ψ of a protein can be determine using the Ramachandran plot. In good-quality homology models, very

283 few dihedral angles are found in forbidden regions (11). In this study, the three-dimensional structure
284 predicted for the r-LigA protein was reliable because only 7.8% of the residues were Ramachandran
285 outliers.

286 Epitopes are small specific segments of antigens that usually affect the specificity of cellular and humoral
287 immune responses. In linear epitopes, the primary amino acid sequence is recognized by the antibodies,
288 whereas in conformational epitopes, amino acids are not neighboring and are brought into close proximity
289 in the folded protein and directly bind to a receptor of the immune system(21, 22). Interestingly, the linear
290 epitopes with high scores as follow AIYGSIHSDSIDF (347-359),
291 ITISSQVLTDKGSAAQQFKAIGTFQGGSQLD (278-308), and SNSNDDRGL (326-334) composing a
292 conformational epitope with the highest score as well.

293 Binding to MHC class I molecules (MHC-I) plays a pivotal role in antigen presentation and induction of
294 cytotoxic T-cell responses. MHC class I and II genes in humans are called “*Human Leukocyte Antigen*”
295 (HLA). NetMHCpan 4.1 uses artificial neural networks (ANNs) and has been trained on a combination of
296 more than 850,000 quantitative binding affinity (BA) and mass spectrometry, Eluted Ligands (EL) peptides
297 that make it as one of the most reliable prediction servers(11, 23). Limited amount of data available for
298 most MHC class II molecules and considerable differences in sequence polymorphism and corresponding
299 details in the molecular structures across the different MHC class II loci hinder the development of cross-
300 species training strategies and is known as main limitation of pan-specific or cross species approach for
301 MHC class II prediction (11). The results for MHC-I binding peptides showed that epitope binding with
302 HLA-A*03:01allele had the highest binding affinity scores at 0.892.

303 Behera et al. applied similar approaches and found that, three out of the most efficient peptides for MHC-I
304 grooves predicted by NetMHCpan 4.1 server were presented by MHC Class II molecules, indicating that
305 both CMI and humoral immune response can be induced by those peptides (11).

306 The NetMHCIIpan 4.0 server was used to determine Helper T lymphocyte (HTL) epitopes (23). The
307 Number of epitope binding with the HLA-DQA1*01:01/DQB1*02:01 allele was higher than other alleles

308 in MHC-II binding peptides. Interestingly, NRAWIFVNDEKLV core sequence binding with HLA-
309 DQA1*01:01/DQB1*02:01 allele was present in top three NetMHCIIpan 4.0 prediction server, followed
310 by the NTFSLAGSATAIDD sequence binding with HLA-DRB1*01:01 in Consensus method.

311 For subunit vaccines, several expression systems were used, including bacteria, yeast, mammalian or plant
312 cells (24). In this study, the r-LigA construct transferred into *E. coli* Top10-DH5a and *E. coli* BL21 StarTM
313 (DE3). r-LigA was expressed in *E. coli* BL21 StarTM (DE3), it was observed that it is mostly in the form of
314 inclusion bodies. It has been reported that LigA and LigB normally contain a lipobox, whereas LigA
315 encodes a lipoprotein signal peptide distinct from the motifs of *E. coli* and other gram-negative bacteria.
316 Due to the presence of the lipobox, there is a *sec*-dependent export pathway for LigA exists across the
317 cytoplasmic membrane. Expression of LigA in *leptospira sp.* results in a surface protein with a signal
318 peptide subject to proteolytic removal of its lipid anchor. Therefore, a portion of LigA protein can be
319 recovered from the culture supernatant in the normal system(25).

320 Palaniappan et al. demonstrated that intact ligA without its signal sequence expression in *E. coli* was toxic
321 to *E. coli*, whereas the expression of a 90-kDa truncated LigA was not toxic to *E. coli* cells. They found
322 that complete LigA protein can be expressed only in leptospira-infected hamster kidney (26). Interestingly,
323 in a study LigA lipoproteins were expressed and exposed on the surface of the saprophyte *L. biflexa* cells
324 and suggested that *L. biflexa* is an appropriate surrogate host for the expression of at least some *L.*
325 *interrogans* outer membrane proteins (27).

326 Due to differences in leptospiral lipobox sequences, it is anticipated that leptospiral lipoproteins to not be
327 processed correctly in *E. coli* (27). To minimize protein inclusion bodies formation, it is recommended to
328 take some strategies such as changing vectors, host strains, production of endogenous chaperones or
329 /chaperone co-expression, low temperature induction, and using the target protein to a soluble protein or
330 peptide tags (28).

331 High levels of r-ligA (834 µg/mL) were purified from inclusion bodies with high efficiency by serial
332 washing with 6–8 molar urea. It can be concluded that r-LigA protein could be recovered from inclusion
333 bodies, using the urea method, which is a simple and low-cost technique. Hartwig *et al.*, (2010) expressed
334 rLigANI (61 kDa) in the eukaryotic expression system of *Pichia pastoris*, resulting in a significantly lower
335 protein yield (24).

336 In this study, the r-LigA construct produced an approximately 38-kDa recombinant protein, and high yield
337 of protein obtained at 6m urea solution. Urea purification method is a low-cost purification strategy in
338 comparison to Ni-NTA affinity chromatography purification. The optimum urea concentration in
339 purification procedure of this study were in accordance with a study focused on LipI41 recombinant protein
340 expression in *E. coli* BL21 (DE3) carrying pET32a+ expression vector (13)

341 r-LigA protein production was confirmed by immunoblotting analysis with HRP-conjugated Anti-6x His-
342 Tag antibodies. The performed dot-blot technique showed promising results in discriminating between
343 positive and negative serum samples.

344 Although expression of r-LigA was successful, due to the limited commercial availability of immunological
345 reagents for use in hamsters, we were not able to determine the cell-mediated immune response of hamsters
346 to r-LigA protein immunizations. Although this study focused on a comprehensive immunoinformatics, it
347 have not yet been assessed as vaccine candidates; and hence, could be worthy of further investigation as
348 novel vaccine candidates.

349 In conclusion, these findings suggest this immunoinformatics study represents novel vaccine candidates
350 that will further aid in the development of improved vaccines for leptospirosis. However, further refinement
351 of this technique is required before it can be used for leptospirosis diagnosis or vaccine development.

352

353 **Acknowledgment**

354 The authors would like to thank the Razi Vaccine and Serum Research Institute, Agricultural Research,
355 Education and Extension Organization (AREEO), Karaj, Iran, for providing laboratory space, facilities and
356 funding from the project source No. 12-18-18-106-96045-961025 on developing Leptospira vaccines.

357 **Authors' Contribution**

358 Conceptualization, Methodology, and Supervision: P. KH.

359 Formal analysis and investigation: All authors.

360 Drafting of the manuscript: A. C.

361 Writing - review and editing: All authors.

362 Funding acquisition: P. KH. And S. MB.

363 All authors checked and approved the final version of the manuscript.

364 **Ethics**

365 All the procedures for animal experiments were approved by the Research Ethics Committees of Islamic
366 Azad University- North Tehran Branch, under IR.IAU.TNB.REC.1401.049 Approval ID and performed in
367 accordance to the ethical principles and the national norms and standards for conducting Medical Research
368 in Iran guidelines.

369 **Conflict of Interest**

370 The authors declare that they have no known conflicts of interest or personal relationships that could have
371 appeared to influence the work reported in this paper.

372 **Data Availability**

373 The datasets generated during and/or analyzed during the current study are available from the corresponding
374 author upon reasonable request.

375

376 References

377

- 378 1. Breda LC, Hsieh C-L, Castiblanco Valencia MM, da Silva LB, Barbosa AS, Blom AM, et al. Fine
379 mapping of the interaction between C4b-binding protein and outer membrane proteins LigA and LigB of
380 pathogenic *Leptospira interrogans*. PLoS neglected tropical diseases. 2015;9(10):e0004192.
- 381 2. Bashiru G, Bahaman ARJJoMR. Advances & challenges in leptospiral vaccine development.
382 2018;147(1):15-22.
- 383 3. Dellagostin OA, Grassmann AA, Hartwig DD, Félix SR, da Silva EF, McBride AJ. Recombinant
384 vaccines against leptospirosis. Human vaccines. 2011;7(11):1215-24.
- 385 4. Koizumi N, Watanabe H. Leptospirosis vaccines: past, present, and future. Journal of
386 postgraduate medicine. 2005;51(3):210-4.
- 387 5. Haake DA, Zückert WRJL, Leptospirosis. The leptospiral outer membrane. 2015:187-221.
- 388 6. Barazzone GC, Teixeira AF, Azevedo BO, Damiano DK, Oliveira MP, Nascimento AL, et al.
389 Revisiting the development of vaccines against pathogenic *Leptospira*: innovative approaches, present
390 challenges, and future perspectives. Frontiers in Immunology. 2022;12:760291.
- 391 7. Haake DA, Matsunaga J. Leptospiral immunoglobulin-like domain proteins: Roles in virulence
392 and immunity. Frontiers in Immunology. 2021;11:579907.
- 393 8. Croda J, Ramos JG, Matsunaga J, Queiroz A, Homma A, Riley LW, et al. Leptospira
394 immunoglobulin-like proteins as a serodiagnostic marker for acute leptospirosis. 2007;45(5):1528-34.
- 395 9. Zhang YJBb. I-TASSER server for protein 3D structure prediction. 2008;9:1-8.
- 396 10. Matsunaga J, Barocchi MA, Croda J, Young TA, Sanchez Y, Siqueira I, et al. Pathogenic *Leptospira*
397 species express surface-exposed proteins belonging to the bacterial immunoglobulin superfamily.
398 2003;49(4):929-46.
- 399 11. Behera SK, Sabarinath T, Mishra PKK, Deneke Y, Kumar A, ChandraSekar S, et al.
400 Immunoinformatic study of recombinant liga/bcon1-5 antigen and evaluation of its diagnostic potential
401 in primary and secondary binding tests for serodiagnosis of porcine leptospirosis. 2021;10(9):1082.
- 402 12. Andreatta M, Nielsen MJB. Gapped sequence alignment using artificial neural networks:
403 application to the MHC class I system. 2016;32(4):511-7.
- 404 13. Golab N, Khaki P, Harzandi N, Esmaelizad M, Tebianian M. Expression and purification of the
405 LipI41, a surface-exposed lipoprotein antigen of pathogenic *Leptospira* spp. Veterinary Archives. 2020;
406 90(3):297-305.
- 407 14. Srivastava SJIJoMM. Supplement-Prospects of developing leptospiral vaccines for animals.
408 2006;24(4):331-6.
- 409 15. Techawiwattanaboon T, Patarakul KJAB. Update on molecular diagnosis of human leptospirosis.
410 2019;13(6):207-16.
- 411 16. Varma VP, Kadivella M, Kavela S, Faisal SM. *Leptospira* Lipid A Is a Potent Adjuvant That Induces
412 Sterilizing Immunity against Leptospirosis. Vaccines. 2023;11(12):1824.
- 413 17. Koizumi N, Watanabe H. Leptospiral immunoglobulin-like proteins elicit protective immunity.
414 Vaccine. 2004;22 (11-12):1545-52.
- 415 18. Silva EF, Medeiros MA, McBride AJ, Matsunaga J, Esteves GS, Ramos JG, et al. The terminal
416 portion of leptospiral immunoglobulin-like protein LigA confers protective immunity against lethal
417 infection in the hamster model of leptospirosis. Vaccine. 2007;25(33):6277-86.
- 418 19. McBride AJ, Cerqueira GM, Suchard MA, Moreira AN, Zuerner RL, Reis MG, et al. Genetic
419 diversity of the Leptospiral immunoglobulin-like (Lig) genes in pathogenic *Leptospira* spp. Infection,
420 Genetics and Evolution. 2009;9(2):196-205.

- 421 20. Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang YJNm. The I-TASSER Suite: protein structure and
422 function prediction. 2015;12(1):7-8.
- 423 21. Lin Xa, Sun A, Ruan P, Zhang Z, Yan JJBm. Characterization of conserved combined T and B cell
424 epitopes in *Leptospira interrogans* major outer membrane proteins OmpL1 and LipL41. 2011;11:1-9.
- 425 22. Fookolae SP, Talebshelimaki S, Rad MTS, Rokni MAJJoCBR. In silico prediction of B cell epitopes
426 of the hemolysis-associated protein 1 for vaccine design against leptospirosis. 2020;1(1):32-7.
- 427 23. Kapoor K, Patel A, Shekh S, Joshi CG. Designing of Multiepitope Subunit Vaccine Against
428 Leptospirosis Using Immunoinformatics Approach. 2022.
- 429 24. Hartwig DD, Oliveira TL, Seixas FK, Forster KM, Rizzi C, Hartleben CP, et al. High yield expression
430 of leptospirosis vaccine candidates LigA and LipL32 in the methylotrophic yeast *Pichia pastoris*. Microbial
431 cell factories. 2010;9:1-7.
- 432 25. Matsunaga J, Sanchez Y, Xu X, Haake DAJI, immunity. Osmolarity, a key environmental signal
433 controlling expression of leptospiral proteins LigA and LigB and the extracellular release of LigA.
434 2005;73(1):70-8.
- 435 26. Palaniappan RU, Chang Y-F, Jusuf S, Artiushin S, Timoney JF, McDonough SP, et al. Cloning and
436 molecular characterization of an immunogenic LigA protein of *Leptospira interrogans*. Infection
437 immunity. 2002;70(11):5924-30.
- 438 27. Figueira CP, Croda J, Choy HA, Haake DA, Reis MG, Ko AI, et al. Heterologous expression of
439 pathogen-specific genes *ligA* and *ligB* in the saprophyte *Leptospira biflexa* confers enhanced adhesion to
440 cultured cells and fibronectin. BMC microbiology. 2011;11:1-9.
- 441 28. Bhatwa A, Wang W, Hassan YI, Abraham N, Li X-Z, Zhou TJFiB, et al. Challenges associated with
442 the formation of recombinant protein inclusion bodies in *Escherichia coli* and strategies to address them
443 for industrial applications. 2021;9:630551.

444