

A Multi-Epitope Chimeric Hemagglutinin Vaccine Elicits Cross-Protective Immunity against Avian Influenza H5N8 and H9N2 Subtypes in Poultry

Zahra Bozorgkhoo¹, Behzad Hemmati¹, Morteza Taghizadeh², Majid Tebianian², Mostafa Ghaderi¹

1. Department of Microbiology, Karaj Branch, Islamic Azad University, Karaj, Iran

2. Department of Biotechnology, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Karaj, Iran

Abstract

Avian influenza viruses, notably H5N8 (HPAI) and H9N2 (LPAI), threaten public health due to zoonotic potential and genetic adaptability. Avian influenza viruses circulate in poultry and can sometimes directly infect humans after contact with infected birds. Some existing vaccines have proven useful in controlling these severe infections. Traditional vaccines targeting variable hemagglutinin (HA) head domains necessitate frequent updates. This study designed a chimeric HA (cHA H9/H5) vaccine targeting conserved epitopes from H5N8 and H9N2 HA proteins to elicit broad immunity.

A multi-epitope construct, integrating B-cell and cytotoxic T lymphocyte epitopes linked via stabilizing sequences (KK, AAY, GPGPG), was a codon-optimized construct, expressed in *E. coli* BL21(DE3), and purified (>95% purity, 22 mg/L yield). Specific pathogen-free chickens (n=20/group) received two doses of cHA H9/H5 formulated with Alum or Freund's adjuvants, compared to commercial H5N8/H9N2 vaccines. Humoral responses were assessed via hemagglutination inhibition (HI) assays.

The cHA H9/H5 vaccine induced robust HI titers against homologous H9N2 (log₂ GMT 10–12) and hetero-subtypic H5N8 (log₂ GMT 6–8), surpassing commercial vaccines. While H9N2 vaccines lacked cross-reactive H5N8 antibodies, cHA H9/H5 elicited neutralizing titers (20–80) against H5N8. Freund's adjuvant enhanced immunogenicity significantly, with sustained post-boost antibody levels.

These results highlight the cHA H9/H5 vaccine's ability to overcome strain-specific limitations by targeting conserved epitopes, inducing cross-reactive immunity critical for pandemic preparedness. Adjuvant selection proved pivotal in optimizing responses, aligning with prior chimeric HA vaccine research. This study advances the development of universal influenza vaccines, offering a promising strategy to mitigate risks posed by different evolving avian influenza variants.

Keywords: Avian Influenza, Chimeric vaccine, H5N8, H9N2

1. Introduction

The influenza virus, a member of the *Orthomyxoviridae* family, is classified into three subgroups (A, B, and C) based on the antigenic properties of its nucleoprotein and matrix protein(1). Among these, influenza type A is highly prevalent and exhibits greater pathogenicity compared to the other subgroups. This virus can infect various species of mammals and birds, potentially causing epidemics of varying intensities. Nearly all known subtypes of influenza A viruses have been found in birds. While numerous other animals can also act as hosts for influenza, historically, avian influenza has posed the greatest threat to human health(2, 3).

The influenza virus, particularly subtypes H5N8 (highly pathogenic avian influenza, HPAI) and H9N2 (low pathogenic avian influenza, LPAI), poses significant threats to both poultry and human health due to its potential for genetic reassortment and zoonotic transmission (4, 5). Existing commercial vaccines, such as inactivated H5N1 and H9N2 vaccines, have been employed to curb poultry outbreaks and reduce human exposure, yet their efficacy is challenged by viral mutations and limited cross-protection (6, 7). The immune response to influenza vaccines relies on eliciting neutralizing antibodies, primarily against HA, which can be assessed through the hemagglutination inhibition (HI) test—a standard measure of vaccine-induced protection, with HI titers ≥ 40 commonly indicating sufficient immunity in protection studies (8). By focusing on conserved HA epitopes, this multi-epitope vaccine seeks to enhance immunogenicity and provide broader, longer-lasting protection against evolving H5N8 and H9N2 strains, addressing gaps in current vaccination strategies (7, 9). However, mutations in the HA gene could potentially enable avian influenza viruses to bind human receptors, enhancing their transmissibility(10).

Simultaneous infections with H5N1 and H9N2 may result in genetic re-assortment, producing new subtypes with high pathogenicity in humans(11) . Consequently, the emergence of avian influenza viruses, particularly H5 and H9 subtypes, represents a dual threat to both the poultry industry and public health. The World Health Organization (WHO) lists these subtypes among the most significant pandemic threats. To mitigate risks, some countries have employed various poultry vaccines targeting H5N1 (HPAI) and H9N2 (LPAI) to protect the poultry industry and reduce direct human exposure to infected birds (12) . However, addressing the challenges of

increasing viral pressures and controlling outbreaks in poultry requires timely development of new vaccines. Such vaccines must account for international trade implications and aim to prevent the spread of viral diseases. Based on these insights, this study focuses on designing a multi-epitope influenza vaccine targeting highly conserved regions of the HA protein in H5N8 and H9N2 strains, which are prone to mutation over time.

This study aims to design a multi-epitope vaccine targeting conserved regions of the hemagglutinin (HA) protein in these strains, a key immunogenic component responsible for viral entry by binding to host sialic acid receptors (9).

2. Materials and methods

2.1. Recombinant construct

The recombinant construct was developed in accordance with our previous report (13). Briefly, the target multi-epitope HA recombinant protein was constructed by selecting antigenic epitopes capable of inducing robust immune responses. B-cell and cytotoxic T lymphocyte (CTL) epitopes were connected linearly using specialized linkers forming a multi-epitope vaccine construct. The design was verified through various bioinformatics tools to confirm its structural and functional properties.

Key evaluations included allergenicity and antigenicity prediction using AllerTOP v. 2.0 and VaxiJen v. 2.0, respectively, to ensure the construct was non-allergenic and immunogenic. Physicochemical properties, such as molecular weight, isoelectric pH, hydropathicity, and stability, were analyzed with ProtParam. Secondary structures, including alpha helices and beta sheets, were modeled with PSIPRED v. 3.3, and tertiary structure prediction was completed using the SwissModel server. The pET-41a+ vector was chosen for expression due to its efficiency and high yield in *E. coli*.

2.2. Expression and purification of recombinant protein

The recombinant pET-41a (+)/HA construct was transformed into *Escherichia coli* BL21 (DE3) cells, optimized for protein expression. Cultures of the transformed *E. coli* were grown in Luria-Bertani (LB) medium supplemented with kanamycin to select for cells harboring the plasmid. Protein expression was induced at the mid-log phase of bacterial growth (OD₆₀₀ ≈ 0.6) by the addition of isopropyl β-D-1-thiogalactopyranoside (IPTG) at a final concentration

of 1 mM. The cultures were then incubated for an additional 4 hours at 37°C with constant shaking to optimize protein production.

Following induction, the cells were harvested by centrifugation and lysed using a combination of lysozyme treatment and sonication. The lysate was clarified by centrifugation to remove cell debris, and the supernatant containing the soluble recombinant chimeric HA protein (cHA H9/H5) was collected.

Purification of the cHA H9/H5 protein was performed using nickel-affinity chromatography. The clarified lysate was applied to a nickel-nitrilotriacetic acid (Ni-NTA) agarose column pre-equilibrated with binding buffer (50 mM NaH₂PO₄, 300 mM NaCl, 10 mM imidazole, and pH 8.0). After washing to remove non-specifically bound proteins, the HA protein was eluted using a high concentration of imidazole (250 mM) in the elution buffer.

The purity and identity of the cHA H9/H5 protein were confirmed by SDS-PAGE and Western blotting using anti-HisTaq antibodies.

2.3. Immunization and experimental infection of chickens

7day-old SPF White Leghorn chickens (Venky Lab Co., Ltd., Pune, India) were randomly allocated into five experimental groups (n=20 per group) and immunized with antigen-adjuvant formulations as delineated in Table 1. cHA H9/H5 protein was prepared at a concentration of 1 mg/ml, with each chicken receiving 30 µg of antigen diluted in PBS. Freund's adjuvant (complete for the initial dose and incomplete for the subsequent dose), and alum were employed as adjuvants in different groups, prepared in specified ratios with the antigen. Two commercial vaccine groups received inactivated influenza antigens derived from the H9N2 and H5N8 strains formulated by ISA 71 adjuvant (Seppic, France) : Influenza A virus (A/chicken/Iran/B308B/2019(H9N2) ARAK/2009(H9N2)), and Influenza A virus (A/Poultry/Iran/clade 2344/2017(H5N8)), respectively.

Immunizations were administered subcutaneously in two doses with 14-days intervals. Blood samples were collected weekly from Day 0 to Day 56 post-immunization, centrifuged to isolate serum, and stored at -20°C for subsequent immunological analysis.

Table 1: Experimental groups and antigen administration

Groups	Antigen Administered	Adjuvant Type	Dose (µg/chick)	Volume (µl)
cHA H9/H5-Alum	cHA H9/H5	Alum	30	200
cHA H9/H5- Freund's	cHA H9/H5	Complete/Incomplete Freund's	30	200
H5N8	H5N8 inactivated antigen	ISA71 Seppic®	300 HA/U	200
H9N2	H9N2 inactivated antigen	ISA71 Seppic®	300 HA/U	200
Mock	PBS	None	-	200

2.4. Evaluation of Humoral Immune Response

2.4.1. Hemagglutination Inhibition (HI) Assay Protocol

Antibody responses following immunization were evaluated using hemagglutination inhibition (HI) assays with H9N2 2019 and H9N2 ARAK/2009 and H5N8 viral antigens.

For the HI assay, serum samples were subjected to two-fold serial dilutions incubated with 4 HA units of viral antigen at 37°C for 30 minutes, followed by the addition of 1% chicken red blood cells (RBCs). After 30-minute incubation at room temperature, HI titers were determined as the reciprocal of the highest serum dilution that completely inhibited hemagglutination and converted into log₂ values.

2.4.2. Statistical Analysis

Statistical analysis was performed using GraphPad Prism software version 9.0 (GraphPad Software, San Diego, CA, USA). Data were expressed as mean ± standard deviation (SD). Group comparisons were performed using one-way ANOVA followed by Tukey's post hoc test to account for multiple comparisons. A p-value of < 0.05 was considered statistically significant. All analyses were conducted in accordance with established guidelines to ensure the validity and reproducibility of results.

3. Results

The second spherical head of HA in different influenza virus subgroups shows limited homology, while HA2 ecto-domains among various subtypes are highly conserved. The amino acid homology level between HA2 regions of H5 or H9 viruses was about 95% within the same subgroup, but only 64 to 65% identity was observed between H9 and H5 viruses. To develop a multi-subtype vaccine through reverse genetics, we produced a chimeric (cHA) H9/H5 construct, including HA2 regions from A/chicken/Iran/B308B/2004 (H9N2) and A/Poultry/Iran/clade 2344/2018 (H5N8)

The retrieved sequences were evaluated using the VaxiJen server to identify the strongest antigenic region. Prediction scores for hemagglutinin proteins H9N2 and H5N8 were 0.4333 and 0.5202, respectively, higher than the threshold of 0.4. Epitope scores and chemical properties were evaluated, selecting epitopes E1 to E8 (three from H9N2 and five from H5N8), with two glycines inserted between them to create a linear structure. The final sequence, consisting of 296 amino acids, with an isoelectric point of approximately 7 and an estimated molecular weight of 32.46 kDa, indicating good antigenic potential. The protein is non-allergenic with an antigenicity score of 0.4828 (Figure 1, Table 2) .




Figure 1: (A) The primary structure of the recombinant protein, highlighting the selected epitopes from H9N2 and H5N8 strains. (B) The predicted secondary and tertiary structures of the designed recombinant protein, showcasing the spatial arrangement of the selected epitopes. (C) The three-

dimensional model of the recombinant protein, illustrating the overall folding and potential antigenic sites.

Table 2: Physicochemical properties of the designed recombinant protein

Molecular Weight	Instability Index	Aliphatic Index	Theoretical pI	No. of Amino acids	Total Number of Atoms	Extinction Co-Efficient	Gravy
32465.09	17.93	71.45	7.19	296	4499	64080	-0.568

The cHA H9/H5 recombinant protein gene was cloned into the pET-41a (+) vector, which includes an N-terminal His-tag sequence, between *XhoI* and *NcoI* sites after codon optimization. The recombinant plasmid was transformed into *E. coli* BL21 (DE3) cells for protein expression. After expression induction, SDS-PAGE and Western blot analyses using anti-His antibodies confirmed successful expression of the target protein, with an expected molecular weight of approximately 32 kDa (Figure 2). Densitometry analysis confirmed that the purity exceeded 95%. The yield of purified cHA H9/H5 protein was approximately 22 mg per liter of bacterial culture, determined by Bradford protein assay.




Figure 2: Analysis of Protein Expression, Purification and Western Blot Analysis. A: SDS-PAGE analysis of protein expression, showing a prominent band at approximately 32 kDa.

Lane M: Molecular weight markers, Lane 1: non-induced sample, Lanes 2–4: Recombinant protein expression at 2, 3, and 4 hour's post-induction, respectively. B: Purification of the expressed protein using Ni-NTA affinity chromatography, as indicated by the SDS-PAGE analysis (Lanes 1-2: Sequential elution of purified recombinant protein). C: Western blot analysis confirming the presence of the target protein at approximately 32 kDa.

To determine the immunogenicity and protective efficacy of the cHA H9/H5 antigen, we immunized groups of specific pathogen-free (SPF) chickens with antigen, formulated with different adjuvants. Subsequently, the humoral immune response was measured by hemagglutination inhibition (HI) assay. It revealed distinct immune responses in chickens immunized with the recombinant chimeric cHA H9/H5 antigen, as well as H9N2 and H5N8 vaccine antigens (Figure 3).

Chickens receiving the cHA H9/H5 antigen formulated with Alum or Freund's adjuvants exhibited the elevated HI geometric mean titers (GMT) against the homologous cHA H9/H5 antigen, indicating robust immunogenicity. Notably, when cHA H9/H5 antigen formulated with Freund's adjuvant, it demonstrated elevated cross-reactive responses to the heterologous H9N2 antigen ($p < 0.05$). Commercial vaccine groups who immunized with inactivated H5N8 and H9N2 antigens showed antigen-specific reactivity, the H5N8 vaccine induced strong HI responses to both cHA H9/H5 and H5N8 antigens, while the H9N2 vaccine elicited high titers against cHA H9/H5 and H9N2 antigens but lower reactivity to H5N8. These findings underscore the enhanced immunogenicity of adjuvanted cHA H9/H5 and highlight cross-reactivity patterns between the chimeric antigen and conventional H5N8/H9N2 vaccine strains.




Figure 3: Hemagglutination Inhibition (HI) geometric mean titers (GMT) against different influenza virus strains. The y-axis represents SN GMT in log base 2 ($10\log_2$) and the x-axis lists different vaccine groups: cHA H9/H5, H9N2, and H5N8.

4. Discussion

The development of Hemagglutinin (HA) protein chimeric vaccines represents a transformative approach to improving influenza immunization strategies. Traditional influenza vaccines target the highly variable HA head domain, necessitating annual updates due to antigenic drift or shift. Chimeric HA vaccines, however, incorporate conserved regions (e.g., the HA stalk) fused with exotic head domains to elicit broad, cross-reactive immunity against diverse influenza strains, including pandemic variants. Preclinical studies demonstrate that such vaccines induce antibodies targeting conserved epitopes, enhancing protection against heterosubtypic viruses and reducing the need for frequent reformulation(14). This strategy aligns with the WHO's priority for universal influenza vaccines to mitigate global health burdens. By leveraging conserved antigenic regions, chimeric HA vaccines offer a promising path toward durable, pan-influenza immunity.

In this study, we aimed to elicit a broad and robust immune response by targeting multiple conserved epitopes from different influenza strains. This innovative approach not only enhances cross-protection against diverse influenza subtypes but also paves the way for the development of a universal influenza vaccine, which is crucial for improving global pandemic preparedness and reducing the impact of influenza outbreaks.

In comparison to our study, one study utilized an immunoinformatic approach to create a vaccine based on conserved epitopes from hemagglutinin, neuraminidase, and matrix proteins. This vaccine, designed with specific linkers and evaluated for immunogenicity using computational frameworks, demonstrated promising immune responses in terms of IgG, T Helper 1 cells, and interferon gamma levels(15) . Another research effort focused on creating a universal T-cell vaccine using a recombinant vaccinia virus expressing multiple epitopes from influenza virus proteins such as NP, M1, NS1, PB1, and PA, resulting in increased influenza virus-specific IFN- γ secreting splenocytes and decreased viral load in vaccinated mice (16). Additionally, a study combining the M2e peptide with stalk HA epitopes of influenza A virus observed enhanced immunogenicity and protective properties, with recombinant proteins providing broad protection against various influenza viruses, including A/H3N2, A/H2N2, and A/H5N1(17).

It's evident that all studies emphasize the use of conserved epitopes and strategic linkers to enhance stability and immune responses. Use of EAAAK, KK, AAY, and GPGPG linkers for epitope stability and immune activity is consistent with similar approaches in these studies. Additionally, the confirmation of protein expression through Western blotting and SDS-PAGE staining aligns with common practices in the field.

The study successfully cloned the cHA H9/H5 recombinant protein gene into the pET-41a (+) vector, which was then transformed into E. coli BL21 (DE3) cells for expression, confirmed by SDS-PAGE and Western blot analyses. The protein was purified using Ni-NTA agarose Affinity Chromatography, achieving over 95% purity with a yield of approximately 22 mg per liter of bacterial culture. These results indicate a robust and scalable method for producing cHA H9/H5 recombinant protein for further immunogenicity and functional studies, aligning with other research utilizing E. coli for high-yield expression of recombinant influenza proteins, such as those that fused the globular head domain of HA with flagellin for robust antibody responses in mice (18). Similarly, the study aligns with approaches highlighted in reviews of protein subunit vaccines targeting H1N1/09, emphasizing antigen selection and expression

systems (19). Additionally, the study's robust method for producing recombinant protein could be further evaluated for immunogenicity and safety, as demonstrated in comparisons of quadrivalent recombinant and inactivated influenza vaccines (20). Overall, the results provide a scalable method for producing cHA H9/H5 recombinant protein, consistent with studies emphasizing high-yield bacterial expression and purification techniques.

Our study demonstrates that the cHA H9/H5 vaccine formulations with ALUM and Freund's adjuvants significantly enhance hemagglutination inhibition (HI) titers against multiple influenza strains, highlighting the superior efficacy of these adjuvants in boosting antibody production. This finding aligns with other studies, such as the one by Kim et al., which showed that a chimeric H9/H5N2 vaccine induced robust HI titers and provided protection against HPAI H5N8 viruses in chickens(21) .

Overall, our results underscore the potential of the cHA H9/H5 antigen combined with adjuvant formulations in eliciting robust immune protection against a range of influenza virus strains, consistent with similar research efforts in the field.

Acknowledgment

This research constitutes part of the doctoral dissertation of the first author and was made possible by the generous support of the Karaj Branch, Islamic Azad University, Iran, and Razi Vaccine and Serum Research Institute (Marand Branch). We sincerely thank all the dedicated staff of the Marand branch of the Razi Vaccine and Serum Research Institute for their invaluable efforts and contributions. Their collective expertise, hard work, and commitment have been vital to the success of this project.

Authors' Contribution

Concept and design of the study: M.T, M.G

Acquisition of data: Z.B.

Analysis and interpretation of data: Z.B, M.T

Drafting of the manuscript: Z.B, M.T

Statistical analysis: B.H

Administrative, technical, and material support: M,T, Z.B.

Ethics

The procedure of this trial was approved by the Ethics Committee of the Karaj Branch of Islamic Azad University (Approval No: IR.IAU.K.REC.1398.026).

Conflict of Interest

The authors declare that they have no conflict of interest.

Data Availability

Data will be available upon request to the corresponding author.

References

1. Krammer F, Smith GJD, Fouchier RAM, et al. Influenza. *Nat Rev Dis Primers*. 2018;4(1):3. doi:10.1038/s41572-018-0002-y
2. Yuen KY, Chan PK, Peiris M, et al. Clinical features and rapid viral diagnosis of human disease associated with avian influenza A H5N1 virus. *Lancet*. 1998;351(9101):467-471. doi:10.1016/s0140-6736(98)01182-9
3. Lee CW, Saif YM. Avian influenza virus. *Comp Immunol Microbiol Infect Dis*. 2009;32(4):301-310. doi:10.1016/j.cimid.2008.01.007
4. Peacock THP, James J, Sealy JE, Iqbal M. A Global Perspective on H9N2 Avian Influenza Virus. *Viruses*. 2019;11(7):620. doi:10.3390/v11070620
5. Lee DH, Torchetti MK, Winker K, Ip HS, Song CS, Swayne DE. Intercontinental Spread of Asian-Origin H5N8 to North America through Beringia by Migratory Birds. *J Virol*. 2015;89(12):6521-6524. doi:10.1128/JVI.00728-15

6. Swayne DE. Impact of Vaccines and Vaccination on Global Control of Avian Influenza. *Avian Diseases*. 2012; 56 (4): 818–28. doi:10.1637/10183-041012-Review.1
7. World Health Organization. Antigenic and genetic characteristics of zoonotic influenza viruses and development of candidate vaccine viruses for pandemic preparedness. 2022. <https://www.who.int/publications/m/item/zoonotic-influenza-update>
8. WHO Manual on Animal Influenza Diagnosis and Surveillance. Hemagglutination inhibition (HI) test for avian influenza. 2022. <https://www.fao.org/3/a1521e/a1521e.pdf>
9. Islam MSB, Miah M, Hossain ME, Kibria KMK. A conserved multi-epitope-based vaccine designed by targeting hemagglutinin protein of highly pathogenic avian H5 influenza viruses. *3 Biotech*. 2020;10(12):546. doi:10.1007/s13205-020-02544-3
10. Matrosovich M, Zhou N, Kawaoka Y, Webster R. The surface glycoproteins of H5 influenza viruses isolated from humans, chickens, and wild aquatic birds have distinguishable properties. *J Virol*. 1999; 73(2):1146-1155. doi:10.1128/JVI.73.2.1146-1155.1999
11. Schrauwen EJ, Fouchier RA. Host adaptation and transmission of influenza A viruses in mammals. *Emerg Microbes Infect*. 2014;3(2):e9. doi:10.1038/emi.2014.9
12. Capua I, Marangon S. Control and prevention of avian influenza in an evolving scenario. *Vaccine*. 2007;25(30):5645-5652. doi:10.1016/j.vaccine.2006.10.053
13. Bozorgkhoo z, Hemmati B, Taghizadeh M, Tebianian M, Ghaderi M. Bioinformatic Anaysis of Multi-epitope Protein Designed Based on Highly Conserved Regions of Hemagglutinin Protein of H9N2 and H5N8 Influenza Virus Strains %J Veterinary Research & Biological Products. 2023;36(4):2-13. Doi: 10.22092/VJ.2022.360912.2034
14. Krammer F, Palese P. Advances in the development of influenza virus vaccines. *Nat Rev Drug Discov*. 2015;14(3):167-182. doi:10.1038/nrd4529
15. Maleki A, Russo G, Parasiliti Palumbo GA, Pappalardo FJBb. In silico design of recombinant multi-epitope vaccine against influenza A virus. *BMC bioinformatics* 2021; 22(Suppl 14):617.
16. Goodman AG, Heinen PP, Guerra S, Vijayan A, Sorzano COS, Gomez CE, et al. A human multi-epitope recombinant vaccinia virus as a universal T cell vaccine candidate against influenza virus. *PloS one*. 2011;6(10):e25938. doi:10.1371/journal.pone.0025938
17. Tsybalova LM, Stepanova LA, Shuklina MA, et al. Combination of M2e peptide with stalk HA epitopes of influenza A virus enhances protective properties of recombinant vaccine. *PLoS One*. 2018; 13(8):e0201429. doi:10.1371/journal.pone.0201429
18. Song L, Nakaar V, Kavita U, et al. Efficacious recombinant influenza vaccines produced by high yield bacterial expression: a solution to global pandemic and seasonal needs. *PLoS One*. 2008;3(5):e2257. doi:10.1371/journal.pone.0002257
19. Zhang Y, Gao J, Xu W, et al. Advances in protein subunit vaccines against H1N1/09 influenza. *Front Immunol*. 2024;15:1499754. doi:10.3389/fimmu.2024.1499754

- 359 20. Dunkle LM, Izikson R, Patriarca PA, Goldenthal KL, Muse D, Cox MMJ. Randomized
360 Comparison of Immunogenicity and Safety of Quadrivalent Recombinant Versus Inactivated
361 Influenza Vaccine in Healthy Adults 18-49 Years of Age. *J Infect Dis.* 2017;216(10):1219-
362 1226. doi:10.1093/infdis/jix478
- 363 21. Kim SM, Kim YI, Park SJ, et al. Vaccine Efficacy of Inactivated, Chimeric Hemagglutinin
364 H9/H5N2 Avian Influenza Virus and Its Suitability for the Marker Vaccine Strategy. *J Virol.*
365 2017;91(6):e01693-16. doi:10.1128/JVI.01693-16

preprint