

# Molecular Characterization and Phylogenetic analysis of Deformed Wing Virus in Honey Bees (*Apis Mellifera*), North of Iran

3

4 **Fahimeh Jamiri<sup>1</sup>, Zahra Ziafati Kafi<sup>1</sup>, Shabnam Jozghasemi<sup>2</sup>, Soroush Sarmadi<sup>1</sup>,**  
5 **Alireza Bakhshi<sup>1</sup>, Nazanin Sarvian<sup>1</sup>, Arash Ghalyanchilangeroudi<sup>1\*</sup>**

6 1- Department of Microbiology and Immunology, Faculty of Veterinary Medicine,  
7 University of Tehran, Tehran, Iran

8 2- Head of Laboratory, Alborz Province Veterinary Organization, Alborz, Iran

9

## 10 **\*Corresponding**

11 Arash Ghalyanchi Langeroudi

12 Department of microbiology and immunology, Faculty of veterinary medicine,  
13 University of Tehran, Tehran, Iran

14 Email: arashghalyanchi@gmail.com

15 Phone: 02161117154

16 Fax: +98-21-66933222

17

18

## 19 **Abstract**

20 Deformed wing virus (DWV) is one of the major bee viruses which can be transmitted by the *Varroa*  
21 *destructor* mite and is a key contributor to honey bee colony collapse. To date, three main variants of the  
22 Deformed wing virus (DWV-A, DWV-B, DWV-C) have been described. In the present study, bee samples  
23 showing signs of DWV infection and *Varroa* infestation were collected from two different farms in Alborz  
24 province, northern Iran. The samples were screened for the presence of DWV genome and also other bee  
25 viral pathogens including Chronic bee paralysis virus, Acute bee paralysis virus, Kashmir bee virus and sac  
26 brood virus by using PCR assay. PCR results indicated that among collected samples, two samples were  
27 positive for DWV and all samples were negative for other viral disease. Phylogenetic analysis revealed that  
28 the detected DWVs clustered within the DWV-A genotype. Homology analysis of isolate UT-SHN07

showed approximately 99% similarity to isolate PA and DWV-A-OR-OCT2018-EVR23 from USA, as well as to DWV-Iraq-2023 and the reference sequence NC004830 from Iraq and Italy respectively. For the other isolate, UT-SHN62, homology analysis showed 100% similarity with isolate PA isolated from USA and 99% similarity with DWV-A-OR-OCT2018-EVR23, DWV-Iraq-2023, NC004830 and Gotland-2 sequences from the USA, Iraq, Italy and Sweden respectively. Iran is ranked among top 10 honey-producing countries, and continuous monitoring of circulating viruses and *Varroa* mites (key contributors to colony losses) is essential. Further molecular and phylogenetic studies are recommended to better understand the current status of honey bee viral disease in Iran.

## Keywords

Honeybee, Deformed wing virus, Colony loss, Phylogeny, Homology

## 1. Introduction

Since the beginning of 21<sup>st</sup> century, substantial honey bee losses have been reported across the northern hemisphere (1). It's estimated that nearly one-third of the typical western diet depends on bee pollination. The honey bee, *Apis mellifera*, is recognized as the primary insect responsible for pollinating a wide range of crops, including fruits, nuts, vegetables and oilseeds, with an estimated annual added market value exceeding 15 billion dollars (2, 3). In addition to pollination, honey bees contribute to the economy by producing various hive products such as honey, beeswax, propolis, royal jelly and others (4). Over the past 15 years, severe winter colony losses have been reported from different regions worldwide raising growing concern about the health status of bee colonies. Nevertheless, over the past 60 years, the number of managed bee colonies has increased all over the world (5). Like many other species, honey bees are susceptible to a variety of pathogens that can cause considerable colony loss. Among these, viruses are considered one of major threats to honey bee health and survival (4).

Viruses that infect honey bees are primarily classified within the order *Picornavirales*, and are positive sense single-stranded RNA viruses (+ssRNA). Common bee viruses include members of the Dicistroviruses such as Israeli acute paralysis virus (IAPV), Kashmir bee virus (KBV), Acute bee paralysis virus (ABPV) and Black queen cell virus (BQCV); the Iflaviruses such as Deformed wing virus (DWV), Kakugo virus, *Varroa destructor* virus-1 (also known as DWV-B), Sacbrood virus (SBV), and Slow bee paralysis virus (SBPV) and a taxonomically unclassified virus Chronic bee paralysis virus (CBPV) and Lake Sinai viruses (LSVs) (6).

Deformed wing virus (DWV), transmitted by the ectoparasite mite *Varroa destructor*, causes wing deformities in honeybees and is considered one of the main viruses associated with the colony collapse, premature death and reduced performance in asymptomatic bees (7, 8). Transmission of DWV by *V.destructor* to pupae can result in clinical symptoms such as pupal death, the emergence of adult bees with deformed wings, bloated and shortened abdomen, and discoloration. However, in the absence of

1 *V.destructor* mites, DWV infection appears to cause no visible symptoms or detectable impact on host  
2 fitness (7). Although infection with DWV in honey bees is strongly associated with *V.destructor* mite  
3 infestation, it can also exist as a covert (symptomless) infection in colonies where the mite is absent (9).  
4 DWV is a picorna-like virus classified in Iflaviridae family. Its genome encodes a poly protein that is  
5 subsequently cleaved by viral and/or cellular enzyme into structural and non-structural proteins required  
6 for the viral replication cycle (10). Mature Virions of Iflaviridae family, are spherical without envelope and  
7 measuring 22-30 nm in diameter. They contain three major structural proteins (VP1, VP2 and VP3) and a  
8 smaller capsid protein VP4 which found in some species (11).

9 Three main variants of DWV have been described: DWV-A, DWV-B and DWV-C, with DWV-A and  
10 DWV-B being the most prevalent globally in honey bee colonies (12). However, more recently a newly  
11 identified virus named Egypt bee virus (EBV) has been proposed as a potential fourth variant of DWV.  
12 EBV appears to be more closely related to DWV-C than to either DWV-A and DWV-B (13). The DWV-A  
13 variant is associated with colony loss. However, the effects of DWV-B on colony health remain a subject  
14 of debate. While DWV-B has been detected in healthy colonies, it has also been suggested that this variant  
15 may offer protective benefits to colonies through a mechanism known as superinfection exclusion (14).

16 According to FAOSTAT 2023 report, Iran ranked third among the top ten natural honey-producing  
17 countries, with an annual production of approximately 100000 tons of honey (15). It has been reported that  
18 in Iran, pesticides and diseases, particularly *Varroa* mite infestations, are among the primary causes of  
19 colony losses and reduction in honey production. Consequently, viruses transmitted by this mite are  
20 believed to contribute significantly to these losses. However, limited information is available regarding  
21 viral infections and their relationship with *Varroa* mites and colony losses in Iran (16). DWV was first  
22 reported in Iran by Ghorani et al in 2017 and since then, several studies have confirmed the presence and  
23 circulation of this virus among bee colonies and mites in the country (16-20).

24 Since Iran ranks among the top ten honey-producing countries, monitoring the presence and circulation of  
25 viruses and mites in bee colonies is of great importance, as these factors can significantly affect colony  
26 health and, consequently performance. In this study, we investigated the simultaneous presence of  
27 Deformed Wing Virus (DWV) and *Varroa* mites in honey bee colonies located in Alborz Province, northern  
28 Iran.

## 29 30 **2. Material and methods**

### 31 **2.1 Sample collection**

32 Samples were collected from honey bee colonies showing signs of wing deformities at two separate farms  
33 located in Alborz Province, northern Iran. A total of 24 colonies were sampled from first farm (farm A) and  
34 15 colonies from the second farm (farm B). The sampled colonies showed darkened brood cells, deformed  
35 and atrophied wings, and, in some cases, symptoms suggestive of wing paralysis. Samples from each farm  
36 were first screened for *Varroa destructor* infestation, then pooled and homogenized. All specimens were

submitted to the Virology Laboratory of the Faculty of Veterinary Medicine, University of Tehran, and stored at -20 °C until further processing.

## **2.2 RNA extraction and RT-PCR**

Samples from each colony were pooled to generate a single representative sample per farm. The pooled samples were then crushed and homogenized using a ceramic mortar with diethylpyrocarbonate (DEPC) treated water. The homogenates were centrifuged in 20,000 × g for 1 minute, and 200 µl of the supernatant was used for total RNA extraction. RNA extraction was performed using Sinapure™ RNA extraction kit according to the manufacturer's instructions. The extracted RNA was eluted into 50 µl of RNAase free water and stored at -20 °C until synthesis of cDNA. For cDNA synthesis, 1µl of random hexamer primer was added to 10µl of total RNA. The mixture was incubated at 70 °C for 5 minutes, followed by immediate cooling at 5°C for 1 minute. In the next step, 9µl of cDNA master mix containing 1µl of MMLV reverse transcriptase enzyme and 4µl of 5X RT-buffer (Yekta Tajhiz Azma, Iran), 1µl of dNTPs (SinaClone, Iran) and 2µl of DEPC-treated water were added resulting in a final reaction volume of 20µl. For the second step, the mixture kept in 25 °C for 5 minutes, after than incubated in 42 °C for 1 hour and then heated to 85 °C for 5 minutes and then cooled to 5 °C for 1 minute.

To detect DWV, a pair of primers targeting the polyprotein coding region were used: DWVF: 5'-CTTACTCTGCCGTCGCCCA-3' and DWVR: 5'-CCGTTAGGAACTCATTATCG-3'). These primers amplify a 160 bp fragment of DWV genome. The PCR conditions were as follows: initial denaturation at 94 °C for 1 minute; 35 cycles of denaturation at 94 °C for 1 minute, annealing at 56 °C for 1 minute, and extension at 72 °C for 1 minute; followed by a final extension at 72 °C for 10 minutes. To evaluate PCR products, 5 µL of each reaction was loaded onto a 1.5% agarose gel, stained with ethidium bromide, and visualized under ultraviolet (UV) light using a transilluminator.

## **2.3 Sequencing and phylogenetic analysis**

Positive samples were submitted for Sanger sequencing by Codon Genetic Group (Tehran, Iran). The quality of the obtained sequences was evaluated using NCBI BLAST and FinchTV software (version 1.4.0). Phylogenetic analysis was performed using MEGA 7 software with the Maximum Likelihood method based on the General Time Reversible (GTR) model with 1000 bootstrap replicates. For this analysis, the sequences obtained in the present study were compared with previously submitted DWV sequences available in the NCBI database. The sequences of the current isolates were submitted to GenBank and available under accession numbers: PV975901 (isolate UT-SHN07) and PV975902 (isolate UT-SHN62).

## **3. Results**

Phylogenetic analysis of the detected DWV isolates revealed that they belong to Deformed wing virus variant A (DWV-A) (Figure 1). Homology analysis showed that the sequence UT-SHN07 from the present study share approximately 99% nucleotide identity with sequences AY292384 and OR361536 from the

USA OR130297 and NC004830 from Iraq and Italy respectively. Results of homology analysis for other isolate in this study, UT-SHN62, indicated 100% similarity between the sequence of this isolate and sequence AY292384 isolated from USA. The sequence of UT-SHN62 has also showed almost 99% similarity with sequences OR361536 (USA), OR130297 (Iraq), NC004830 (Italy) and, MZ867711 (Sweden) (table 1). Investigation of Varroa infestation demonstrated that, 13 out of 24 colonies (54%) from farm A and 3 out of 15 (20%) from farm B, were infested with mites.

#### 4. Discussion

The global population of honeybees (*Apis mellifera*) is under pressure due to habitat loss, environmental stress, and specially pathogens such as viruses that can cause lethal epidemics (21). Honeybees play a vital role in pollination and contribute billions of dollars in added value to agriculture (22). Although many viral infections can be asymptomatic, they are of great concern as they can harm honeybees at various developmental stages, including egg, larva, pupa, adult, worker, drone, and queen (22). Among these viruses, Deformed Wing Virus (DWV), along with its vector, the Varroa destructor mite, appears to be a major global threat to honeybee populations (21).

The RT-PCR method developed to amplify a specific nucleotide sequence located in virus genome, present in sample is considered as valuable tool for diagnosis of viral disease in honey bees as it is fast, reliable and specific (especially when target-specific primers are used) (17). In addition to molecular detection, Phylogenetic analysis serves as a useful approach for defining the diversity and differences between nucleotide sequences of various virus strains. These differences are affected through different ways including the replication cycles, the accuracy of replication and the phenotypic effects of mutations. The quantity of these variation is evaluated by phylogenetic algorithms in order to define the probable genetic relationships between isolates (23).

Colony losses have been associated with infestation by an exotic mite, *Varroa destructor*, which feeds on honey bee hemolymph and can harbor different honeybee viral pathogens specially deformed wing virus. Severe DWV infection can result in pupal death, wing deformities, shortened abdomen and cuticle discoloration in adult bees, which often die shortly after pupation, causing colony, collapse (21). In colonies that don't show the symptoms of DWV disease, *Varroa destructor* mites are either absent or present in low levels. The severity of DWV infection has been shown to be directly correlated with the prevalence and intensity of *Varroa destructor* infestation within the colony (16).

In Iran different studies have reported the presence and circulation of bee viruses among colonies. In 2017, Ghorani et al examined 89 Iranian honey bee samples showing signs of depopulation, sudden collapse, paralysis or dark coloring. Using RT-PCR method with DWV specific primers targeting the structural protein encoding gene, they found that 16 samples (17.97%) were positive for DWV making it the most prevalent virus detected in this study (17). In 2018, Moharrami et al screened 156 apiaries for DWV infection using RT-PCR method with specific primers targeting a 435 bp fragment of the polyprotein

1 encoding gene. DWV was detected 34 apiaries (21.8%) with 21.8% of adult bees and 20.51% of pupae  
2 testing positive (24). In another study by Moharrami et al in 2018, analyzed 160 adult bee samples from 23  
3 provinces, detecting DWV in 34 samples (21.8%) ranking it as the second most prevalent virus (20). In a  
4 more recent study by same group (2023), 45 samples were tested for six major bee viruses with emerging  
5 as the most prevalent, present in 37.7% of samples (19). Additionally, Shojaei et al, screened 30 apiaries  
6 using RT-PCR with primers targeting partial VP2, VP4 and VP1 and partial RNA helicase gene. Their  
7 findings showed DWV infection in 36.6% of apiaries were infected with DWV including 8, 0, and 3 positive  
8 samples on capped larvae and workers, and Varroa mites respectively (25). All together, these studies  
9 suggest that DWV is one of the most prevalent honey bee viruses, currently circulating among honeybees  
10 in Iran.

11 In addition to Iran, neighboring and nearby countries have also reported the presence of DWV in their  
12 honey bee colonies. In a study conducted by ÇAĞIRGAN et al in 2021, a total of 111 samples were  
13 collected from seven provinces in Turkey to screen for seven major honey bee viruses. Among the detected  
14 viruses, 22 out of 111 samples tested positive for DWV making it the most prevalent virus in this study  
15 (19.8%) (26). In another study by Karapinar et al, samples were collected from 26 hives, of which 18 hives  
16 (69.23%) were positive for DWV infection (27). Haddad et al, investigated the distribution of DWV in  
17 Middle Eastern and North African (MENA) countries. For this propose, 111 adult worker bee samples were  
18 collected from 12 countries including Lebanon, Syria, Iraq, Palestine, Jordan, Egypt, Libya, Tunisia,  
19 Algeria, Morocco, Yemen and Sudan. According to the results, the highest prevalence was observed in  
20 Syria and Lebanon with lower rates reported in Palestine, Jordan and Iraq (28).

21 In the present study, homology analysis revealed 100% and 99 % sequence similarity of isolates UT-SHN62  
22 and UT-SHN07, respectively, with isolate PA/USA isolate reported from the United States. The genome of  
23 this isolate (PA/USA) was studied completely and phylogenetic analysis revealed that it clustered in  
24 genotype A of DWV (29).

25 Besides their vital role in ecosystem, honey bees contribute significantly to the economy through products  
26 like honey, beeswax, propolis and royal jelly. Viruses, particularly DWV, pose a serious threat to bee health  
27 and colony sustainability. Present study is performed for detection and phylogeny of DWV isolated from 2  
28 different farms in Alborz province in the north of Iran. According to this study, both isolates were clustered  
29 in genotype A of deformed wing viruses. Monitoring of common circulating viruses is necessary for finding  
30 new ways in order to prevent and control viral bee disease and subsequently prevent colony losses and  
31 decrease of production. Constant monitoring can help to improve the quality of production and increasing  
32 the exportation potential of honey bee products. Further studies are recommended for genotyping of  
33 circulating deformed wing viruses in Iran.

34

35

36

## Conflict of Interest

The authors declare no conflict of interest.

## Author's contribution

Conceptualization, Methodology and Supervision: A. G.

Sample collection: S. J.

Formal Analysis and Software: Z. Z K, S. S.

Original Draft and Writing: F. J, Z. ZK, S. S, A. B, N. S.

Review and Editing: A. G, S. J.

## Ethics committee Approval

Our study did not involve any invasive procedures on animals therefore, an ethics committee approval is unnecessary.

## Data Availability

The data that supporting the findings of this study are available upon request from the corresponding author.

## References

1. Coulon M, Dalmon A, Di Prisco G, Prado A, Arban F, Dubois E, et al. Interactions between thiamethoxam and deformed wing virus can drastically impair flight behavior of honey bees. *Frontiers in microbiology*. 2020;11:766.
2. Chen Y, Zhao Y, Hammond J, Hsu H-t, Evans J, Feldlaufer M. Multiple virus infections in the honey bee and genome divergence of honey bee viruses. *Journal of Invertebrate Pathology*. 2004;87(2):84-93.
3. Brutscher LM, McMenamin AJ, Flenniken ML. The buzz about honey bee viruses. *PLoS Pathogens*. 2016;12(8):e1005757.
4. Chen YP, Siede R. Honey Bee Viruses. *Advances in Virus Research*. 70: Academic Press; 2007. p. 33-80.
5. Gisder S, Genersch E. Special Issue: Honey Bee Viruses. *Viruses*. 2015;7(10):5603-8.
6. McMenamin AJ, Flenniken ML. Recently identified bee viruses and their impact on bee pollinators. *Current Opinion in Insect Science*. 2018;26:120-9.
7. de Miranda JR, Genersch E. Deformed wing virus. *Journal of Invertebrate Pathology*. 2010;103:S48-S61.
8. Desai S, Eu YJ, Whyard S, Currie R. Reduction in deformed wing virus infection in larval and adult honey bees (*Apis mellifera* L.) by double-stranded RNA ingestion. *Insect molecular biology*. 2012;21(4):446-55.



9. Gisder S, Aumeier P, Genersch E. Deformed wing virus: replication and viral load in mites (*Varroa destructor*). *Journal of General Virology*. 2009;90(2):463-7.
10. Gusachenko ON, Woodford L, Balbirnie-Cumming K, Campbell EM, Christie CR, Bowman AS, et al. Green Bees: Reverse Genetic Analysis of Deformed Wing Virus Transmission, Replication, and Tropism. *Viruses*. 2020;12(5).
11. Valles S, Chen Y, Firth A, Guérin DA, Hashimoto Y, Herrero S, et al. ICTV virus taxonomy profile: Iflaviridae. *Journal of General Virology*. 2017;98(4):527-8.
12. Brettell LE, Schroeder DC, Martin SJ. RNAseq of Deformed Wing Virus and Other Honey Bee-Associated Viruses in Eight Insect Taxa with or without *Varroa* Infestation. *Viruses*. 2020;12(11):1229.
13. de Miranda JR, Brettell LE, Chejanovsky N, Childers AK, Dalmon A, Deboutte W, et al. Cold case: The disappearance of Egypt bee virus, a fourth distinct master strain of deformed wing virus linked to honeybee mortality in 1970's Egypt. *Virology journal*. 2022;19(1):12.
14. Kevill JL, Stainton KC, Schroeder DC, Martin SJ. Deformed wing virus variant shift from 2010 to 2016 in managed and feral UK honey bee colonies. *Archives of Virology*. 2021;166(10):2693-702.
15. FAO. Production of natural honey in the world 2022 [Available from: [https://www.fao.org/faostat/en/#rankings/countries\\_by\\_commodity](https://www.fao.org/faostat/en/#rankings/countries_by_commodity)].
16. Sabahi Q, Morfin N, Nehzati-Paghaleh G, Guzman-Novoa E. Detection and replication of deformed wing virus and black queen cell virus in parasitic mites, *Varroa destructor*, from Iranian honey bee (*Apis mellifera*) colonies. *Journal of Apicultural Research*. 2020;59(2):211-7.
17. Ghorani M, Madadgar O, Langeroudi AG, Rezapanah M, Nabian S, Akbarein H, et al. The first comprehensive molecular detection of six honey bee viruses in Iran in 2015-2016. *Archives of Virology*. 2017;162(8):2287-91.
18. Shojaei A, Khanjani M, Nourian A, Mahmoodi P. Detection and phylogenetic analysis of deformed wing virus (DWV) in honey bees, *Apis mellifera* L. and parasitic mite, *Varroa destructor* Anderson and Trueman (Acari: Varroidae). *Systematic and Applied Acarology*. 2021;26(6):1177-84, 8.
19. Moharrami M, Modirrousta H, Torkaman M, Jalali HR, Ahmadi K, Fallah Mehrabadi MH. Molecular epidemiology of six pathogenic viruses in apiaries of eight provinces of Iran. *Veterinary Research & Biological Products*. 2023;36(2):52-9.
20. Moharrami M, Modirrousta H. Molecular identification of six honeybee viruses in Iranian apiaries. *Archives of Razi Institute*. 2018;73(4):311-8.
21. Skubnik K, Novacek J, Fuzik T, Pridal A, Paxton RJ, Plevka P. Structure of deformed wing virus, a major honey bee pathogen. *Proc Natl Acad Sci U S A*. 2017;114(12):3210-5.
22. Arzumanyan H, Avagyan H, Voskanyan H, Simonyan L, Simonyan J, Semirjyan Z, et al. First molecular detection of the presence of honey bee viruses in insects, *Varroa destructor* mites, and pollinated plants in an isolated region of Armenia. *Vet World*. 2023;16(5):1029-34.
23. Berényi O, Bakonyi T, Derakhshifar I, Köglberger H, Topolska G, Ritter W, et al. Phylogenetic analysis of deformed wing virus genotypes from diverse geographic origins indicates recent global distribution of the virus. *Applied and Environmental Microbiology*. 2007;73(11):3605-11.
24. Moharrami M, Modirrousta H, Torkaman M, Hemmati B, Shojaei M. Identification of deformed wing virus (DWV) in honeybee pupae by RT-PCR. *Veterinary Research & Biological Products*. 2018;31(4):26-33.
25. Shojaei A, Khanjani M, Nourian A, Mahmoodi P. Detection and phylogenetic analysis of deformed wing virus (DWV) in honey bees, *Apis mellifera* L. and parasitic mite, *Varroa destructor* Anderson and Trueman (Acari: Varroidae). *Systematic and Applied Acarology*. 2021;26(6):1177-84.
26. Çağırhan AA, Yazici Z. The prevalence of seven crucial honeybee viruses using multiplex RT-PCR and their phylogenetic analysis. *Turkish Journal of Veterinary & Animal Sciences*. 2021;45(1):44-55.
27. Karapınar Z, Oğuz B, Dinçer E, Öztürk C. Phylogenetic analysis of black queen cell virus and deformed wing virus in honeybee colonies infected by mites in Van, Eastern Turkey. 2018.
28. Haddad NJ, Nouredine A, Al-Shagour B, Loucif-Ayad W, El-Niweiri MA, Anaswah E, et al. Distribution and variability of deformed wing virus of honeybees (*Apis mellifera*) in the Middle East and North Africa. *Insect science*. 2017;24(1):103-13.
29. Lanzi G, de Miranda JR, Boniotti MB, Cameron CE, Lavazza A, Capucci L, et al. Molecular and biological characterization of deformed wing virus of honeybees (*Apis mellifera* L.). *J Virol*. 2006;80(10):4998-5009.



1

2

preprint

3 Table 1- Nucleotide sequence variation for deformed wing virus of isolates in present study compared with previous sequences retrieved from  
4 NCBI GeneBank.

|    |                                                                        | 1    | 2    | 3    | 4    | 5    | 6    | 7    | 8    | 9    | 10   | 11   | 12   | 13   | 14   | 15   | 16   | 17   | 18   | 19   | 20   | 21 |
|----|------------------------------------------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|----|
| 1  | UT-SHN07 (PV975901)                                                    |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 2  | UT-SHN62 (PV975902)                                                    | 99.3 |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 3  | PA/USA (AY292384.1)                                                    | 99.3 | 100  |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 4  | DWV-A-OR-OCT2018-EVR23/USA (OR361536.1)                                | 99.3 | 98.6 | 98.6 |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 5  | DWV-Iraq-202/Iraq (OR130297.1)                                         | 98.6 | 99.3 | 99.3 | 98   |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 6  | Deformed wing virus Reference sequence/Italy (NC_004830.2)             | 98.6 | 99.3 | 99.3 | 97.9 | 98.6 |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 7  | Gotland-2/Sweden (MZ867711.1)                                          | 98   | 98.6 | 98.6 | 97.2 | 99.3 | 98.6 |      |      |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 8  | 85-DWV/France (KX373899.2)                                             | 97.9 | 97.2 | 97.2 | 97.2 | 96.5 | 97.2 | 97.2 |      |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 9  | Chilensis_A1/Chilie (JQ413340.1)                                       | 97.2 | 96.5 | 96.5 | 96.5 | 95.7 | 96.4 | 96.5 | 97.2 |      |      |      |      |      |      |      |      |      |      |      |      |    |
| 10 | Warwick-2009/United Kingdom (GU109335.1)                               | 96.5 | 97.2 | 97.2 | 95.8 | 97.9 | 97.2 | 98.6 | 96.5 | 96.5 |      |      |      |      |      |      |      |      |      |      |      |    |
| 11 | Austria 1414/ Austria (KU847397.1)                                     | 96.5 | 97.2 | 97.2 | 95.8 | 96.5 | 97.2 | 97.3 | 95.8 | 95   | 95.8 |      |      |      |      |      |      |      |      |      |      |    |
| 12 | AmE711/USA (KT004425.1)                                                | 95.7 | 95   | 95   | 96.5 | 94.3 | 94.2 | 93.5 | 93.4 | 94.2 | 91.8 | 93.4 |      |      |      |      |      |      |      |      |      |    |
| 13 | Kakugo virus/ japan (AB070959.1)                                       | 94.3 | 95.1 | 95.1 | 93.6 | 94.3 | 95   | 95.1 | 93.6 | 92.7 | 93.5 | 96.5 | 89.4 |      |      |      |      |      |      |      |      |    |
| 14 | Korea1/ South Korea (JX878304.1)                                       | 94.1 | 94.9 | 94.9 | 93.4 | 95.7 | 94.9 | 96.4 | 93.4 | 94.2 | 94.9 | 94.9 | 91   | 92.6 |      |      |      |      |      |      |      |    |
| 15 | Korea2/ South_Korea (JX878305.1)                                       | 93.4 | 94.2 | 94.2 | 92.6 | 95   | 94.1 | 95.7 | 92.6 | 93.5 | 94.1 | 95.7 | 90.3 | 93.5 | 94.9 |      |      |      |      |      |      |    |
| 16 | DWV-B-MI-OCT2018-EVR22/ USA (OR361558.1)                               | 70.7 | 71.8 | 71.8 | 71.9 | 70.7 | 73.4 | 70.7 | 70.7 | 66   | 70.3 | 72.9 | 68.9 | 66.8 | 65.3 | 67.3 |      |      |      |      |      |    |
| 17 | Leuven-dw1/ Belgium (KX783225.1)                                       | 69.8 | 71   | 71   | 71.1 | 69.8 | 72.6 | 69.8 | 69.8 | 65.2 | 69.5 | 72.1 | 68.1 | 68.7 | 64.5 | 66.4 | 97.9 |      |      |      |      |    |
| 18 | DWV-B-OR-OCT2018-EVR24/ USA (OR361560.1)                               | 69.4 | 70.6 | 70.6 | 70.7 | 69.4 | 72.2 | 69.4 | 69.4 | 64.6 | 69   | 71.6 | 67.5 | 68.2 | 63.9 | 65.9 | 99.3 | 98.6 |      |      |      |    |
| 19 | DWV_B_Netherlands/ Netherlands (MN538209.1)                            | 69.4 | 70.6 | 70.6 | 70.7 | 69.4 | 72.2 | 69.4 | 69.4 | 64.6 | 69   | 71.6 | 67.5 | 68.2 | 63.9 | 65.9 | 99.3 | 98.6 | 100  |      |      |    |
| 20 | VDV-1_Ox/ United Kingdom (KC786222.1)                                  | 69.4 | 70.6 | 70.6 | 70.7 | 69.4 | 72.2 | 69.4 | 69.4 | 64.6 | 69   | 71.6 | 67.5 | 68.2 | 63.9 | 65.9 | 99.3 | 98.6 | 100  | 100  |      |    |
| 21 | DWV-B-ND-JUN2018-EVR03/ USA (OR361540.1)                               | 68.2 | 69.4 | 69.4 | 69.5 | 68.2 | 71   | 68.2 | 68.2 | 63.3 | 67.8 | 70.5 | 66.3 | 64.2 | 62.6 | 64.6 | 98.6 | 96.5 | 97.9 | 97.9 | 97.9 |    |
| 22 | Varroa destructor virus-1 Reference sequence/ Netherland (NC_006494.1) | 67.1 | 68.3 | 68.3 | 68.4 | 69.5 | 70   | 69.5 | 67.1 | 62.1 | 69.1 | 69.4 | 65.2 | 63.1 | 64   | 66   | 98   | 97.2 | 97.3 | 97.3 | 97.3 | 98 |

5

6

7

8

9




Figure 1- Phylogenetic analysis of deformed wing virus. Phylogenetic analysis was performed with MEGA 7 software, using Maximum Likelihood method based on General Time Reversible model with 1000 bootstrap. Black circles represent for isolates in this study. According to this phylogenetic tree, current isolates are clustered with isolates in genotype A of deformed wing virus.